



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) EP 1 352 650 A1

(12) **EUROPEAN PATENT APPLICATION**
published in accordance with Art. 158(3) EPC

(43) Date of publication:
15.10.2003 Bulletin 2003/42

(21) Application number: 01271124.8

(22) Date of filing: 18.12.2001

(51) Int Cl.⁷: **A61K 31/055**, A61K 31/166,
A61K 31/12, A61K 31/18,
A61K 31/167, A61K 31/136,
A61K 31/17, A61K 31/695,
A61K 31/5375, A61K 31/357,
A61K 31/404, A61K 31/44,
A61K 31/498, A61K 31/403,
A61K 31/415, A61K 31/421,
A61K 31/422, A61K 31/433,
A61K 31/428, A61K 31/505,
A61K 31/47

(86) International application number:
PCT/JP01/11084

(87) International publication number:
WO 02/049632 (27.06.2002 Gazette 2002/26)

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 18.12.2000 JP 2000383202

(71) Applicant: Institute of Medicinal Molecular
Design, Inc.
Tokyo 113-0033 (JP)


(72) Inventors:
• MUTO, Susumu
Koganei-shi, Tokyo 184-0003 (JP)

- NAGANO, Tatsuo
Shibuya-ku, Tokyo 151-0072 (JP)
- SAOTOME, Tomomi
Nakano-ku, Tokyo 165-0022 (JP)
- ITAI, Akiko
Bunkyo-ku, Tokyo 113-0033 (JP)

(74) Representative:
Sternagel, Fleischer, Godemeyer & Partner
Patentanwälte
An den Gärten 7
51491 Overath (DE)

(54) **INHIBITORS AGAINST THE PRODUCTION AND RELEASE OF INFLAMMATORY CYTOKINES**

(57) A medicament having inhibitory activity against NF- κ B activation, which comprises a compound represented by the following general formula (I) or a pharmacologically acceptable salt as an active ingredient:



wherein X represents a connecting group, A represents hydrogen atom or acetyl group, E represents an aryl group or a heteroaryl group, and ring Z represents an arene or a heteroarene.

BEST AVAILABLE COPY

EP 1 352 650 A1

Description

Field of Invention

[0001] The present invention relates to pharmaceutical compositions having inhibitory activity against the production and release of inflammatory cytokines such as interleukin (IL)-1, IL-6, IL-8 and tumor necrosis factor (TNF- α), and having inhibitory activity against the activation of NF- κ B.

Background Art

[0002] Inflammation is a basic defense mechanism to various infestations, where inflammatory cytokine such as interleukin (IL)-1 and TNF- α (tumor necrosis factor) are known to play important roles. Due to the progress of gene analysis of inflammatory cytokines and inflammatory cell adhesion factors, it has been revealed that these cytokines are controlled by a common transcription factor (also called as transcription regulatory factor). This transcription factor is a protein called as NF- κ B (also described as NF- κ B, Clark B. D., et al., Nucl. Acids Res., 14, 7898(1984); Nedospasov S. A., et al., Cold Spring Harb. Symp. Quant. Biol., 51, 611(1986)).

[0003] This NF- κ B is a hetero dimer(also called as complex) of p65(also called as Rel A) and p50(also called as NF- κ B-1), usually binds to I- κ B when external stimulation does not exist, and exists in cytoplasm as an inactive type. I- κ B is phosphorylated by various external stimulations such as oxidative stress, cytokine, lipopolysaccharide, virus, UV, free radical, protein kinase C to become ubiquitin, and then decomposed by proteasome (Verma I. M., Stevenson J. K., et al., Genes Dev., 9, 2723-2735(1995)). NF- κ B separated from I- κ B immediately move into nucleus, and plays a role as a transcription factor by binding to promoter region which has recognition sequence of NF- κ B.

[0004] In 1997, phosphoenzyme (called as I- κ B kinase abbreviated as "IKK"), which participates in phosphorylation of I- κ B, was identified (DiDonato J., Hayakawa M., et al., Nature, 388, 548-554(1997); Regnier C. H., Song H. Y., et al., Cell, 90, 373-383(1997)). IKK- α (also called as IKK1) and IKK- β (also called as IKK2) which resemble each other and exist among a class of IKK, and they are known to form a complex to bind directly to I- κ B and phosphorylate I- κ B (Woronicz J. D., et al., Science, 278, 866-869(1997); Zandi, E., et al., Cell, 91, 243-252(1997)).

[0005] Recently, a mechanism except cyclooxygenase inhibition is suggested for aspirin which is a widely used anti-inflammatory agent, which is known to be based on inhibition of NF- κ B activation (Kopp E., et al., Science, 265, 956-959(1994)). Moreover, it was revealed that aspirin regulates release and activation of NF- κ B by binding reversibly to IKK- β which is I- κ B kinase competing with ATP and by inhibiting phosphorylation of I- κ B (Yin M. J., et al., Nature, 396, 77-80(1998)). However, since huge amount of aspirin needs to be administered to sufficiently suppress NF- κ B activation, and as a result, since possibility of side effects such as gastrointestinal disorders by prostaglandin synthesis inhibition and increase of bleeding tendency by anticoagulation is expected with high probability, aspirin is not suitable for long term application.

[0006] Besides aspirin, some pharmaceuticals are known to have inhibitory action against NF- κ B activation. Glucocorticoids (steroid hormones) such as dexamethasone suppress NF- κ B activation by binding to their receptors (called as glucocorticoid receptor, Scheinman R. I., et al., Science, 270, 283(1995)). However, long term use is not suitable, because they have serious side effects such as aggravation of an infectious disease, generation of peptic ulcer, degradation of bone density, and central action. Leflunomide as an immunosuppressive agent, as an isoxazole-type agent, also has NF- κ B inhibitory action (Manna S., et al., J. Immunol., 164, 2095-2102(1999)), however, the drug is also not suitable for long term use due to serious side effects. Furthermore, substituted pyrimidine derivatives (Japanese Patent Publication of International Application (Kohyo) (Hei)11-512399, J. Med. Chem., 41, 413(1998)), xanthine derivatives (Japanese Patent Unexamined Publication (KOKAI) No.(Hei)9-227561), isoquinoline derivatives (Japanese Patent Unexamined Publication (KOKAI) No.(Hei) 10-87491), indan derivatives (WO00/05234 epoxyquinomycin C, D, and their derivatives (Japanese Patent Unexamined Publication (KOKAI) No.(Hei)10-45738, Bioorg. Med. Chem. Lett., 10, 865-869(2000)) are known as inhibitors against NF- κ B activation. However, mechanism of inhibition against NF- κ B activation and participating receptors or proteins have not been revealed.

Disclosure of the Invention

[0007] Compounds having specific inhibitory action against IKK- β , found by using IKK- β as a target which directly induces phosphorylation of IKK- β , are expected to have inhibitory action against production and release of the target inflammatory cytokine and inhibitory action against production of inflammatory cell adhesion molecules, without affecting other signal transfer pathway, that is, without serious side effects. NF- κ B activation is induced by the aforementioned external stimulation, and as a result, proteins such as inflammatory cytokine are expressed. Among the inflammatory cytokines, TNF- α and interleukin (IL)-1 whose gene expression itself is considered to be regulated positively by NF- κ B to form positive feedback loop (TNF- α →NF- κ B→TNF- α) and is considered to participate in chronicity


of inflammation (18th Meeting of The Japanese Inflammatory Society, Symposium "Mechanism of Antirheumatic Pharmaceutical composition and New Development" Tokyo, 2000). Accordingly, the compounds which specifically inhibit IKK- β as a target are expected to be useful drugs for inflammatory diseases advanced in a chronic stage and diseases caused by TNF- α and IL-1.

[0008] Therefore, an object of the present invention is to provide medicaments useful for preventive and/or therapeutic treatment of inflammatory disorders, autoimmune disease such as chronic arthrorheumatism, and bone disease such as osteoporosis, in which inflammatory cytokine is participated. Another object of the present invention is to provide an inhibitor against release of an inflammatory cytokine which avoids side effects by specifically inhibiting IKK- β , and has inhibitory activity against NF- κ B activation.

[0009] The inventors of the present invention carried out search for compounds having inhibitory action against NF- κ B activation by selective inhibition of IKK- β by using computerized molecular design technology to solve the aforementioned object. Appropriate protein kinases with high homology with IKK- β were selected from the kinases whose structures are registered in PDB (Protein Data Bank), and three-dimensional structure model of IKK- β was constructed by applying the homology modeling technique employing the chosen kinase as a template, and then binding mode of aspirin to the ATP binding region or IKK- β and characteristic intermolecular interactions were analyzed by using automatic search program for binding modes of a drug molecule to a protein. On the basis of the results obtained, an automatic search program of a ligand from a three-dimensional compound database based on the three-dimensional structure of the protein was carried out, and compounds potentially be specific inhibitors against IKK- β were selected by a virtual screening out of compounds registered in a database of commercial compounds. Further, inhibitory activity of those compounds against NF- κ B activation was confirmed by a reporter assay method under TNF- α stimulation. Among them, the compounds with potent activities were further studied on the binding mode to IKK- β and interactions. On the basis of these results, the present invention was achieved by further carrying out search from compound databases of analogous compounds and syntheses

[0010] The medicament of the present invention is:

(1) that having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of a compound represented by the general formula (I) and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof:



wherein

X represents a connecting group whose number of atoms in the main chain is 2 to 4 (said connecting group may be substituted),

A represents hydrogen atom or acetyl group,

E represents an aryl group which may be substituted or a hetero aryl group which may be substituted,





ring Z represents an arene which may have one or more substituents in addition to the group represented by formula -O-A wherein A has the same meaning as that defined above and the group represented by formula -X-E wherein each of X and E has the same meaning as that defined above, or a hetero arene which may have one or more substituents in addition to the group represented by formula -O-A wherein A has the same meaning as that defined above and the group represented by formula -X-E wherein each of X and E has the same meaning as that defined above.

[0011] Among them, preferred medicaments include:

(2) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein X is a group selected from the following connecting group α which may be substituted:


EP 1 352 650 A1

[Connecting groups α]: The groups of the following formulas:




wherein the bond at the left end binds to ring Z and the bond at the right end binds to E;

(3) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein X is a group which may be substituted and represented by the following formula:



wherein the bond at the left end binds to ring Z and the bond at the right end binds to E;

(4) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein X is a group represented by the following formula:



wherein the bond at the left end binds to ring Z and the bond at the right end binds to E;

(5) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein A is a hydrogen atom;

(6) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein ring Z is a C₆ to C₁₀ arene which may have one or more substituents in addition to the group represented by formula -O-A wherein A has the same meaning as that defined in the general formula(I) and the group represented by formula -X-E wherein each of X and E has the same meaning as that defined in the general formula (I), or a 6 to 13-membered hetero arene which may have one or more substituents in addition to the group represented by formula -O-A wherein A has the same meaning as that defined in the general formula(I) and the group represented by formula -X-E wherein each of X and E has the same meaning as that defined in the general formula(I);

(7) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein ring Z is selected from the following ring group β which may have one or more substituents in addition to the group represented by formula -O-A wherein A has the same meaning as that defined in the general formula(I) and the group represented by formula -X-E wherein each of X and E has the same meaning as that defined in the general formula(I):


[Ring Group β] benzene ring, naphthalene ring, pyridine ring, indole ring, quinoxaline ring, carbazole ring;

(8) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein ring Z is a benzene ring which may have one or more substituents in addition to the group represented by formula -O-A wherein A has the same meaning as that defined in the general formula (I) and the group represented by formula -X-E wherein each of X and E has the same meaning as that defined in the general formula (I);


(9) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein ring Z is a benzene ring which may further have one or more substituents selected from the following substituent group γ -1z in addition to the group represented by formula -O-A wherein A has the same meaning as that defined in the general formula (I) and the group represented by formula -X-E wherein each of X and E has the same meaning as that defined in the general formula (I):

[substituent group γ -1z] a halogen atom, nitro group, cyano group, hydroxy group which may be substituted, amino group which may be substituted, hydrocarbon group which may be substituted, heterocyclic group which may be substituted, acyl group which may be substituted, ureido group which may be substituted, thioureido group which may be substituted, diazenyl group which may be substituted;

(10) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein the following partial structural formula (Iz-1) including ring Z in the general formula(I) :



is a group represented by the following formula (Iz-2)



(1 z - 2)

wherein R^2 represents hydrogen atom, a halogen atom, nitro group, cyano group, hydroxy group which may be substituted, an amino group which may be substituted, a hydrocarbon group which may be substituted, a heterocyclic group which may be substituted, an acyl group which may be substituted, an ureido group which may be substituted, a thioureido group which may be substituted, or a diazenyl group which may be substituted;

(11) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein R^2 is hydrogen atom, a halogen atom, nitro group, cyano group, a C_1 to C_6 alkoxy group which may be substituted, a di(C_1 to C_6 alkyl)-amino group, a C_6 to C_{10} aryl-carbonyl-amino group, a C_1 to C_6 alkyl group which may be substituted, a C_1 to C_6 halogenated alkyl group which may be substituted, a C_2 to C_6 alkenyl group which may be substituted, a C_2 to C_6 alkynyl group which may be substituted, a C_7 to C_{16} aralkyl group which may be substituted, a 5 to 6 membered heteroaryl group which may be substituted, a carbamoyl group which may be substituted, a sulfamoyl group which may be substituted, a C_1 to C_6 alkyl-carbamoyl group which may be substituted, a C_1 to C_6 alkoxy-carbamoyl group which may be substituted, a 5 membered heteroaryl-sulfonyl group which may be substituted, a 6 membered nonaromatic heterocyclic-sulfonyl group which may be substituted, or a diazenyl group which may be substituted;

(12) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein R^2 is hydrogen atom, a halogen atom, nitro group, cyano group, methoxy group, dimethylamino group, benzoylamino group, methyl group, tert-butyl group, 1-hydroxyethyl group, 1-(methoxyimino)ethyl group, 1-((benzyloxy)imino)ethyl group, trifluoromethyl group, pentafluoroethyl group, phenyl group, 4-(trifluoromethyl)phenyl group, 4-fluorophenyl group, 2,4-difluorophenyl group, 2-phenylethen-1-yl group, 2,2-dicyanoethen-1-yl group, 2-cyano-2-(methoxycarbonyl)ethen-1-yl group, 2-carboxy-2-cyanoethen-1-yl group, ethynyl group, phenylethynyl group, (trimethylsilyl)ethynyl group, phenyl group, 2-phenethyl group, 2-thienyl group, 3-thienyl group, 1-pyrrolyl group, 2-methylthiazol-4-yl group, 2-pyridyl group, N-[3,5-bis(trifluoromethyl)-phenyl]carbamoyl group, dimethylcarbamoyl group, dimethylsulfamoyl group, acetyl group, isobutyryl group, methoxycarbonyl group, piperidinocarbonyl group, 4-benzylpiperidino group, (pyrrol-1-yl)sulfonyl group, 3-phenylureido group, (3-phenyl)thioureido group, (4-nitrophenyl)diazenyl group, or {[(4-pyridin-2-yl)sulfamoyl]phenyl}diazenyl group;

(13) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein R^2 is a halogen atom;

(14) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a C_6 to C_{10} aryl group which may be substituted or a 5 to 13 membered heteroaryl group which may be substituted;

(15) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a C_6 to C_{10} aryl group which may be substituted;

(16) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a phenyl group which may be substituted;

(17) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a phenyl group substituted with two C_1 to C_6 halogenated alkyl groups and said phenyl group may further have one or more substituents in addition to the two C_1 to C_6 halogenated alkyl groups;

(18) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient

a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a phenyl group substituted with two C₁ to C₆ halogenated alkyl groups;

(19) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is 3,5-bis(trifluoromethyl)phenyl group or 2,5-bis(trifluoromethyl)phenyl group;

(20) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is 3,5-bis(trifluoromethyl)phenyl group;

(21) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a phenyl group substituted with one C₁ to C₆ halogenated alkyl group and said phenyl group may further have one or more substituents, except a C₁ to C₆ halogenated alkyl group, in addition to the C₁ to C₆ halogenated alkyl group;

(22) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a phenyl group substituted with one C₁ to C₆ halogenated alkyl group and said phenyl group may further have one or more substituents selected from the following substituent group γ -1e in addition to the C₁ to C₆ halogenated alkyl group:

[Substituent group γ -1e] a halogen atom, nitro group, cyano group, hydroxy group which may be substituted, a hydrocarbon group which may be substituted, a heterocyclic group which may be substituted, sulfanyl group which may be substituted;

(23) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a phenyl group substituted with one C₁ to C₆ halogenated alkyl group and said phenyl group may further have one or more substituents selected from the following substituent group γ -2e in addition to the C₁ to C₆ halogenated alkyl group:

[Substituent group γ -2e] a halogen atom, nitro group, cyano group, a C₁ to C₆ alkyl group which may be substituted, a 5 to 6 membered nonaromatic heterocyclic group which may be substituted, a C₁ to C₆ alkoxy group which may be substituted, a C₁ to C₆ alkyl-sulfanyl group which may be substituted;

(24) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is 2-(trifluoromethyl)phenyl group, 3-(trifluoromethyl)phenyl group, 4-(trifluoromethyl)phenyl group, 2-fluoro-3-(trifluoromethyl)phenyl group, 2-chloro-4-(trifluoromethyl)phenyl group, 2-fluoro-5-(trifluoromethyl)phenyl group, 2-chloro-5-(trifluoromethyl)phenyl group, 3-fluoro-5-(trifluoromethyl)phenyl group, 3-bromo-5-(trifluoromethyl)phenyl group, 4-chloro-2-(trifluoromethyl)phenyl group, 4-fluoro-3-(trifluoromethyl)phenyl group, 4-chloro-3-(trifluoromethyl)phenyl group, 2-nitro-5-(trifluoromethyl)phenyl group, 4-nitro-3-(trifluoromethyl)phenyl group, 4-cyano-3-(trifluoromethyl)phenyl group, 2-methyl-3-(trifluoromethyl)phenyl group, 2-methyl-5-(trifluoromethyl)phenyl group, 4-methyl-3-(trifluoromethyl)phenyl group, 2-methoxy-5-(trifluoromethyl)phenyl group, 3-methoxy-5-(trifluoromethyl)phenyl group, 4-methoxy-3-(trifluoromethyl)phenyl group, 2-(methylsulfanyl)-5-(trifluoromethyl)phenyl group, 2-(1-pyrrolizino)-5-(trifluoromethyl)phenyl group, or 2-morpholino-5-(trifluoromethyl)phenyl group;

(25) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is 2-chloro-5-(trifluoromethyl)phenyl group, 4-chloro-3-(trifluoromethyl)phenyl group, 2-methoxy-5-(trifluoromethyl)phenyl group, or 3-methoxy-5-(trifluoromethyl)phenyl group;

(26) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is 2-chloro-5-(trifluoromethyl)phenyl group;

(27) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a phenyl group which may have one or more substituent

except a C₁ to C₆ halogenated alkyl group;

(28) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a phenyl group which may have one or more substituents selected from the following substituent group γ -3e:

[Substituent group γ -3e] a halogen atom, nitro group, hydroxy group which may be substituted, hydrocarbon group which may be substituted, acyl group which may be substituted;

(29) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a phenyl group which is substituted with one or more substituents selected from the following substituent group γ -4e;

[Substituent group γ -4e] a halogen atom, nitro group, hydroxy group, a C₁ to C₆ alkyl group which may be substituted, a C₆ to C₁₀ aryl group which may be substituted, a C₁ to C₆ alkylene group which may be substituted, a C₁ to C₆ alkoxy group which may be substituted, a C₁ to C₆ alkyl-carbonyl group which may be substituted, a C₁ to C₆ alkoxy-carbonyl group which may be substituted;

(30) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is phenyl group, 3-chlorophenyl group, 4-chlorophenyl group, 2,5-dichlorophenyl group, 3,4-dichlorophenyl group, 3,5-difluorophenyl group, 3,5-dichlorophenyl group, 3,4,5-trichlorophenyl group, pentafluorophenyl group, 3,5-dinitrophenyl group, 3,5-dichloro-4-hydroxyphenyl group, 2,5-dimethoxyphenyl group, 3,5-dimethoxyphenyl group, 3,5-dimethylphenyl group, 2,5-bis[(1,1-dimethyl)ethyl]phenyl group, 3,5-bis[(1,1-dimethyl)ethyl]phenyl group, 5-(1,1-dimethyl)ethyl-2-methoxyphenyl group, 3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydronaphthalen-2-yl group, biphenyl-3-yl group, 4-methoxybiphenyl-3-yl group, 3-acetylphenyl group, or 3,5-bis(methoxycarbonyl)phenyl group;

(31) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is 2,5-bis[(1,1-dimethyl)ethyl]phenyl group, 3,5-bis[(1,1-dimethyl)ethyl]phenyl group, or 5-(1,1-dimethyl)ethyl-2-methoxyphenyl group;

(32) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a 5 to 13 membered heteroaryl group which may be substituted;

(33) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a thienyl group which may be substituted, a pyrazolyl group which may be substituted, an oxazolyl group which may be substituted, a thiazolyl group which may be substituted, a thiadiazolyl group which may be substituted, a pyridyl group which may be substituted, a pyrimidinyl group which may be substituted, an indolyl group which may be substituted, a quinolyl group which may be substituted, or a carbazolyl group which may be substituted;

(34) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a thiazolyl group which may be substituted;

(35) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a thiazolyl group which may have one or more substituents selected from the following substituent group γ -5e:

[Substituent group γ -5e] a halogen atom, cyano group, a hydrocarbon group which may be substituted, a heteroring group which may be substituted, an acyl group which may be substituted;

(36) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is a thiazolyl group which may have one or more substituents selected from the following substituent group γ -6e:

[Substituent group γ -6e] a halogen atom, cyano group, a C_1 to C_6 alkyl group which may be substituted, a C_1 to C_6 halogenated alkyl group which may be substituted, a C_6 to C_{10} aryl group which may be substituted, a C_7 to C_{16} aralkyl group which may be substituted, a 6 membered nonaromatic heteroring group, a C_1 to C_6 alkyl-carbonyl group which may be substituted, a C_6 to C_{10} aryl-carbonyl group which may be substituted, a C_1 to C_6 alkoxy-carbonyl group which may be substituted;

(37) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is 5-bromo-4-[(1,1-dimethyl)ethylthiazol-2-yl] group, 5-bromo-4-(trifluoromethyl)thiazol-2-yl group, 5-cyano-4-[(1,1-dimethyl)ethylthiazol-2-yl] group, 4-[(1,1-dimethyl)ethylthiazol-2-yl] group, 5-phenyl-4-(trifluoromethyl)thiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-ethylthiazol-2-yl group, 5-methyl-4-phenylthiazol-2-yl group, 4-isopropyl-5-phenylthiazol-2-yl group, 4-benzyl-5-phenylthiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionylthiazol-2-yl] group, 5-acetyl-4-phenylthiazol-2-yl group, 5-benzoyl-4-phenylthiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-(ethoxycarbonyl)thiazol-2-yl group, 5-ethoxycarbonyl-4-(trifluoromethyl)thiazol-2-yl group, 5-ethoxycarbonyl-4-phenylthiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-piperidin-4-(trifluoromethyl)thiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-morpholinethiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-(4-phenylpiperidin-1-yl)thiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-(4-methylpiperidin-1-yl)thiazol-2-yl group, 4,5-diphenylthiazol-2-yl group, 4-phenylthiazol-2-yl group, 4,5-dimethylthiazol-2-yl group, 2-thiazolyl group, 5-methylthiazol-2-yl group, 4-ethyl-5-phenylthiazol-2-yl group, 5-carboxymethyl-4-phenylthiazol-2-yl group, 5-methylcarbamoyl-4-phenylthiazol-2-yl group, 5-ethylcarbamoyl-4-phenylthiazol-2-yl group, 5-isopropylcarbamoyl-4-phenylthiazol-2-yl group, 5-(2-phenethyl)carbamoyl-4-phenylthiazol-2-yl group, 4-(n-butyl)-5-phenylthiazol-2-yl group, 4-methyl-5-[(3-trifluoromethyl)phenyl]thiazol-2-yl group, or 5-(4-fluorophenyl)-4-methylthiazol-2-yl group; and

(38) the medicament having inhibitory action against NF- κ B activation which comprises as an active ingredient a substance selected from the group consisting of the compound and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, wherein E is 4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl group.

[0012] The medicament of the present invention may be used as gene expression inhibitor of one or more substances selected from the following substance group δ :

[Substance group δ] tumor necrosis factor (TNF), interleukin-1, interleukin-2, interleukin-6, interleukin-8, granulocyte colony-stimulating factor, interferon β , cell adhesion factor ICAM-1, VCAM-1, ELAM-1, nitric oxide synthetase, major histocompatibility antigen family class I, major histocompatibility antigen family class II, β 2-microglobulin, immunoglobulin light chain, serum amyloid A, angiotensinogen, complement B, complement C4, c-myc, transcript derived from HIV gene, transcript derived from HTLV gene, transcript derived from simian virus 40 gene, transcript derived from cytomegalovirus gene, and transcript derived from adenovirus gene.

[0013] The medicament of the present invention may be used as an inhibitor against production and release of an inflammatory cytokine or as an immunosuppressive agent. The medicament of the present invention may be used for preventive and/or therapeutic treatment of one or more diseases selected from the following disease group ϵ -1:

[Disease group ϵ -1] an inflammatory disease, an autoimmune disease, an allergic disease, a cancers such as carcinoma and sarcoma, a metabolic disease, a cardiovascular disease, an angioproliferation disease, a septic disease, a viral disease, or from the following disease group ϵ -2 which is resulted from NF- κ B activation or overproduction of an inflammatory cytokine:


[Disease group ϵ -2] an autoimmune diseases such as chronic rheumatism, osteoarthritis, systematic lupus erythematosus, systematic scleroderma, polymyositis, Sjogren's syndrome, vasculitis syndrome, antiphospholipid syndrome, Still's disease, Behcet's disease, periarteritis nodosa, ulcerative colitis, Crohn's disease, active chronic hepatitis, glomerulonephritis, and chronic nephritis, chronic pancreatitis, gout, atherosclerosis, multiple sclerosis, arteriosclerosis, endothelial hypertrophy, psoriasis, psoriatic arthritis, contact dermatitis, atopic dermatitis, allergic disease such as pollinosis, asthma, bronchitis, interstitial pneumonia, lung disease involving granuloma, chronic obstructive pulmonary disease, chronic pulmonary thromboembolism, inflammatory colitis, insulin resistance, obesity, diabetes and its complications (nephropathy, retinopathy, neurosis, hyperinsulinemia, arteriosclerosis, hypertension, peripheral vessel obstruction and the like) a disease with abnormal vascular proliferation such as hyperlipemia, retinopathy, pneumonia, Alzheimer's disease, encephalomyelitis, acute hepatitis, chronic hepatitis, pharmaceutical composition induced toxic hepatopathy, alcoholic hepatitis, viral hepatitis, icterus, cirrhosis, hepatic insufficiency, atrial myxoma, Caslemani's syndrome, mesangial nephritis, kidney cancer, lung cancer, liver cancer, breast cancer, uterine cancer, pancreatic cancer, other solid cancer, sarcoma, osteosarcoma, metastatic invasion

of cancer, carcination of inflammatory focus, cancerous cachexia, metastasis of cancer, leukemia such as acute myeloblastic leukemia, multiple myeloma, Lennert's lymphoma, malignant lymphoma, development of carcinostatic resistance of cancer, carcination of foci such as viral hepatitis and cirrhosis, carcination from polyp of colon, brain tumor, nervous tumor, endotoxic shock, sepsis, cytomegaloviral pneumonia, cytomegaloviral retinopathy, adenoviral cold, adenoviral pool fever, adenoviral ophthalmia, conjunctivitis, AIDS, uveitis, diseases or complications provoked by infections of other bacteria, viruses, and mycete, complications after surgery such as generalized inflammatory symptoms, restenosis after percutaneous tubal coronary artery plastic surgery, reperfusion disorders after vascular occulusion opening such as ischemia reperfusion disorders, organ transplantation rejection and reperfusion disorders of heart, liver, or kidney, etc., itch, anorexia, malaise, chronic fatigue syndrome, osteoporosis, metabolic bone disease such as osteocarcinomic pain, deterioration of organ during organ conservation before transplantation.

[0014] From another aspect, the present invention provides use of each of the substances for manufacture of the medicament according to the aforementioned (1) to (38).

[0015] Furthermore, the present invention provides: a method for inhibiting activation of NF- κ B in a mammal including a human, which comprises the step of administering the aforementioned medicament (1) to (38) to a mammal including a human; a method for inhibiting expression of one or more substances selected from the aforementioned substance group δ in a mammal including a human, which comprises the step of administering the aforementioned medicament (1) to (38) to a mammal including a human; a method for inhibiting production and release of an inflammatory cytokine in a mammal including a human, which comprises the step of administering the aforementioned medicament (1) to (38) to a mammal including a human; a method for immune inhibition in a mammal including a human, which comprises the step of administering the aforementioned medicament (1) to (38) to a mammal including a human; a method for preventive and/or therapeutic treatment of one or more diseases selected from aforementioned disease group ϵ -1, which comprises the step of administering the aforementioned medicament (1) to (38) to a mammal including a human; and a method for preventive and/or therapeutic treatment of one or more diseases selected from aforementioned disease group ϵ -2 caused by NF- κ B activation or inflammatory cytokine overproduction, which comprises the step of administering the aforementioned medicament (1) to (38) to a mammal including a human.

[0016] The present invention further provides (1) a compound represented by the general formula (I-1) or a salt thereof, or a hydrate thereof or a solvate thereof:



wherein Z^1 represents 2-hydroxyphenyl group which may have a substituent in the 5-position or 2-acetoxyphenyl group, which may have a substituent in the 5-position, E^1 represents a phenyl group substituted with two C_1 to C_6 halogenated alkyl groups wherein said phenyl group may further have one or more substituents in addition to the two C_1 to C_6 halogenated alkyl groups, provided that the following compounds are excluded:

N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide,
N-[3,5-bis(trifluoromethyl)phenyl]-5-chloro-2-hydroxybenzamide,
N-[3,5-bis(trifluoromethyl)phenyl]-5-bromo-2-hydroxybenzamide,
N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-iodobenzamide,
N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-nitrobenzamide, and
2-hydroxy-N-[2,3,5-tris(trifluoromethyl)phenyl]benzamide).


[0017] Preferred examples include:

(2) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E^1 is a phenyl group substituted with two C_1 to C_6 alkyl groups each of which is substituted with one or more fluorine atoms wherein said phenyl group may further have one or more substituents in addition to the two C_1 to C_6 alkyl groups which are substituted with one or more fluorine atoms;
(3) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E^1 is a phenyl group substituted with two C_1 to C_6 alkyl groups each of which is substituted with three or more

fluorine atoms wherein said phenyl group may further have one or more substituents in addition to the two C₁ to C₆ alkyl groups each of which is substituted with three or more fluorine atoms;

(4) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E¹ is a phenyl group substituted with two C₁ to C₆ alkyl groups each of which is substituted with three or more fluorine atoms;

(5) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E¹ is a group represented by the following formula :




wherein one of R^{1e2} and R^{1e3} represents hydrogen atom and the other represents a C₁ to C₆ alkyl group substituted with three or more fluorine atoms, and R^{1e5} represents a C₁ to C₆ alkyl group substituted with three or more fluorine atoms;

(6) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E¹ is 3,5-bis(trifluoromethyl)phenyl group, or 2,5-bis(trifluoromethyl)phenyl group;

(7) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E¹ is 2,5-bis(trifluoromethyl)phenyl group;

(8) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein Z¹ is a group represented by the following formula :



wherein A¹ represents hydrogen atom or acetyl group, R^{1z} represents hydrogen atom, a halogen atom, nitro group, cyano group, hydroxy group which may be substituted, amino group which may be substituted, a hydrocarbon group which may be substituted, a heteroring group which may be substituted, an acyl group which may be substituted, an ureido group which may be substituted, a thioureido group which may be substituted, a diazenyl group which may be substituted;

(9) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein A¹ is hydrogen atom;

(10) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein R^{1z} is hydrogen atom, a halogen atom, nitro group, cyano group, a C₁ to C₆ alkoxy group which may be substituted, a di(C₁ to C₆ alkyl)-amino group, a C₁ to C₆ alkyl group which may be substituted, a C₁ to C₆ halogenated alkyl group which may be substituted, a C₂ to C₆ alkenyl group which may be substituted, a C₂ to C₆ alkynyl group which may be substituted, a C₆ to C₁₀ aryl group which may be substituted, a C₇ to C₁₆ aralkyl group which may be substituted, a 5 to 6 membered heteroaryl group which may be substituted, a C₁ to C₆ alkyl-carbonyl group which may be substituted, a C₁ to C₆ alkoxy-carbonyl group which may be substituted, a 5-membered heteroaryl-sulfonyl group which may be substituted, or diazenyl group which may be substituted;

(11) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein R^{1z} is a halogen atom, a C₁ to C₆ alkyl groups which may be substituted, a C₁ to C₆ halogenated alkyl group which


may be substituted; and

(12) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein R¹² is a halogen atom.

5 **[0018]** Most preferably, the following compounds or pharmacologically acceptable salts thereof, or hydrates thereof or solvates thereof are provided.

N-[3,5-bis(trifluoromethyl)phenyl]-5-fluoro-2-hydroxybenzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-5-cyano-2-hydroxybenzamide,
 10 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-methylbenzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-5-(1,1-dimethyl)ethyl-2-hydroxybenzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(trifluoromethyl)benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(1,1,2,2,2-pentafluoroethyl)benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(2-phenylethen-1-yl)benzamide,
 15 N-[3,5-bis(trifluoromethyl)phenyl]-5-(2,2-dicyanoethen-1-yl)-2-hydroxybenzamide,
 3-({[3,5-bis(trifluoromethyl)phenyl]carbamoyl}-4-hydroxyphenyl)-2-cyanoacrylic acid methyl ester,
 3-({[3,5-bis(trifluoromethyl)phenyl]carbamoyl}-4-hydroxyphenyl)-2-cyanoacrylic acid, ...
 N-[3,5-bis(trifluoromethyl)phenyl]-5-ethynyl-2-hydroxybenzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(phenylethynyl)benzamide,
 20 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-[(trimethylsilyl)ethynyl]benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-4-hydroxybiphenyl-3-carboxamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(2-phenylethyl)benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(3-thienyl)benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(1-pyrrolyl)benzamide,
 25 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(2-methylthiazol-4-yl)benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(2-pyridyl)benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-5-dimethylamino-2-hydroxybenzamide,
 5-benzoylamino-N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 N³-[3,5-bis(trifluoromethyl)phenyl]-4-hydroxy-N¹,N¹-dimethyliso phthalamide,
 30 N¹,N³-bis[3,5-bis(trifluoromethyl)phenyl]-4-hydroxyisophthalamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(piperidine-1-carbonyl)benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(4-benzylpiperidine-1-carbonyl)-benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-5-dimethylsulfamoyl-2-hydroxybenzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(pyrrole-1-sulfonyl)benzamide,
 35 5-acetyl-N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-isobutyrylbenzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-4-hydroxyisophthalamide acid methylester,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-[(4-nitrophenyl)diazenyl]benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-[[[(4-pyridin-2-yl)sulfamoyl]phenyl]diazanyl]benzamide,
 40 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-[(3-phenyl)ureido]benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-[(3-phenyl)thioureido]benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(1-hydroxyethyl)benzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-methoxybenzamide,
 N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-[1-(methoxyimino)ethyl]benzamide,
 45 5-[1-[(benzyloxy)imino]ethyl]-N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 N-[2,5-bis(trifluoromethyl)phenyl]-5-chloro-2-hydroxybenzamide,
 N-[2,5-bis(trifluoromethyl)phenyl]-5-bromo-2-hydroxybenzamide,
 2-acetoxy-N-[3,5-bis(trifluoromethyl)phenyl]benzamide,
 2-acetoxy-N-[2,5-bis(trifluoromethyl)phenyl]-5-chlorobenzamide, and
 50 2-acetoxy-N-[3,5-bis(trifluoromethyl)phenyl]-5-chlorobenzamide.

[0019] The present invention further provides the compound represented by the following general formula (I-2) or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof:



wherein Z^2 represents 2-hydroxyphenyl group which may be substituted in the 5-position, or 2-acetoxyphenyl group which may be substituted in the 5-position, E^2 represents a phenyl group whose 3-position or 5-position is substituted with a C_1 to C_6 halogenated alkyl group wherein said phenyl group may further have one or more substituents (except when the substituent is a C_1 to C_6 halogenated alkyl group) in addition to the C_1 to C_6 halogenated alkyl group in the 3-position or 5-position, provided that the following compounds are excluded:

5-chloro-2-hydroxy-N-[3-(trifluoromethyl)phenyl]benzamide,
 5-bromo-2-hydroxy-N-[3-(trifluoromethyl)phenyl]benzamide,
 2-hydroxy-5-iodo-N-[3-(trifluoromethyl)phenyl]benzamide,
 5-chloro-N-(4-chloro-3-(trifluoromethyl)phenyl)-2-hydroxybenzamide,
 5-chloro-N-[5-chloro-3-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-chloro-2-hydroxy-N-[4-nitro-3-(trifluoromethyl)phenyl]benzamide,
 5-fluoro-2-hydroxy-N-[2-(2,2,2-trifluoroethoxy)-5-(trifluoromethyl)phenyl]benzamide,
 5-fluoro-2-hydroxy-N-[2-(6,6,6-trifluorohexyloxy)-5-(trifluoromethyl)phenyl]benzamide
 5-chloro-2-hydroxy-N-[3-(trifluoromethyl)-4-[(4-(trifluoromethyl)sulfanyl]phenoxy)-phenyl]benzamide
 N-[4-(benzothiazol-2-yl)sulfanyl-3-(trifluoromethyl)phenyl]-5-chloro-2-hydroxybenzamide
 5-chloro-N-[2-(4-chlorophenoxy)-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-chloro-2-hydroxy-N-[2-(4-methylphenoxy)-5-(trifluoromethyl)phenyl]benzamide,
 5-chloro-N-[2-(4-chlorophenyl)sulfanyl-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide
 5-chloro-2-hydroxy-N-[2-(1-naphthyloxy)-5-(trifluoromethyl)phenyl]benzamide, and
 5-chloro-2-hydroxy-N-[2-(2-naphthyloxy)-5-(trifluoromethyl)phenyl]benzamide.

[0020] Preferred examples include:

(2) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E^2 is a phenyl group whose 3-position or 5-position is substituted with a C_1 to C_6 alkyl group which is substituted with one or more fluorine atoms wherein said phenyl group may have one or more substituents (except when the substituent is a C_1 to C_6 halogenated alkyl group) in addition to the C_1 to C_6 alkyl group in the 3-position or the 5-position which is substituted with one or more fluorine atoms;

(3) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E^2 is a phenyl group whose 3-position or 5-position is substituted with a C_1 to C_6 alkyl group which is substituted with three or more fluorine atoms wherein said phenyl group may have one or more substituents (except when the substituent is a C_1 to C_6 halogenated alkyl group) in addition to the C_1 to C_6 alkyl group in the 3-position or the 5-position which is substituted with three or more fluorine atoms;


(4) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E^2 is a phenyl group whose 3-position or 5-position is substituted with a C_1 to C_6 alkyl group which is substituted with three or more fluorine atoms wherein said phenyl group may have one or more substituents selected from the following substituent group γ -7e in addition to the C_1 to C_6 alkyl group in the 3-position or 5-position which is substituted with three or more fluorine atoms:

[substituent group γ -7e] a halogen atom, nitro group, cyano group, a C_1 to C_6 alkyl group which may be substituted, a 5 to 6 membered nonaromatic heteroring group which may be substituted, a C_1 to C_6 alkoxy group which may be substituted, a C_1 to C_6 alkyl-sulfanyl group which may be substituted;

(6) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E^2 is 3-(trifluoromethyl)phenyl group, 2-fluoro-3-(trifluoromethyl)phenyl group, 2-fluoro-5-(trifluoromethyl)phenyl group, 2-chloro-5-(trifluoromethyl)phenyl group, 3-fluoro-5-(trifluoromethyl)phenyl group, 3-bromo-5-(trifluoromethyl)phenyl group, 4-fluoro-3-(trifluoromethyl)phenyl group, 4-chloro-3-(trifluoromethyl)phenyl group, 2-nitro-5-(trifluoromethyl)phenyl group, 4-nitro-3-(trifluoromethyl)phenyl group, 4-cyano-3-(trifluoromethyl)phenyl group, 2-methyl-3-(trifluoromethyl)phenyl group, 2-methyl-5-(trifluoromethyl)phenyl group, 4-methyl-3-(trifluoromethyl)phenyl-

group, 2-methoxy-5-(trifluoromethyl)phenyl group, 3-methoxy-5-(trifluoromethyl)phenyl group, 4-methoxy-3-(trifluoromethyl)phenyl group, 2-(methylsulfanyl)-5-(trifluoromethyl)phenyl group, 2-(1-pyrrolidino)-5-(trifluoromethyl)phenyl group, or 2-morpholino-5-(trifluoromethyl)phenyl group;

(7) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein Z^2 is a group represented by the following formula:



wherein A^2 represents hydrogen atom or acetyl group, R^Z represents a halogen atom, a C_1 to C_6 alkyl group or a C_1 to C_6 halogenated alkyl group;


(8) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein A^2 is hydrogen atom; and

(9) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein R^Z is halogen atom.


[0021] Most preferably, the following compounds or pharmacologically acceptable salts thereof, or hydrates thereof or solvates thereof are provided:

5-chloro-N-[2-fluoro-3-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-chloro-N-[2-fluoro-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-chloro-N-[2-chloro-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-bromo-N-[2-chloro-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 2-acetoxy-5-chloro-N-[2-chloro-5-(trifluoromethyl)phenyl]benzamide,
 N-[2-chloro-5-(trifluoromethyl)phenyl]-2-hydroxy-5-methylbenzamide,
 5-chloro-N-[3-fluoro-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-bromo-N-[3-bromo-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-chloro-N-[3-fluoro-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-chloro-N-[4-fluoro-3-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-bromo-N-[4-chloro-3-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 N-[4-chloro-3-(trifluoromethyl)phenyl]-2-hydroxy-5-methylbenzamide,
 5-chloro-2-hydroxy-N-[2-nitro-5-(trifluoromethyl)phenyl]benzamide,
 5-bromo-N-[4-cyano-3-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-chloro-2-hydroxy-N-[2-methyl-3-(trifluoromethyl)phenyl]benzamide,
 5-chloro-2-hydroxy-N-[2-methyl-5-(trifluoromethyl)phenyl]benzamide,
 2-hydroxy-5-methyl-N-[2-methyl-5-(trifluoromethyl)phenyl]benzamide,
 5-chloro-2-hydroxy-N-[4-methyl-3-(trifluoromethyl)phenyl]benzamide,
 2-hydroxy-5-methyl-N-[4-methyl-3-(trifluoromethyl)phenyl]benzamide,
 5-bromo-2-hydroxy-N-[2-methoxy-5-(trifluoromethyl)phenyl]benzamide,
 5-chloro-2-hydroxy-N-[2-methoxy-5-(trifluoromethyl)phenyl]benzamide,
 2-hydroxy-N-[2-methoxy-5-(trifluoromethyl)phenyl]-5-methylbenzamide,
 5-bromo-2-hydroxy-N-[3-methoxy-5-(trifluoromethyl)phenyl]benzamide,
 5-chloro-2-hydroxy-N-[4-methoxy-3-(trifluoromethyl)phenyl]benzamide,
 2-hydroxy-N-[4-methoxy-3-(trifluoromethyl)phenyl]-5-methylbenzamide,
 5-chloro-2-hydroxy-N-[2-methylsulfanyl-5-(trifluoromethyl)phenyl]benzamide,
 5-chloro-2-hydroxy-N-[2-(1-pyrrolidino)-5-(trifluoromethyl)phenyl]benzamide, and
 5-chloro-2-hydroxy-N-[2-morpholino-5-(trifluoromethyl)phenyl]benzamide.

[0022] Furthermore, the present invention provides (1) the compound represented by the following general formula (I-3) or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof



wherein Z^3 represents 2-hydroxyphenyl group which may be substituted in the 5-position or 2-acetoxyphenyl group which may be substituted in the 5-position, E^3 represents a group represented by the following formula:




wherein R^{3e2} and R^{3e3} may be the same or different and each represents hydrogen atom, a hydrocarbon group which may be substituted, or hydroxy group which may be substituted, except that where both of R^{3e2} and R^{3e3} are hydrogen atoms is excluded, R^{3e5} represents a C_2 to C_6 hydrocarbon group which may be substituted.

[0023] Preferred examples include:

(2) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein R^{3e2} and R^{3e3} may be the same or different and each represents hydrogen atom, a C_1 to C_6 alkyl group which may be substituted, or a C_1 to C_6 alkoxy group which may be substituted, provided that where both of R^{3e2} and R^{3e3} are hydrogen atoms is excluded, R^{3e5} is a C_2 to C_6 alkyl group which may be substituted;

(3) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E^3 is 2,5-bis[(1,1-dimethyl)ethyl]phenyl group, 3,5-bis[(1,1-dimethyl)ethyl]phenyl group, or 5-(1,1-dimethyl)ethyl-2-methoxyphenyl group;

(4) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein Z^3 is a group represented by the following formula:



wherein A^3 represents hydrogen atom or acetyl group, R^{3z} represents a halogen atom, a C_1 to C_6 alkyl group, or a C_1 to C_6 halogenated alkyl group;

(5) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein A^3 is hydrogen atom;

(6) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein R^{3z} is halogen atom.


[0024] Most preferably, the following compounds or pharmacologically acceptable salts thereof, or hydrates thereof or solvates thereof are provided:

N-{2,5-bis[(1,1-dimethyl)ethyl]phenyl}-5-chloro-2-hydroxybenzamide,


EP 1 352 650 A1

N-(2,5-bis [(1,1-dimethyl)ethyl] phenyl)-2-hydroxy-5-methylbenzamide,
 N-(3,5-bis[(1,1-dimethyl)ethyl] phenyl)-5-chloro-2-hydroxybenzamide,
 N-(3,5-bis[(1,1-dimethyl)ethyl] phenyl)-5-bromo-2-hydroxybenzamide,
 N-(3,5-bis[(1,1-dimethyl)ethyl] phenyl)-2-hydroxy-5-methylbenzamide,
 2-acetoxy-N-(3,5-bis[(1,1-dimethyl)ethyl] phenyl)-5-chlorobenzamide,
 5-chloro-N-[5-(1,1-dimethyl)ethyl-2-methoxyphenyl]-2-hydroxybenzamide,
 N-[5-(1,1-dimethyl)ethyl-2-methoxyphenyl]-2-hydroxy-5-benzamide, and
 2-acetoxy-5-chloro-N-(5-(1,1-dimethyl)ethyl-2-methoxyphenyl)benzamide.

[0025] The present invention further provides (1) the compound represented by the following general formula (I-4) or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof:



wherein Z⁴ represents 2-hydroxyphenyl group which may be substituted in the 5-position or 2-acetoxyphenyl group which may be substituted in the 5-position, E⁴ is a group represented by the following formula :



wherein R^{4e4} represents a hydrocarbon group which may be substituted, R^{4e5} represents a halogen atom, cyano group, acyl group which may be substituted, or a heteroring group which may be substituted.

[0026] Preferred examples include:

(2) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein R^{4e4} represents a C₁ to C₆ alkyl group which may be substituted, a C₁ to C₆ halogenated alkyl group which may be substituted, or a C₆ to C₁₀ aryl group which may be substituted, R^{4e5} represents a halogen atom, cyano group, a C₁ to C₆ alkyl-carbonyl group which may be substituted, a C₆ to C₁₀ aryl-carbonyl group which may be substituted, a C₁ to C₆ alkoxy-carbonyl group which may be substituted, or a 6-membered nonaromatic heteroring group which may be substituted;


(3) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E⁴ is 5-bromo-4-[(1,1-dimethyl)ethyl]thiazol-2-yl group,

5-bromo-4-(trifluoromethyl)thiazol-2-yl group,
 5-cyano-4-[(1,1-dimethyl)ethyl]thiazol-2-yl group,
 4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl group,
 5-acetyl-4-phenylthiazol-2-yl group, 5-benzoyl-4-phenylthiazol-2-yl group,
 4-(1,1-dimethyl)ethyl-5-(ethoxycarbonyl)thiazol-2-yl group,
 5-ethoxycarbonyl-4-(trifluoromethyl)thiazol-2-yl group,
 5-ethoxycarbonyl-4-phenylthiazol-2-yl group,
 5-ethoxycarbonyl-4-(pentafluorophenyl)thiazol-2-yl group,
 4-(1,1-dimethyl)ethyl-5-piperidinethiazol-2-yl group,
 4-(1,1-dimethyl)ethyl-5-morpholinethiazol-2-yl group,
 4-(1,1-dimethyl)ethyl-5-(4-methylpiperidin-1-yl)thiazol-2-yl group, or
 4-(1,1-dimethyl)ethyl-5-(4-phenylpiperidin-1-yl)thiazol-2-yl group;

EP 1 352 650 A1

(4) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein E⁴ is 4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl group;

(5) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein Z⁴ is a group represented by the following formula:



wherein A⁴ represents hydrogen atom or acetyl group, R^{4z} represents a halogen atom, a C₆ to C₁₀ aryl group which may be substituted, or a 5-membered heteroaryl group;

(6) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein A⁴ is hydrogen atom; and

(7) the compound or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof wherein R^{4z} is a halogen atoms.

[0027] Most preferably, the following compounds or pharmacologically acceptable salts thereof, or hydrates thereof or solvates thereof are provided.

5-bromo-N-{5-bromo-4-[(1,1-dimethyl)ethyl]thiazol-2-yl}-2-hydroxybenzamide,
 5-bromo-N-{5-bromo-4-(trifluoromethyl)thiazol-2-yl}-2-hydroxybenzamide,
 5-chloro-N-{5-cyano-4-[(1,1-dimethyl)ethyl]thiazol-2-yl}-2-hydroxybenzamide,
 5-bromo-N-{5-cyano-4-[(1,1-dimethyl)ethyl]thiazol-2-yl}-2-hydroxybenzamide,
 5-chloro-N-{4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl}-2-hydroxybenzamide,
 5-bromo-N-{4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl}-2-hydroxybenzamide,
 N-(5-acetyl-4-phenylthiazol-2-yl)-5-bromo-2-hydroxybenzamide,
 N-(5-benzoyl-4-phenylthiazol-2-yl)-5-bromo-2-hydroxybenzamide,
 2-(5-bromo-2-hydroxybenzoyl)amino-4-[(1,1-dimethyl)ethyl]thiazol-5-carboxylic acid ethylester,
 2-(5-bromo-2-hydroxybenzoyl)amino-4-(trifluoromethyl)thiazol-5-carboxylic acid ethylester,
 2-(5-chloro-2-hydroxybenzoyl)amino-4-phenylthiazol-5-carboxylic acid ethylester,
 2-(5-bromo-2-hydroxybenzoyl)amino-4-phenylthiazol-5-carboxylic acid ethylester,
 2-[(4-hydroxybiphenyl)-3-carbonyl]amino-4-phenylthiazol-5-carboxylic acid ethylester,
 2-[(4'-fluoro-4-hydroxybiphenyl)-3-carbonyl]amino-4-phenylthiazol-5-carboxylic acid ethylester,
 2-[(2',4'-difluoro-4-hydroxybiphenyl)-3-carbonyl]amino-4-phenylthiazol-5-carboxylic acid ethylester,
 2-[(4-hydroxy-4'-(trifluoromethyl)biphenyl)-3-carbonyl]amino-4-phenylthiazol-5-carboxylic acid ethylester,
 2-[2-hydroxy-5-(1-pyrrolyl)benzoyl]amino-4-phenylthiazol-5-carboxylic acid ethylester,
 2-[2-hydroxy-5-(1-thienyl)benzoyl]amino-4-phenylthiazol-5-carboxylic acid ethylester,
 2-(5-bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)thiazol-5-carboxylic acid ethylester,
 5-bromo-N-{4-(1,1-dimethyl)ethyl-5-piperidinethiazol-2-yl}-2-hydroxybenzamide,
 5-bromo-N-{4-(1,1-dimethyl)ethyl-5-morpholinethiazol-2-yl}-2-hydroxybenzamide,
 5-bromo-N-{4-(1,1-dimethyl)ethyl-5-(4-methylpiperidin-1-yl)thiazol-2-yl}-2-hydroxybenzamide, and
 5-bromo-N-{4-(1,1-dimethyl)ethyl-5-(4-phenylpiperidin-1-yl)thiazol-2-yl}-2-hydroxybenzamide.

Best Mode for Carrying out the Invention

[0028] The terms used in the present specification have the following meanings.

[0029] As the halogen atom, any of fluorine atom, chlorine atom, bromine atom, or iodine atom may be used unless otherwise specifically referred to.

[0030] Examples of the hydrocarbon group include, for example, an aliphatic hydrocarbon group, an aryl group, an arylene group, an aralkyl group, a bridged cyclic hydrocarbon group, a spiro cyclic hydrocarbon group, and a terpene hydrocarbon.

[0031] Examples of the aliphatic hydrocarbon group include, for example, alkyl group, alkenyl group, alkynyl group,

alkylene group, alkenylene group, alkylidene group and the like which are straight chain or branched chain monovalent or bivalent acyclic hydrocarbon groups; cycloalkyl group, cycloalkenyl group, cycloalkanedieryl group, cycloalkyl-alkyl group, cycloalkylene group, and cycloalkenylene group, which are saturated or unsaturated monovalent or bivalent alicyclic hydrocarbon groups.

[0032] Examples of the alkyl group include, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, 2-methylbutyl, 1-methylbutyl, neopentyl, 1,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 4-methylpentyl, 3-methylpentyl, 2-methylpentyl, 1-methylpentyl, 3,3-dimethylbutyl, 2,2-dimethylbutyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,3-dimethylbutyl, 2-ethylbutyl, 1-ethylbutyl, 1-ethyl-1-methylpropyl, n-heptyl, n-octyl, n-nonyl, n-decyl, n-undecyl, n-dodecyl, n-tridecyl, n-tetradecyl, and n-pentadecyl group, which are C₁ to C₁₅ straight chain or branched chain alkyl groups.

[0033] Examples of the alkenyl group include, for example, vinyl, prop-1-en-1-yl, allyl, isopropenyl, but-1-en-1-yl, but-2-en-1-yl, but-3-en-1-yl, 2-methylprop-2-en-1-yl, 1-methylprop-2-en-1-yl, pent-1-en-1-yl, pent-2-en-1-yl, pent-3-en-1-yl, pent-4-en-1-yl, 3-methylbut-2-en-1-yl, 3-methylbut-3-en-1-yl, hex-1-en-1-yl, hex-2-en-1-yl, hex-3-en-1-yl, hex-4-en-1-yl, hex-5-en-1-yl, 4-methylpent-3-en-1-yl, 4-methylpent-3-en-1-yl, hept-1-en-1-yl, hept-6-en-1-yl, oct-1-en-1-yl, oct-7-en-1-yl, non-1-en-1-yl, non-8-en-1-yl, dec-1-en-1-yl, dec-9-en-1-yl, undec-1-en-1-yl, undec-10-en-1-yl, dodec-1-en-1-yl, dodec-11-en-1-yl, tridec-1-en-1-yl, tridec-12-en-1-yl, tetradec-1-en-1-yl, tetradec-13-en-1-yl, pentadec-1-en-1-yl, and pentadec-14-en-1-yl group, which are C₂ to C₁₅ straight chain or branched chain alkenyl groups.

[0034] Examples of the alkynyl group include, for example, ethynyl, prop-1-yn-1-yl, prop-2-yn-1-yl, but-1-yn-1-yl, but-3-yn-1-yl, 1-methylprop-2-yn-1-yl, pent-1-yn-1-yl, pent-4-yn-1-yl, hex-1-yn-1-yl, hex-5-yn-1-yl, hept-1-yn-1-yl, hept-6-yn-1-yl, oct-1-yn-1-yl, oct-7-yn-1-yl, non-1-yn-1-yl, non-8-yn-1-yl, dec-1-yn-1-yl, dec-9-yn-1-yl, undec-1-yn-1-yl, undec-10-yn-1-yl, dodec-1-yn-1-yl, dodec-11-yn-1-yl, tridec-1-yn-1-yl, tridec-12-yn-1-yl, tetradec-1-yn-1-yl, tetradec-13-yn-1-yl, pentadec-1-yn-1-yl, and pentadec-14-yn-1-yl group, which are C₂ to C₁₅ straight chain or branched chain alkynyl groups.

[0035] Examples of the alkylene group include, for example, methylene, ethylene, ethane-1,1-diyl, propane-1,3-diyl, propane-1,2-diyl, propane-2,2-diyl, butane-1,4-diyl, pentane-1,5-diyl, hexane-1,6-diyl, and 1,1,4,4-tetramethylbutane-1,4-diyl group, which are C₁ to C₈ straight chain or branched chain alkylene groups.

[0036] Examples of the alkenylene group include, for example, ethene-1,2-diyl, propene-1,3-diyl, but-1-ene-1,4-diyl, but-2-ene-1,4-diyl, 2-methylpropene-1,3-diyl, pent-2-ene-1,5-diyl, and hex-3-ene-1,6-diyl group, which are C₁ to C₆ straight chain or branched chain alkenylene groups.

[0037] Examples of the alkylidene group include, for example, methylidene, ethylidene, propylidene, isopropylidene, butylidene, pentylidene, and hexylidene, which are C₁ to C₆ straight chain or branched chain alkylidene groups.

[0038] Examples of the cycloalkyl group include, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl, which are C₃ to C₈ cycloalkyl groups.

[0039] The aforementioned cycloalkyl group may be fused with benzene ring, naphthalene ring and the like, and examples include, for example, 1-indanyl, 2-indanyl, 1,2,3,4-tetrahydronaphthalen-1-yl, and 1,2,3,4-tetrahydronaphthalen-2-yl group.

[0040] Examples of the cycloalkenyl group includes, for example, 2-cyclopropen-1-yl, 2-cyclobuten-1-yl, 2-cyclopenten-1-yl, 3-cyclopenten-1-yl, 2-cyclohexen-1-yl, 3-cyclohexen-1-yl, 1-cyclobuten-1- and 1-cyclopenten-1-yl group, which are C₃ to C₆ cycloalkenyl groups.

[0041] The aforementioned cycloalkenyl group may be fused with benzene ring, naphthalene ring and the like, and examples include, for example, 1-indanyl, 2-indanyl, 1,2,3,4-tetrahydronaphthalen-1-yl, 1,2,3,4-tetrahydronaphthalen-2-yl, 1-indenyl, and 2-indenyl group.

[0042] Examples of the cycloalkanedieryl group include, for example, 2,4-cyclopentadien-1-yl, 2,4-cyclohexanedieryl-1-yl, and 2,5-cyclohexanedieryl-1-yl group, which are C₅ to C₆ cycloalkanedieryl groups.

[0043] The aforementioned cycloalkanedieryl group may be fused with benzene ring, naphthalene ring and the like, and examples of the group include, for example, 1-indenyl and 2-indenyl group.

[0044] Examples of the cycloalkyl-alkyl group include the group in which one hydrogen atom of alkyl group is substituted with cycloalkyl group, and include, for example, cyclopropylmethyl, 1-cyclopropylethyl, 2-cyclopropylethyl, 3-cyclopropylpropyl, 4-cyclopropylbutyl, 5-cyclopropylpentyl, 6-cyclopropylhexyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, cycloheptylmethyl, cyclooctylmethyl, and 6-cyclooctylhexyl group, which are C₄ to C₁₄ cycloalkyl-alkyl groups.

[0045] Examples of the cycloalkylene group include, for example, cyclopropane-1,1-diyl, cyclopropane-1,2-diyl, cyclobutane-1,1-diyl, cyclobutane-1,2-diyl, cyclobutane-1,3-diyl, cyclopentane-1,1-diyl, cyclopentane-1,2-diyl, cyclopentane-1,3-diyl, cyclohexane-1,1-diyl, cyclohexane-1,2-diyl, cyclohexane-1,3-diyl, cyclohexane-1,4-diyl, cycloheptane-1,1-diyl, cycloheptane-1,2-diyl, cyclooctane-1,1-diyl, and cyclooctane-1,2-diyl group, which are C₃ to C₈ cycloalkylene groups.

[0046] Examples of the cycloalkenylene group include, for example, 2-cyclopropene-1,1-diyl, 2-cyclobutene-1,1-diyl, 2-cyclopentene-1,1-diyl, 3-cyclopentene-1,1-diyl, 2-cyclohexene-1,1-diyl, 2-cyclohexene-1,2-diyl, 2-cyclohexene-

1,4-diyl, 3-cyclohexene-1,1-diyl, 1-cyclobutene-1,2-diyl, 1-cyclopentene-1,2-diyl, and 1-cyclohexene-1,2-diyl group, which are C₃ to C₆ cycloalkenylene groups.

[0047] Examples of the aryl group include a monocyclic or a fused polycyclic aromatic hydrocarbon group, and include, for example, phenyl, 1-naphthyl, 2-naphthyl, anthryl, phenanthryl, and acenaphthylenyl group, which are C₆ to C₁₄ aryl groups.

[0048] The aforementioned aryl group may be fused with aforementioned C₃ to C₈ cycloalkyl group, C₃ to C₆ cycloalkenyl group, C₅ to C₆ cycloalkanedienyl group or the like, and the example of the group include, for example, 4-indanyl, 5-indanyl, 1,2,3,4-tetrahydronaphthalen-5-yl, 1,2,3,4-tetrahydronaphthalen-6-yl, 3-acenaphthenyl, 4-acenaphthenyl, inden-4-yl, inden-5-yl, inden-6-yl, inden-7-yl, 4-phenalenyl, 5-phenalenyl, 6-phenalenyl, 7-phenalenyl, 8-phenalenyl, and 9-phenalenyl group.

[0049] Examples of the arylene group include, for example, 1,2-phenylene, 1,3-phenylene, 1,4-phenylene, naphthalene-1,2-diyl, naphthalene-1,3-diyl, naphthalene-1,4-diyl, naphthalene-1,5-diyl, naphthalene-1,6-diyl, naphthalene-1,7-diyl, naphthalene-1,8-diyl, naphthalene-2,3-diyl, naphthalene-2,4-diyl, naphthalene-2,5-diyl, naphthalene-2,6-diyl, naphthalene-2,7-diyl, naphthalene-2,8-diyl, and anthracene-1,4-diyl, which are C₆ to C₁₄ arylene groups.

[0050] Examples of the aralkyl group include the group in which one hydrogen atom of alkyl group is substituted with aryl group, and include, for example, benzyl, 1-naphthylmethyl, 2-naphthylmethyl, anthracenylmethyl, phenanthrenyl methyl, acenaphthylenylmethyl, diphenylmethyl, 1-phenethyl, 2-phenethyl, 1-(1-naphthyl)ethyl, 1-(2-naphthyl)ethyl, 2-(1-naphthyl)ethyl, 2-(2-naphthyl)ethyl, 3-phenylpropyl, 3-(1-naphthyl)propyl, 3-(2-naphthyl)propyl, 4-phenylbutyl, 4-(1-naphthyl)butyl, 4-(2-naphthyl)butyl, 5-phenylpentyl, 5-(1-naphthyl)pentyl, 5-(2-naphthyl)pentyl, 6-phenylhexyl, 6-(1-naphthyl)hexyl, and 6-(2-naphthyl)hexyl group, which are C₇ to C₁₆ aralkyl groups.

[0051] Examples of the bridged cyclic hydrocarbon group include, for example, bicyclo[2.1.0]pentyl, bicyclo[2.2.1]heptyl, bicyclo[2.2.1]octyl, and adamantyl group.

[0052] Examples of the spiro cyclic hydrocarbon group include, for example, spiro[3.4]octyl, and spiro[4.5]decane-1,6-dienyl group.

[0053] Examples of the terpene hydrocarbon include, for example, geranyl, neryl, linalyl, phytyl, menthyl, and bornyl group.

[0054] Examples of the halogenated alkyl group include the group in which one hydrogen atom of alkyl group is substituted with a halogen atom, and include, for example, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, bromomethyl, dibromomethyl, tribromomethyl, iodomethyl, diiodomethyl, triiodomethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, 3,3,3-trifluoropropyl, heptafluoropropyl, heptafluoroisopropyl, nonafluorobutyl, and perfluorohexyl group, which are C₁ to C₆ straight chain or branched chain halogenated alkyl groups substituted with 1 to 13 halogen atoms.

[0055] Examples of the heterocyclic group include, for example, a monocyclic or a fused polycyclic hetero aryl group which comprises at least one atom of 1 to 3 kinds of hetero atoms selected from oxygen atom, sulfur atom, nitrogen atom and the like as ring-constituting atoms (ring forming atoms), and a monocyclic or a fused polycyclic nonaromatic heterocyclic group which comprises at least one atom of 1 to 3 kinds of hetero atoms selected from oxygen atom, sulfur atom, nitrogen atom and the like as ring-constituting atoms (ring forming atoms).

[0056] Examples of the monocyclic heteroaryl group include, for example, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, (1,2,3-oxadiazol)-4-yl, (1,2,3-oxadiazol)-5-yl, (1,2,4-oxadiazol)-3-yl, (1,2,4-oxadiazol)-5-yl, (1,2,5-oxadiazol)-3-yl, (1,2,5-oxadiazol)-4-yl, (1,3,4-oxadiazol)-2-yl, (1,3,4-oxadiazol)-5-yl, furazanyl, (1,2,3-thiadiazol)-4-yl, (1,2,3-thiadiazol)-5-yl, (1,2,4-thiadiazol)-3-yl, (1,2,4-thiadiazol)-5-yl, (1,2,5-thiadiazol)-3-yl, (1,2,5-thiadiazol)-4-yl, (1,3,4-thiadiazol)-2-yl, (1,3,4-thiadiazol)-5-yl, (1H-1,2,3-triazol)-1-yl, (1H-1,2,3-triazol)-4-yl, (1H-1,2,3-triazol)-5-yl, (2H-1,2,3-triazol)-2-yl, (2H-1,2,3-triazol)-4-yl, (1H-1,2,4-triazol)-1-yl, (1H-1,2,4-triazol)-3-yl, (1H-1,2,4-triazol)-5-yl, (4H-1,2,4-triazol)-3-yl, (4H-1,2,4-triazol)-4-yl, (1H-tetrazol)-1-yl, (1H-tetrazol)-5-yl, (2H-tetrazol)-2-yl, (2H-tetrazol)-5-yl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 3-pyridazinyl, 4-pyridazinyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 2-pyrazinyl, (1,2,3-triazin)-4-yl, (1,2,3-triazin)-5-yl, (1,2,4-triazin)-3-yl, (1,2,4-triazin)-5-yl, (1,2,4-triazin)-6-yl, (1,3,5-triazin)-2-yl, 1-azepinyl, 2-azepinyl, 3-azepinyl, 4-azepinyl, (1,4-oxazepin)-2-yl, (1,4-oxazepin)-3-yl, (1,4-oxazepin)-5-yl, (1,4-oxazepin)-6-yl, (1,4-oxazepin)-7-yl, (1,4-thiazepin)-2-yl, (1,4-thiazepin)-3-yl, (1,4-thiazepin)-5-yl, (1,4-thiazepin)-6-yl, and (1,4-thiazepin)-7-yl group, which are 5- to 7-membered monocyclic heteroaryl groups.

[0057] Examples of the fused polycyclic heteroaryl group include, for example, 2-benzofuranyl, 3-benzofuranyl, 4-benzofuranyl, 5-benzofuranyl, 6-benzofuranyl, 7-benzofuranyl, 1-isobenzofuranyl, 4-isobenzofuranyl, 5-isobenzofuranyl, 2-benzo[b]thienyl, 3-benzo[b]thienyl, 4-benzo[b]thienyl, 5-benzo[b]thienyl, 6-benzo[b]thienyl, 7-benzo[b]thienyl, 1-benzo[c]thienyl, 4-benzo[c]thienyl, 5-benzo[c]thienyl, 1-indolyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl, (2H-isoidol)-1-yl, (2H-isoidol)-2-yl, (2H-isoidol)-4-yl, (2H-isoidol)-5-yl, (1H-indazol)-1-yl, (1H-indazol)-4-yl, (1H-indazol)-5-yl, (1H-indazol)-6-yl, (1H-indazol)-7-yl, (2H-indazol)-1-yl, (2H-indazol)-2-yl, (2H-indazol)-4-yl, (2H-indazol)-5-yl, 2-benzoxazolyl, 2-benzoxazolyl, 4-benzoxazolyl, 5-benzoxazolyl, 6-benzox-

azolyl, 7-benzoxazolyl, (1,2-benzisoxazol)-3-yl, (1,2-benzisoxazol)-4-yl, (1,2-benzisoxazol)-5-yl, (1,2-benzisoxazol)-6-yl, (1,2-benzisoxazol)-7-yl, (2,1-benzisoxazol)-3-yl, (2,1-benzisoxazol)-4-yl, (2,1-benzisoxazol)-5-yl, (2,1-benzisoxazol)-6-yl, (2,1-benzisoxazol)-7-yl, 2-benzothiazolyl, 4-benzothiazolyl, 5-benzothiazolyl, 6-benzothiazolyl, 7-benzothiazolyl, (1,2-benzisothiazol)-3-yl, (1,2-benzisothiazol)-4-yl, (1,2-benzisothiazol)-5-yl, (1,2-benzisothiazol)-6-yl, (1,2-benzisothiazol)-7-yl, (2,1-benzisothiazol)-3-yl, (2,1-benzisothiazol)-4-yl, (2,1-benzisothiazol)-5-yl, (2,1-benzisothiazol)-6-yl, (2,1-benzisothiazol)-7-yl, (1,2,3-benzoxadiazol)-4-yl, (1,2,3-benzoxadiazol)-5-yl, (1,2,3-benzoxadiazol)-6-yl, (1,2,3-benzoxadiazol)-7-yl, (2,1,3-benzoxadiazol)-4-yl, (2,1,3-benzoxadiazol)-5-yl, (1,2,3-benzothiadiazol)-4-yl, (1,2,3-benzothiadiazol)-5-yl, (1,2,3-benzothiadiazol)-6-yl, (1,2,3-benzothiadiazol)-7-yl, (2,1,3-benzothiadiazol)-4-yl, (2,1,3-benzothiadiazol)-5-yl, (1H-benzotriazol)-1-yl, (1H-benzotriazol)-4-yl, (1H-benzotriazol)-5-yl, (1H-benzotriazol)-6-yl, (1H-benzotriazol)-7-yl, (2H-benzotriazol)-2-yl, (2H-benzotriazol)-4-yl, (2H-benzotriazol)-5-yl, 2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-quinolyl, 7-quinolyl, 8-quinolyl, 1-isoquinolyl, 3-isoquinolyl, 4-isoquinolyl, 5-isoquinolyl, 6-isoquinolyl, 7-isoquinolyl, 8-isoquinolyl, 3-cinnolyl, 4-cinnolyl, 5-cinnolyl, 6-cinnolyl, 7-cinnolyl, 8-cinnolyl, 2-quinazolyl, 4-quinazolyl, 5-quinazolyl, 6-quinazolyl, 7-quinazolyl, 8-quinazolyl, 2-quinoxalyl, 5-quinoxalyl, 6-quinoxalyl, 1-phthalazyl, 5-phthalazyl, 6-phthalazyl, 2-naphthyridyl, 3-naphthyridyl, 4-naphthyridyl, 2-purlyl, 6-purlyl, 7-purlyl, 8-purlyl, 2-pteridyl, 4-pteridyl, 6-pteridyl, 7-pteridyl, 1-carbazolyl, 2-carbazolyl, 3-carbazolyl, 4-carbazolyl, 9-carbazolyl, 2-(α -carbollyl), 3-(α -carbollyl), 4-(α -carbollyl), 5-(α -carbollyl), 6-(α -carbollyl), 7-(α -carbollyl), 8-(α -carbollyl), 9-(α -carbollyl), 1-(β -carbollyl), 3-(β -carbollyl), 4-(β -carbollyl), 5-(β -carbollyl), 6-(β -carbollyl), 7-(β -carbollyl), 8-(β -carbollyl), 9-(β -carbollyl), 1-(γ -carbollyl), 2-(γ -carbollyl), 4-(γ -carbollyl), 5-(γ -carbollyl), 6-(γ -carbollyl), 7-(γ -carbollyl), 8-(γ -carbollyl), 9-(γ -carbollyl), 1-acridyl, 2-acridyl, 3-acridyl, 4-acridyl, 9-acridyl, 1-phenoxazyl, 2-phenoxazyl, 3-phenoxazyl, 4-phenoxazyl, 10-phenoxazyl, 1-phenothiazyl, 2-phenothiazyl, 3-phenothiazyl, 4-phenothiazyl, 10-phenothiazyl, 1-phenazyl, 2-phenazyl, 1-phenanthridyl, 2-phenanthridyl, 3-phenanthridyl, 4-phenanthridyl, 6-phenanthridyl, 7-phenanthridyl, 8-phenanthridyl, 9-phenanthridyl, 10-phenanthridyl, 2-phenanthrolyl, 3-phenanthrolyl, 4-phenanthrolyl, 5-phenanthrolyl, 6-phenanthrolyl, 7-phenanthrolyl, 8-phenanthrolyl, 9-phenanthrolyl, 10-phenanthrolyl, 1-thianthrenyl, 2-thianthrenyl, 1-indolizyl, 2-indolizyl, 3-indolizyl, 5-indolizyl, 6-indolizyl, 7-indolizyl, 8-indolizyl, 1-phenoxathiyl, 2-phenoxathiyl, 3-phenoxathiyl, 4-phenoxathiyl, thieno[2,3-b]furyl, pyrrolo[1,2-b]pyridazyl, pyrazolo[1,5-a]pyridyl, imidazo[11,2-a]pyridyl, imidazo[1,5-a]pyridyl, imidazo[1,2-b]pyridazyl, imidazo[1,2-a]pyrimidyl, 1,2,4-triazolo[4,3-a]pyridyl, and 1,2,4-triazolo[4,3-a]pyridazyl group, which are 8- to 14-membered fused polycyclic heteroaryl groups.

[0058] Examples of the monocyclic nonaromatic heterocyclic group include, for example, 1-aziridyl, 1-azetidyl, 1-pyrrolidyl, 2-pyrrolidyl, 3-pyrrolidyl, 2-tetrahydrofuryl, 3-tetrahydrofuryl, thiolanyl, 1-imidazolidyl, 2-imidazolidyl, 4-imidazolidyl, 1-pyrazolidyl, 3-pyrazolidyl, 4-pyrazolidyl, 1-(2-pyrrolinyl), 1-(2-imidazolyl), 2-(2-imidazolyl), 1-(2-pyrazolyl), 3-(2-pyrazolyl), piperidino, 2-piperidyl, 3-piperidyl, 4-piperidyl, 1-homopiperidyl, 2-tetrahydropyranyl, morpholino, (thiomorpholin)-4-yl, 1-piperazyl, and 1-homopiperazyl group, which are 3- to 7-membered saturated or unsaturated monocyclic nonaromatic heterocyclic groups.

[0059] Examples of the fused polycyclic nonaromatic heterocyclic group include, for example, 2-quinuclidyl, 2-chromanyl, 3-chromanyl, 4-chromanyl, 5-chromanyl, 6-chromanyl, 7-chromanyl, 8-chromanyl, 1-isochromanyl, 3-isochromanyl, 4-isochromanyl, 5-isochromanyl, 6-isochromanyl, 7-isochromanyl, 8-isochromanyl, 2-thiochromanyl, 3-thiochromanyl, 4-thiochromanyl, 5-thiochromanyl, 6-thiochromanyl, 7-thiochromanyl, 8-thiochromanyl, 1-isothiochromanyl, 3-isothiochromanyl, 4-isothiochromanyl, 5-isothiochromanyl, 6-isothiochromanyl, 7-isothiochromanyl, 8-isothiochromanyl, 1-indolyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl, 1-isindolyl, 2-isindolyl, 4-isindolyl, 5-isindolyl, 2-(4H-chromenyl), 3-(4H-chromenyl), 4-(4H-chromenyl), 5-(4H-chromenyl), 6-(4H-chromenyl), 7-(4H-chromenyl), 8-(4H-chromenyl), 1-isochromenyl, 3-isochromenyl, 4-isochromenyl, 5-isochromenyl, 6-isochromenyl, 7-isochromenyl, 8-isochromenyl, 1-(1H-pyrrolidyl), 2-(1H-pyrrolidyl), 3-(1H-pyrrolidyl), 5-(1H-pyrrolidyl), 6-(1H-pyrrolidyl), and 7-(1H-pyrrolidyl) group, which are 8- to 10-membered saturated or unsaturated fused polycyclic nonaromatic heterocyclic groups.

[0060] Among the aforementioned heterocyclic groups, a monocyclic or a fused polycyclic heteroaryl groups which may have 1 to 3 kinds of hetero atoms selected from oxygen atom, sulfur atom, nitrogen atom and the like, in addition to the nitrogen atom that has the bond, as ring-constituting atoms (ring forming atoms), and a monocyclic or a fused polycyclic nonaromatic heterocyclic groups which may have 1 to 3 kinds of hetero atoms selected from oxygen atom, sulfur atom, nitrogen atom and the like, in addition to the nitrogen atom that has the bond, as ring-constituting atoms (ring forming atoms) are referred to as "cyclic amino groups". Examples include, for example, 1-pyrrolidyl, 1-imidazolidyl, 1-pyrazolidyl, 1-oxazolidyl, 1-thiazolidyl, piperidino, morpholino, 1-piperazyl, thiomorpholin-4-yl, 1-homopiperidyl, 1-homopiperazyl, 2-pyrrolin-1-yl, 2-imidazol-1-yl, 2-pyrazol-1-yl, 1-indolyl, 2-isindolyl, 1,2,3,4-tetrahydroquinolin-1-yl, 1,2,3,4-tetrahydroisoquinolin-2-yl, 1-pyrrolyl, 1-imidazolyl, 1-pyrazolyl, 1-indolyl, 1-indazolyl, and 2-isindolyl group.

[0061] Examples of the hydrocarbon-oxy group include the group in which hydrogen atom of hydroxy group is substituted with hydrocarbon group, and examples of the hydrocarbon include similar groups to the aforementioned hydro-

drocarbon. Examples of the hydrocarbon-oxy group include, for example, alkoxy group (alkyl-oxy group), alkenyl-oxy group, alkynyl-oxy group, cycloalkyl-oxy group, cycloalkyl-alkyl-oxy group and the like which are aliphatic hydrocarbon-oxy groups, aryl-oxy group, aralkyl-oxy group, and alkylene-dioxy group.

[0062] Examples of the alkoxy(alkyl-oxy group) include, for example, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, n-pentyloxy, isopentyloxy, 2-methylbutoxy, 1-methylbutoxy, neopentyloxy, 1,2-dimethylpropoxy, 1-ethylpropoxy, n-hexyloxy, 4-methylpentyloxy, 3-methylpentyloxy, 2-methylpentyloxy, 1-methylpentyloxy, 3,3-dimethylbutoxy, 2,2-dimethylbutoxy, 1,1-dimethylbutoxy, 1,2-dimethylbutoxy, 1,3-dimethylbutoxy, 2,3-dimethylbutoxy, 2-ethylbutoxy, 1-ethylbutoxy, 1-ethyl-1-methylpropoxy, n-heptyloxy, n-octyloxy, n-nonyloxy, n-decyloxy, n-undecyloxy, n-dodecyloxy, n-tridecyloxy, n-tetradecyloxy, and n-pentadecyloxy group, which are C₁ to C₁₅ straight chain or branched chain alkoxy groups.

[0063] Examples of the alkenyl-oxy group include, for example, vinyloxy, (prop-1-en-1-yl)oxy, allyloxy, isopropenyloxy, (but-1-en-1-yl)oxy, (but-2-en-1-yl)oxy, (but-3-en-1-yl)oxy, (2-methylprop-2-en-1-yl)oxy, (1-methylprop-2-en-1-yl)oxy, (pent-1-en-1-yl)oxy, (pent-2-en-1-yl)oxy, (pent-3-en-1-yl)oxy, (pent-4-en-1-yl)oxy, (3-methylbut-2-en-1-yl)oxy, (3-methylbut-3-en-1-yl)oxy, (hex-1-en-1-yl)oxy, (hex-2-en-1-yl)oxy, (hex-3-en-1-yl)oxy, (hex-4-en-1-yl)oxy, (hex-5-en-1-yl)oxy, (4-methylpent-3-en-1-yl)oxy, (4-methylpent-3-en-1-yl)oxy, (hept-1-en-1-yl)oxy, (hept-6-en-1-yl)oxy, (oct-1-en-1-yl)oxy, (oct-7-en-1-yl)oxy, (non-1-en-1-yl)oxy, (non-8-en-1-yl)oxy, (dec-1-en-1-yl)oxy, (dec-9-en-1-yl)oxy, (undec-1-en-1-yl)oxy, (undec-10-en-1-yl)oxy, (dodec-1-en-1-yl)oxy, (dodec-11-en-1-yl)oxy, (tridec-1-en-1-yl)oxy, (tridec-12-en-1-yl)oxy, (tetradec-1-en-1-yl)oxy, (tetradec-13-en-1-yl)oxy, (pentadec-1-en-1-yl)oxy, and (pentadec-14-en-1-yl)oxy group, which are C₂ to C₁₅ straight chain or branched chain alkenyl-oxy groups.

[0064] Examples of the alkynyl-oxy group include, for example, ethynyloxy, (prop-1-yn-1-yl)oxy, (prop-2-yn-1-yl)oxy, (but-1-yn-1-yl)oxy, (but-3-yn-1-yl)oxy, (1-methylprop-2-yn-1-yl)oxy, (pent-1-yn-1-yl)oxy, (pent-4-yn-1-yl)oxy, (hex-1-yn-1-yl)oxy, (hex-5-yn-1-yl)oxy, (hept-1-yn-1-yl)oxy, (hept-6-yn-1-yl)oxy, (oct-1-yn-1-yl)oxy, (oct-7-yn-1-yl)oxy, (non-1-yn-1-yl)oxy, (non-8-yn-1-yl)oxy, (dec-1-yn-1-yl)oxy, (dec-9-yn-1-yl)oxy, (undec-1-yn-1-yl)oxy, (undec-10-yn-1-yl)oxy, (dodec-1-yn-1-yl)oxy, (dodec-11-yn-1-yl)oxy, (tridec-1-yn-1-yl)oxy, (tridec-12-yn-1-yl)oxy, (tetradec-1-yn-1-yl)oxy, (tetradec-13-yn-1-yl)oxy, (pentadec-1-yn-1-yl)oxy, and (pentadec-14-yn-1-yl)oxy group, which are C₂ to C₁₅ straight chain or branched chain alkynyl-oxy groups.

[0065] Examples of the cycloalkyl-oxy group include, for example, cyclopropoxy, cyclobutoxy, cyclopentyloxy, cyclohexyloxy, cycloheptyloxy, and cyclooctyloxy group, which are C₃ to C₈ cycloalkyl-oxy groups.

[0066] Examples of the cycloalkyl-alkyl-oxy group include, for example, cyclopropylmethoxy, 1-cyclopropylethoxy, 2-cyclopropylethoxy, 3-cyclopropylpropoxy, 4-cyclopropylbutoxy, 5-cyclopropylpentyloxy, 6-cyclopropylhexyloxy, cyclobutylmethoxy, cyclopentylmethoxy, cyclobutylmethoxy, cyclopentylmethoxy, cyclohexylmethoxy, 2-cyclohexylethoxy, 3-cyclohexylpropoxy, 4-cyclohexylbutoxy, cycloheptylmethoxy, cyclooctylmethoxy, and 6-cyclooctylhexyloxy group, which are C₄ to C₁₄ cycloalkyl-alkyl-oxy groups.

[0067] Examples of the aryl-oxy group include, for example, phenoxy, 1-naphthyloxy, 2-naphthyloxy, anthryloxy, phenanthryloxy, and acenaphthylenyloxy group, which are C₆ to C₁₄ aryl-oxy groups.

[0068] Examples of the aralkyl-oxy group include, for example, benzyloxy, 1-naphthylmethoxy, 2-naphthylmethoxy, anthracenylmethoxy, phenanthrenylmethoxy, acenaphthylenyloxy, diphenylmethoxy, 1-phenylethoxy, 2-phenylethoxy, 1-(1-naphthyl)ethoxy, 1-(2-naphthyl)ethoxy, 2-(1-naphthyl)ethoxy, 2-(2-naphthyl)ethoxy, 3-phenylpropoxy, 3-(1-naphthyl)propoxy, 3-(2-naphthyl)propoxy, 4-phenylbutoxy, 4-(1-naphthyl)butoxy, 4-(2-naphthyl)butoxy, 5-phenylpentyloxy, 5-(1-naphthyl)pentyloxy, 5-(2-naphthyl)pentyloxy, 6-phenylhexyloxy, 6-(1-naphthyl)hexyloxy, and 6-(2-naphthyl)hexyloxy group, which are C₇ to C₁₆ aralkyl-oxy groups.

[0069] Examples of the alkylenedioxy group include, for example, methylenedioxy, ethylenedioxy, 1-methylmethylenedioxy, and 1,1-dimethylmethylenedioxy group.

[0070] Examples of the halogenated alkoxy group(halogenated alkyl-oxy group) include the group in which hydrogen atom of hydroxy group is substituted with halogenated alkyl group, and include, for example, fluoromethoxy, difluoromethoxy, chloromethoxy, bromomethoxy, iodomethoxy, trifluoromethoxy, trichloromethoxy, 2,2,2-trifluoroethoxy, pentafluoroethoxy, 3,3,3-trifluoropropoxy, heptafluoropropoxy, heptafluoroisopropoxy, nonafluorobutoxy, and perfluorohexyloxy group, which are C₁ to C₆ straight chain or branched chain halogenated alkoxy groups substituted with 1 to 13 halogen atoms.

[0071] Examples of the heterocyclic-oxy group include the group in which hydrogen atom of hydroxy group is substituted with the heterocyclic group, and examples of the heteroring include similar groups to the aforementioned heterocyclic group. Examples of the heteroring-oxy group include, for example, a monocyclic heteroaryl-oxy group, a fused polycyclic heteroaryl-oxy group, a monocyclic nonaromatic heteroring-oxy group, a fused polycyclic nonaromatic heteroring-oxy group and the like.

[0072] Examples of the monocyclic heteroaryl-oxy group include, for example, 3-thienyloxy, (isoxazol-3-yl)oxy, (thiazol-4-yl)oxy, 2-pyridyloxy, 3-pyridyloxy, 4-pyridyloxy, and (pyrimidin-4-yl)oxy group.

[0073] Examples of the fused polycyclic heteroaryl-oxy group include, 5-indolyloxy, (benzimidazol-2-yl)oxy, 2-quinolyloxy, 3-quinolyloxy, and 4-quinolyloxy group.

[0074] Examples of the monocyclic nonaromatic heteroring-oxy group include, for example, 3-pyrrolidinyloxy, and 4-piperidinyloxy group.

[0075] Examples of the fused polycyclic non-aromatic heteroring-oxy group include, for example, 3-indolinyloxy, and 4-chromanyloxy group.

[0076] Examples of the hydrocarbon-sulfanyl group include the group in which hydrogen atom of sulfanyl group is substituted with a hydrocarbon group, and examples of the hydrocarbon include similar groups to the aforementioned hydrocarbon. Examples of the hydrocarbon-sulfanyl group include, for example, alkyl-sulfanyl group, alkenyl-sulfanyl group, alkynyl-sulfanyl group, cycloalkyl-sulfanyl group, cycloalkyl-alkyl-sulfanyl group and the like which are aliphatic hydrocarbon-sulfanyl groups, aryl-sulfanyl group, and aralkyl-sulfanyl group.

[0077] Examples of the alkyl-sulfanyl group include, for example, methylsulfanyl, ethylsulfanyl, n-propylsulfanyl, isopropylsulfanyl, n-butylsulfanyl, isobutylsulfanyl, sec-butylsulfanyl, tert-butylsulfanyl, n-pentylsulfanyl, isopentylsulfanyl, (2-methylbutyl)sulfanyl, (1-methylbutyl)sulfanyl, neopentylsulfanyl, (1,2-dimethylpropyl)sulfanyl, (1-ethylpropyl)sulfanyl, n-hexylsulfanyl, (4-methylpentyl)sulfanyl, (3-methylpentyl)sulfanyl, (2-methylpentyl)sulfanyl, (1-methylpentyl)sulfanyl, (3,3-dimethylbutyl)sulfanyl, (2,2-dimethylbutyl)sulfanyl, (1,1-dimethylbutyl)sulfanyl, (1,2-dimethylbutyl)sulfanyl, (1,3-dimethylbutyl)sulfanyl, (2,3-dimethylbutyl)sulfanyl, (2-ethylbutyl)sulfanyl, (1-ethylbutyl)sulfanyl, (1-ethyl-1-methylpropyl)sulfanyl, n-heptylsulfanyl, n-octylsulfanyl, n-nonylsulfanyl, n-decylsulfanyl, n-undecylsulfanyl, n-dodecylsulfanyl, n-tridecylsulfanyl, n-tetradecylsulfanyl, and n-pentadecylsulfanyl group, which are C₁ to C₁₅ straight chain or branched chain alkyl-sulfanyl groups.

[0078] Examples of the alkenyl-sulfanyl group include, for example, vinylsulfanyl, (prop-1-en-1-yl)sulfanyl, allylsulfanyl, isopropenylsulfanyl, (but-1-en-1-yl)sulfanyl, (but-2-en-1-yl)sulfanyl, (but-3-en-1-yl)sulfanyl, (2-methylprop-2-en-1-yl)sulfanyl, (1-methylprop-2-en-1-yl)sulfanyl, (pent-1-en-1-yl)sulfanyl, (pent-2-en-1-yl)sulfanyl, (pent-3-en-1-yl)sulfanyl, (pent-4-en-1-yl)sulfanyl, (3-methylbut-2-en-1-yl)sulfanyl, (3-methylbut-3-en-1-yl)sulfanyl, (hex-1-en-1-yl)sulfanyl, (hex-2-en-1-yl)sulfanyl, (hex-3-en-1-yl)sulfanyl, (hex-4-en-1-yl)sulfanyl, (hex-5-en-1-yl)sulfanyl, (4-methylpent-3-en-1-yl)sulfanyl, (4-methylpent-3-en-1-yl)sulfanyl, (hept-1-en-1-yl)sulfanyl, (hept-6-en-1-yl)sulfanyl, (oct-1-en-1-yl)sulfanyl, (oct-7-en-1-yl)sulfanyl, (non-1-en-1-yl)sulfanyl, (non-8-en-1-yl)sulfanyl, (dec-1-en-1-yl)sulfanyl, (dec-9-en-1-yl)sulfanyl, (undec-1-en-1-yl)sulfanyl, (undec-10-en-1-yl)sulfanyl, (dodec-1-en-1-yl)sulfanyl, (dodec-11-en-1-yl)sulfanyl, (tridec-1-en-1-yl)sulfanyl, (tridec-12-en-1-yl)sulfanyl, (tetradec-1-en-1-yl)sulfanyl, (tetradec-13-en-1-yl)sulfanyl, (pentadec-1-en-1-yl)sulfanyl, and (pentadec-14-en-1-yl)sulfanyl, which are C₂ to C₁₅ straight chain or branched chain alkenyl-sulfanyl groups.

[0079] Examples of the alkynyl-sulfanyl group include, for example, ethynylsulfanyl, (prop-1-yn-1-yl)sulfanyl, (prop-2-yn-1-yl)sulfanyl, (but-1-yn-1-yl)sulfanyl, (but-3-yn-1-yl)sulfanyl, (1-methylprop-2-yn-1-yl)sulfanyl, (pent-1-yn-1-yl)sulfanyl, (pent-4-yn-1-yl)sulfanyl, (hex-1-yn-1-yl)sulfanyl, (hex-5-yn-1-yl)sulfanyl, (hept-1-yn-1-yl)sulfanyl, (hept-6-yn-1-yl)sulfanyl, (oct-1-yn-1-yl)sulfanyl, (oct-7-yn-1-yl)sulfanyl, (non-1-yn-1-yl)sulfanyl, (non-8-yn-1-yl)sulfanyl, (dec-1-yn-1-yl)sulfanyl, (dec-9-yn-1-yl)sulfanyl, (undec-1-yn-1-yl)sulfanyl, (undec-10-yn-1-yl)sulfanyl, (dodec-1-yn-1-yl)sulfanyl, (dodec-11-yn-1-yl)sulfanyl, (tridec-1-yn-1-yl)sulfanyl, (tridec-12-yn-1-yl)sulfanyl, (tetradec-1-yn-1-yl)sulfanyl, (tetradec-13-yn-1-yl)sulfanyl, (pentadec-1-yn-1-yl)sulfanyl, and (pentadec-14-yn-1-yl)sulfanyl, which are C₂ to C₁₅ straight chain or branched chain alkynyl-sulfanyl groups.

[0080] Examples of the cycloalkyl-sulfanyl group include, for example, cyclopropylsulfanyl, cyclobutylsulfanyl, cyclopentylsulfanyl, cyclohexylsulfanyl, cycloheptylsulfanyl, and cyclooctylsulfanyl group, which are C₃ to C₈ cycloalkyl-sulfanyl groups.

[0081] Examples of the cycloalkyl-alkyl-sulfanyl group include, for example, (cyclopropylmethyl)sulfanyl, (1-cyclopropylethyl)sulfanyl, (2-cyclopropylethyl)sulfanyl, (3-cyclopropylpropyl)sulfanyl, (4-cyclopropylbutyl)sulfanyl, (5-cyclopropylpentyl)sulfanyl, (6-cyclopropylhexyl)sulfanyl, (cyclobutylmethyl)sulfanyl, (cyclopentylmethyl)sulfanyl, (cyclobutylmethyl)sulfanyl, (cyclopentylmethyl)sulfanyl, (cyclohexylmethyl)sulfanyl, (2-cyclohexylethyl)sulfanyl, (3-cyclohexylpropyl)sulfanyl, (4-cyclohexylbutyl)sulfanyl, (cycloheptylmethyl)sulfanyl, (cyclooctylmethyl)sulfanyl, and (6-cyclooctylhexyl)sulfanyl group, which are C₄ to C₁₄ cycloalkyl-alkyl-sulfanyl groups.

[0082] Examples of the aryl-sulfanyl group include, for example, phenylsulfanyl, 1-naphthylsulfanyl, 2-naphthylsulfanyl, anthrylsulfanyl, fenanthrilysulfanyl, and acenaphthylenylsulfanyl group, which are C₆ to C₁₄ aryl-sulfanyl groups.

[0083] Examples of the aralkyl-sulfanyl group include, for example, benzylsulfanyl, (1-naphthylmethyl)sulfanyl, (2-naphthylmethyl)sulfanyl, (anthracenylmethyl)sulfanyl, (phenanthrenylmethyl)sulfanyl, (acenaphthylenylmethyl)sulfanyl, (diphenylmethyl)sulfanyl, (1-phenethyl)sulfanyl, (2-phenethyl)sulfanyl, (1-(1-naphthyl)ethyl)sulfanyl, (1-(2-naphthyl)ethyl)sulfanyl, (2-(1-naphthyl)ethyl)sulfanyl, (2-(2-naphthyl)ethyl)sulfanyl, (3-phenylpropyl)sulfanyl, (3-(1-naphthyl)propyl)sulfanyl, (3-(2-naphthyl)propyl)sulfanyl, (4-phenylbutyl)sulfanyl, (4-(1-naphthyl)butyl)sulfanyl, (4-(2-naphthyl)butyl)sulfanyl, (5-phenylpentyl)sulfanyl, (5-(1-naphthyl)pentyl)sulfanyl, (5-(2-naphthyl)pentyl)sulfanyl, (6-phenylhexyl)sulfanyl, (6-(1-naphthyl)hexyl)sulfanyl, and (6-(2-naphthyl)hexyl)sulfanyl group, which are C₇ to C₁₆ aralkyl-sulfanyl groups.

[0084] Examples of the halogenated alkyl-sulfanyl group include the group in which hydrogen atom of sulfanyl group is substituted with a halogenated alkyl group, and include, for example, (fluoromethyl)sulfanyl, (chloromethyl)sulfanyl,

(bromomethyl)sulfanyl, (iodomethyl)sulfanyl, (difluoromethyl)sulfanyl, (trifluoromethyl)sulfanyl, (trichloromethyl)sulfanyl, (2,2,2-trifluoroethyl)sulfanyl, (pentafluoroethyl)sulfanyl, (3,3,3-trifluoropropyl)sulfanyl, (heptafluoropropyl)sulfanyl, (heptafluoroisopropyl)sulfanyl, (nonafluorobutyl)sulfanyl, and (perfluorohexyl)sulfanyl group, which are C₁ to C₆ straight chain or branched chain halogenated alkyl-sulfanyl groups substituted with 1 to 13 halogen atoms.

[0085] Examples of the heterocyclic-sulfanyl group include the group in which hydrogen atom of sulfanyl group is substituted with the heterocyclic group, and examples of the heteroring include similar groups to the aforementioned heterocyclic group. Examples of the heteroring-sulfanyl group include, for example, monocyclic heteroaryl-sulfanyl group; fused polycyclic heteroaryl-sulfanyl group, monocyclic nonaromatic heteroring-sulfanyl group, and fused polycyclic nonaromatic heteroring-sulfanyl group.













[0086] Examples of the monocyclic heteroaryl-sulfanyl group include, for example, (imidazol-2-yl)sulfanyl, (1,2,4-triazol-2-yl)sulfanyl, (pyridin-2-yl)sulfanyl, (pyridin-4-yl)sulfanyl, and (pyrimidin-2-yl)sulfanyl group.










[0087] Examples of the fused polycyclic heteroaryl-sulfanyl group include, for example, (benzimidazol-2-yl)sulfanyl, (quinolin-2-yl)sulfanyl, and (quinolin-4-yl)sulfanyl group.

[0088] Examples of the monocyclic non-aromatic heteroring-sulfanyl group include, for example, (3-pyrrolidinyl)sulfanyl group and (4-piperidinyl)sulfanyl group.

[0089] Examples of the fused polycyclic nonaromatic heteroring-sulfanyl group include, for example, (3-indoliny) sulfanyl and (4-chromanyl)sulfanyl group.

[0090] Examples of the acyl group include, for example, formyl group, glyoxyloyl group, and thioformyl group, and the group represented by the following formulas:





wherein R^{a1} and R^{b1} may be the same or different and each represents a hydrocarbon group or heterocyclic group, or R^{a1} and R^{b1} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group.

[0091] In the definition of the aforementioned acyl group, among the groups represented by the formula (ω -1A), those groups in which R^{a1} is a hydrocarbon group are referred to as "hydrocarbon-carbonyl groups" whose examples include, for example, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, lauroyl, myristoyl, palmitoyl, acryloyl, propioloyl, methacryloyl, crotonoyl, isocrotonoyl, cyclohexylcarbonyl, cyclohexylmethylcarbonyl, benzoyl, 1-naphthoyl, 2-naphthoyl, and phenylacetyl group. Those groups in which R^{a1} is a heterocyclic group are referred to as "heteroring-carbonyl" groups whose examples include, for example, 2-thenoyl, 3-furoyl, nicotinoyl, and isonicotinoyl group.

[0092] Among the groups represented by the formula (ω -2A), those groups in which R^{a1} is a hydrocarbon group are referred to as "hydrocarbon-oxy-carbonyl groups" whose examples include, for example, methoxycarbonyl, ethoxycarbonyl, phenoxycarbonyl, and benzyloxycarbonyl, and those groups in which R^{a1} is a heterocyclic group are referred to as "heteroring-oxy-carbonyl groups" whose example includes, for example, 3-pyridyloxycarbonyl group.

[0093] Among the groups represented by the formula (ω -3A), those groups in which R^{a1} is a hydrocarbon group are referred to as "hydrocarbon-carbonyl-carbonyl groups" whose example includes, for example, pyruvoyl group, and those groups in which R^{a1} is a heterocyclic group are referred to as "heteroring-carbonyl-carbonyl groups".

[0094] Among the groups represented by the formula (ω -4A), those groups in which R^{a1} is a hydrocarbon group are referred to as "hydrocarbon-oxy-carbonyl-carbonyl groups" whose examples include, for example, methoxalyl and ethoxalyl groups, and those groups in which R^{a1} is a heterocyclic group are referred to as "heteroring-oxy-carbonyl-carbonyl groups".

[0095] Among the groups represented by the formula (ω -5A), those groups in which R^{a1} is a hydrocarbon group are referred to as "hydrocarbon-sulfanyl-carbonyl groups", and those groups in which R^{a1} is a heterocyclic group are referred to as "heteroring-sulfanyl-carbonyl groups".

[0096] Among the groups represented by the formula (ω -6A), those groups in which R^{a1} is a hydrocarbon group are referred to as "hydrocarbon-thiocarbonyl groups", and those groups in which R^{a1} is a heterocyclic group are referred

to as "heteroring-thiocarbonyl groups".

[0097] Among the groups represented by the formula (ω -7A), those groups in which R^{a1} is a hydrocarbon group are referred to as "hydrocarbon-oxy-thiocarbonyl groups", and those groups in which R^{a1} is a heterocyclic group are referred to as "heteroring-oxy-thiocarbonyl groups".

[0098] Among the groups represented by the formula (ω -8A), those groups in which R^{a1} is a hydrocarbon group are referred to as "hydrocarbon-sulfanyl-thiocarbonyl groups", and those groups in which R^{a1} is a heterocyclic group are referred to as "heteroring-sulfanyl-thiocarbonyl groups".

[0099] Among the groups represented by the formula (ω -9A), those groups in which R^{a1} is a hydrocarbon group are referred to as "N-hydrocarbon-carbamoyl groups" whose example includes, for example, N-methylcarbamoyl group, and those groups in which R^{a1} is a heterocyclic group are referred to as "N-heteroring-carbamoyl groups".

[0100] Among the groups represented by the formula (ω -10A), those groups in which both R^{a1} and R^{b1} are hydrocarbon groups are referred to as "N,N-dihydrocarbon-carbamoyl groups" whose example includes, for example, N,N-dimethylcarbamoyl group, those groups in which both R^{a1} and R^{b1} are heterocyclic groups are referred to as "N,N-di(heteroring)-carbamoyl groups", those groups in which R^{a1} is a hydrocarbon group and R^{b1} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-substituted carbamoyl groups", and those groups in which R^{a1} and R^{b1} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclic amino-carbonyl groups" whose example includes, for example, morpholino-carbonyl.

[0101] Among the groups represented by the formula (ω -11A), those groups in which R^{a1} is a hydrocarbon group are referred to as "N-hydrocarbon-thiocarbamoyl groups", and those groups in which R^{a1} is a heterocyclic group are referred to as "N-heteroring-thiocarbamoyl groups".

[0102] Among the groups represented by the formula (ω -12A), those groups in which both R^{a1} and R^{b1} are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-thiocarbamoyl groups", those groups in which both R^{a1} and R^{b1} are heterocyclic groups are referred to as "N,N-di(heteroring)-thiocarbamoyl groups", those groups in which R^{a1} is a hydrocarbon group and R^{b1} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-thiocarbamoyl groups", and those groups in which R^{a1} and R^{b1} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclic amino-thiocarbonyl groups".

[0103] Among the groups represented by the formula (ω -13A), those groups in which R^{a1} is a hydrocarbon group are referred to as "N-hydrocarbon-sulfamoyl groups", and those groups in which R^{a1} is a heterocyclic group are referred to as "N-heteroring-sulfamoyl groups".

[0104] Among the groups represented by the formula (ω -14A), those groups in which both R^{a1} and R^{b1} are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-sulfamoyl groups" whose example includes, for example, N,N-dimethylsulfamoyl group, those groups in which both R^{a1} and R^{b1} are heterocyclic groups are referred to as "N,N-di(heteroring)sulfamoyl groups", those groups in which R^{a1} is a hydrocarbon group and R^{b1} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-sulfamoyl groups", and those groups in which R^{a1} and R^{b1} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclic amino-sulfonyl groups" which include, for example 1-pyrrolylsulfonyl group.

[0105] Among the groups represented by the formula (ω -15A), those groups in which R^{a1} is a hydrocarbon group are referred to as "N-hydrocarbon-sulfinamoyl groups", and those groups in which R^{a1} is a heterocyclic group are referred to as "N-heteroring-sulfinamoyl groups".

[0106] Among the groups represented by the formula (ω -16A), those groups in which both R^{a1} and R^{b1} are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-sulfinamoyl groups", those groups in which both R^{a1} and R^{b1} are heterocyclic groups are referred to as "N,N-di(heteroring)-sulfinamoyl groups", those groups in which R^{a1} is a hydrocarbon group and R^{b1} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-sulfinamoyl groups", and those groups in which R^{a1} and R^{b1} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclic amino-sulfinyl groups".

[0107] Among the groups represented by the formula (ω -17A), those groups in which R^{a1} is a hydrocarbon group are referred to as "hydrocarbon-oxy-sulfonyl groups", and those groups in which R^{a1} is a heterocyclic group are referred to as "heteroring-oxy-sulfonyl groups".

[0108] Among the groups represented by the formula (ω -18A), those groups in which R^{a1} is a hydrocarbon group are referred to as "hydrocarbon-oxy-sulfinyl groups", and those groups in which R^{a1} is a heterocyclic group are referred to as "heteroring-oxy-sulfinyl groups".

[0109] Among the groups represented by the formula (ω -19A), those groups in which both R^{a1} and R^{b1} are hydrocarbon groups are referred to as "O,O'-di(hydrocarbon)-phosphono groups", those groups in which both R^{a1} and R^{b1} are heterocyclic groups are referred to as "O,O'-di(heteroring)-phosphono groups", and those groups in which R^{a1} is a hydrocarbon group and R^{b1} is a heterocyclic group are referred to as "O-hydrocarbon-O'-heteroring-phosphono groups".

[0110] Among the groups represented by the formula (ω -20A), those groups in which R^{a1} is a hydrocarbon group-

are referred to as "hydrocarbon-sulfonyl groups" whose examples include, for example, methanesulfonyl and benzenesulfonyl, and those groups in which R^{a1} is a heterocyclic group are referred to as "heteroring-sulfonyl groups".

[0111] Among the groups represented by the formula (ω -21A), those groups in which R^{a1} is a hydrocarbon group are referred to as "hydrocarbon-sulfinyl groups" whose examples include, for example, methylsulfinyl and benzenesulfinyl, and those groups in which R^{a1} is a heterocyclic group are referred to as "heteroring-sulfinyl groups".

[0112] Examples of the hydrocarbon in the groups represented by the aforementioned formulas (ω -1A) through (ω -21A) include the similar groups to the aforementioned hydrocarbon group. Examples of the hydrocarbon-carbonyl groups represented by the formula (ω -1A) include, for example, an alkyl-carbonyl group, an alkenyl-carbonyl group, an alkynyl-carbonyl group, a cycloalkyl-carbonyl group, a cycloalkenyl-carbonyl group, a cycloalkanedienyl-carbonyl group, a cycloalkyl-alkyl-carbonyl group which is an aliphatic hydrocarbon-carbonyl group, an aryl-carbonyl group, an aralkyl-carbonyl group, a bridged cyclic hydrocarbon-carbonyl group, a spirocyclic hydrocarbon-carbonyl group, and a terpene family hydrocarbon-carbonyl. In the following, groups represented by the formulas (ω -2A) through (ω -21A) are similar to those explained above.

[0113] Examples of the heteroring in the groups represented by the aforementioned formulas (ω -1A) through (ω -21A) include similar groups to the aforementioned heterocyclic group. Examples of the heteroring-carbonyl group represented by the formula (ω -1A) include, for example, a monocyclic heteroaryl-carbonyl group, a fused polycyclic heteroaryl-carbonyl group, a monocyclic nonaromatic heteroring-carbonyl group, and a fused polycyclic nonaromatic heteroring-carbonyl. In the following, groups represented by the formulas (ω -2A) through (ω -21A) are similar to those explained above.

[0114] Examples of the cyclic amino in the groups represented by the aforementioned formulas (ω -10A) through (ω -16A) include similar groups to the aforementioned cyclic amino group.



















[0115] The aforementioned acyl group, carbamoyl group, thiocarbamoyl group, sulfamoyl group, and sulfinamoyl group are generically referred to as "acyl groups" which may be substituted.




[0116] In the present specification, when a certain functional group is defined as "which may be substituted", the definition means that the functional group may sometimes have one or more substituents at chemically substitutable positions, unless otherwise specifically mentioned. Kind of substituents, number of substituents, and the position of substituents existing in the functional groups are not particularly limited, and when two or more substituents exist, they may be the same or different. Examples of the substituent existing in the functional group include, for example, halogen atoms, oxo group, thioxo group, nitro group, nitroso group, cyano group, isocyano group, cyanato group, thiocyanato group, isocyanato group, isothiocyanato group, hydroxy group, sulfanyl group, carboxy group, sulfanylcabonyl group, oxalo group, methooxalo group, thiocarboxy group, dithiocarboxy group, carbamoyl group, thiocarbamoyl group, sulfo group, sulfamoyl group, sulfinoyl group, sulfinamoyl group, sulfeno group, sulfenamoyl group, phosphono group, hydroxyphosphonyl group, hydrocarbon group, heterocyclic group, hydrocarbon-oxy group, heteroring-oxy group, hydrocarbon-sulfanyl group, heteroring-sulfanyl group, acyl group, amino group, hydrazino group, hydrazono group, diazenyl group, ureido group, thioureido group, guanidino group, carbamoimidoyl group (amidino group), azido group, imino group, hydroxyamino group, hydroxyimino group, aminooxy group, diazo group, semicarbazino group, semicarbazono group, allophanoyl group, hydantoyl group, phosphano group, phosphoroso group, phospho group, boryl group, silyl group, stannyl group, selanyl group, oxido group and the like.

[0117] When two or more substituents exist according to the abovementioned definition of "which may be substituted", said two or more substituents may combine to each other, together with atom(s) to which they bind, to form a ring. For these cyclic groups, as ring-constituting atoms (ring forming atoms), one to three kinds of one or more hetero atoms selected from oxygen atom, sulfur atom, nitrogen atom and the like may be included, and one or more substituents may exist on the ring. The ring may be monocyclic or fused polycyclic, and aromatic or nonaromatic.

[0118] The above substituents according to the abovementioned definition of "which may be substituted" may further be substituted with the aforementioned substituents at the chemically substitutable positions on the substituent. Kind of substituents, number of substituents, and positions of substituents are not particularly limited, and when the substituents are substituted with two or more substituents, they may be the same or different. Examples of the substituent include, for example, a halogenated alkyl-carbonyl group (trifluoroacetyl group as an example), a halogenated alkyl-sulfonyl group (trifluoromethanesulfonyl group as an example), an acyl-oxy group, an acyl-sulfanyl group, an N-hydrocarbon-amino group, an N,N-di(hydrocarbon)-amino group, an N-heteroring-amino group, an N-hydrocarbon-N-heteroring-amino group, an acyl-amino group, and a di(acyl)-amino group. Moreover, substitution on the aforementioned substituents may be repeated multiple orders.

[0119] Examples of the acyl-oxy group include the group in which hydrogen atom of hydroxy group is substituted with acyl group, and include, for example, formyloxy group, glyoxyloxy group, and thioformyloxy group, and the group represented by the following formulas:





wherein $\text{R}^{\text{a}2}$ and $\text{R}^{\text{b}2}$ may be the same or different and represent a hydrocarbon group or a heterocyclic group, or $\text{R}^{\text{a}2}$ and $\text{R}^{\text{b}2}$ combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group.

[0120] In the definition of aforementioned acyl-oxy group, among the groups represented by the formula ($\omega-1\text{B}$), those groups in which $\text{R}^{\text{a}2}$ is a hydrocarbon group are referred to as "hydrocarbon-carbonyl-oxy group" whose examples include, for example, acetoxy and benzyloxy group, and those groups in which $\text{R}^{\text{a}2}$ is a heterocyclic group are referred to as "heteroring-carbonyl-oxy group".

[0121] Among the groups represented by the formula ($\omega-2\text{B}$), those groups in which $\text{R}^{\text{a}2}$ is a hydrocarbon group are referred to as "hydrocarbon-oxy-carbonyl-oxy group", and those groups in which $\text{R}^{\text{a}2}$ is a heterocyclic group are referred to as "heteroring-oxy-carbonyl-oxy group".

[0122] Among the groups represented by the formula ($\omega-3\text{B}$), those groups in which $\text{R}^{\text{a}2}$ is a hydrocarbon group are referred to as "hydrocarbon-carbonyl-carbonyl-oxy group", and groups in which $\text{R}^{\text{a}2}$ is a heterocyclic group are referred to as "heteroring-carbonyl-carbonyl-oxy group".

[0123] Among the groups represented by the formula ($\omega-4\text{B}$), those groups in which $\text{R}^{\text{a}2}$ is a hydrocarbon group are referred to as "hydrocarbon-oxy-carbonyl-carbonyl-oxy group", and groups in which $\text{R}^{\text{a}2}$ is a heterocyclic group are referred to as "heteroring-oxy-carbonyl-carbonyl-oxy group".

[0124] Among the groups represented by the formula ($\omega-5\text{B}$), those groups in which $\text{R}^{\text{a}2}$ is a hydrocarbon group are referred to as "hydrocarbon-sulfanyl-carbonyl-oxy group", and groups where $\text{R}^{\text{a}2}$ is a heterocyclic group are referred to as "heteroring-sulfanyl-carbonyl-oxy group".

[0125] Among the groups represented by the formula ($\omega-6\text{B}$), those groups in which $\text{R}^{\text{a}2}$ is a hydrocarbon group are referred to as "hydrocarbon-thiocarbonyl-oxy group", and groups where $\text{R}^{\text{a}2}$ is a heterocyclic group are referred to as "heteroring-thiocarbonyl-oxy group".

[0126] Among the groups represented by the formula ($\omega-7\text{B}$), those groups in which $\text{R}^{\text{a}2}$ is a hydrocarbon group are referred to as "hydrocarbon-oxy-thiocarbonyl-oxy group", and groups in which $\text{R}^{\text{a}2}$ is a heterocyclic group are referred to as "heteroring-oxy-thiocarbonyl-oxy group".

[0127] Among the groups represented by the formula ($\omega-8\text{B}$), those groups in which $\text{R}^{\text{a}2}$ is a hydrocarbon group are referred to as "hydrocarbon-sulfanyl-thiocarbonyl-oxy group", and groups wherein $\text{R}^{\text{a}2}$ is a heterocyclic group are referred to as "heteroring-sulfanyl-thiocarbonyl-oxy group".

[0128] Among the groups represented by the formula ($\omega-9\text{B}$), those groups in which $\text{R}^{\text{a}2}$ is a hydrocarbon group are referred to as "N-hydrocarbon-carbamoyl-oxy group", and groups in which $\text{R}^{\text{a}2}$ is a heterocyclic group are referred to as "N-heteroring-carbamoyl-oxy group".

[0129] Among the groups represented by the formula ($\omega-10\text{B}$), those groups in which both $\text{R}^{\text{a}2}$ and $\text{R}^{\text{b}2}$ are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-carbamoyl-oxy group", those groups in which both $\text{R}^{\text{a}2}$ and $\text{R}^{\text{b}2}$ are heterocyclic groups are referred to as "N,N-di(heteroring)-carbamoyl-oxy group", those groups in which $\text{R}^{\text{a}2}$ is a hydrocarbon group and $\text{R}^{\text{b}2}$ is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-carbamoyl-oxy group", and those groups in which $\text{R}^{\text{a}2}$ and $\text{R}^{\text{b}2}$ combine each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclicamino-carbonyl-oxy group".

[0130] Among the groups represented by the formula ($\omega-11\text{B}$), those groups in which $\text{R}^{\text{a}2}$ is a hydrocarbon group are referred to as "N-hydrocarbon-thiocarbamoyl-oxy group", and those groups in which $\text{R}^{\text{a}2}$ is a heterocyclic group are referred to as "N-heteroring-thiocarbamoyl-oxy group".

[0131] Among the groups represented by the formula ($\omega-12\text{B}$), those groups in which both $\text{R}^{\text{a}2}$ and $\text{R}^{\text{b}2}$ are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-thiocarbamoyl-oxy group", those groups in which both $\text{R}^{\text{a}2}$ and $\text{R}^{\text{b}2}$ are heterocyclic groups are referred to as "N,N-di(heteroring)-thiocarbamoyl-oxy group", those groups wherein $\text{R}^{\text{a}2}$ is a hydrocarbon group and $\text{R}^{\text{b}2}$ is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-thiocarbamoyl-oxy group", and those groups in which $\text{R}^{\text{a}2}$ and $\text{R}^{\text{b}2}$ combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclicamino-thiocarbonyl-oxy group".

[0132] Among the groups represented by the formula (ω -13B), those groups in which R^{a2} is a hydrocarbon group are referred to as "N-hydrocarbon-sulfamoyl-oxy groups", and those groups in which R^{a2} is a heterocyclic group are referred to as "N-heteroring-sulfamoyl-oxy groups".

[0133] Among the groups represented by the formula (ω -14B), those groups in which both R^{a2} and R^{b2} are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-sulfamoyl-oxy groups", those groups in which both R^{a2} and R^{b2} are heterocyclic groups are referred to as "N,N-di(heteroring)-sulfamoyl-oxy groups", those groups in which R^{a2} is a hydrocarbon group and R^{b2} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-sulfamoyl-oxy groups", and those groups in which R^{a2} and R^{b2} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclic amino-sulfonyl-oxy groups".

[0134] Among the groups represented by the formula (ω -15B), those groups in which R^{a2} is a hydrocarbon group are referred to as "N-hydrocarbon-sulfinamoyl-oxy groups", and those groups where R^{a2} is a heterocyclic group are referred to as "N-heteroring-sulfinamoyl-oxy groups".

[0135] Among the groups represented by the formula (ω -16B), those groups in which both R^{a2} and R^{b2} are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-sulfinamoyl-oxy groups", those groups in which both R^{a2} and R^{b2} are heterocyclic groups are referred to as "N,N-di(heteroring)-sulfinamoyl-oxy groups", those groups in which R^{a2} is a hydrocarbon group and R^{b2} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-sulfinamoyl-oxy groups", and those groups in which R^{a2} and R^{b2} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclic amino-sulfinyl-oxy group".

[0136] Among the groups represented by the formula (ω -17B), those groups in which R^{a2} is a hydrocarbon group are referred to as "hydrocarbon-oxy-sulfonyl-oxy group", and those groups in which R^{a2} is a heterocyclic group are referred to as "heteroring-oxy-sulfonyl-oxy group".

[0137] Among the groups represented by the formula (ω -18B), those groups in which R^{a2} is a hydrocarbon group are referred to as "hydrocarbon-oxy-sulfinyl-oxy groups", those groups in which R^{a2} is a heterocyclic group are referred to as "heteroring-oxy-sulfinyl-oxy groups".

[0138] Among the groups represented by the formula (ω -19B), those groups in which both R^{a2} and R^{b2} are hydrocarbon groups are referred to as "O,O'-di(hydrocarbon)-phosphono-oxy group", groups in which both R^{a2} and R^{b2} are heterocyclic groups are referred to as "O,O'-di(heteroring)-phosphono-oxy group", and those groups in which R^{a2} is a hydrocarbon group and R^{b2} is a heterocyclic group are referred to as "O-hydrocarbon substituted-O'-heteroring substituted phosphono-oxy group".

[0139] Among the groups represented by the formula (ω -20B), those groups in which R^{a2} is a hydrocarbon group are referred to as "hydrocarbon-sulfonyl-oxy group", and those groups in which R^{a2} is a heterocyclic group referred to as "heteroring-sulfonyl-oxy group".

[0140] Among the groups represented by the formula (ω -21B), those groups in which R^{a2} is a hydrocarbon group are referred to as "hydrocarbon-sulfinyl-oxy group", and those groups in which R^{a2} is a heterocyclic group are referred to as "heteroring-sulfinyl-oxy group".



















[0141] Examples of the hydrocarbon in the groups represented by the aforementioned formulas (ω -1B) through (ω -21B) include the similar groups to the aforementioned hydrocarbon group. Examples of the hydrocarbon-carbonyl-oxy groups represented by the formula (ω -1B) include, for example, an alkyl-carbonyl-oxy group, an alkenyl-carbonyl-oxy group, an alkynyl-carbonyl-oxy group, a cycloalkyl-carbonyl-oxy group, a cycloalkenyl-carbonyl-oxy group, a cycloalkenedienyl-carbonyl-oxy group, and a cycloalkyl-alkyl-carbonyl-oxy group, which are aliphatic hydrocarbon-carbonyl-oxy groups, an aryl-carbonyl-oxy group, an aralkyl-carbonyl-oxy group, a bridged cyclic hydrocarbon-carbonyl-oxy group, a spirocyclic hydrocarbon-carbonyl-oxy group, and a terpene family hydrocarbon-carbonyl-oxy group. In the following, groups represented by the formulas (ω -2B) through (ω -21B) are similar to those explained above.




[0142] Examples of the heteroring in the groups represented by the aforementioned formulas (ω -1B) through (ω -21B) include similar groups to the aforementioned heterocyclic group. Examples of the heteroring-carbonyl group represented by the formula (ω -1B) include, for example, a monocyclic heteroaryl-carbonyl group, a fused polycyclic heteroaryl-carbonyl group, a monocyclic nonaromatic heteroring-carbonyl group, and a fused polycyclic nonaromatic heteroring-carbonyl group. In the following, groups represented by the formulas (ω -2B) through (ω -21B) are similar to those groups mentioned above.

[0143] Examples of the cyclic amino in the groups represented by the aforementioned formulas (ω -10B) through (ω -16B) include similar groups to the aforementioned cyclic amino group.

[0144] The aforementioned acyl-oxy group, hydrocarbon-oxy group, and heterocyclic-oxy group are generically referred to as "substituted oxy group". Moreover, these substituted oxy group and hydroxy group are generically referred to as "hydroxy group" which may be substituted.

[0145] Examples of the acyl-sulfanyl include the group in which hydrogen atom of sulfanyl group is substituted with acyl group, and include, for example, formylsulfanyl group, glyoxylsulfanyl group, and thioformylsulfanyl group, and groups represented by the following formulas:





wherein R^{a3} and R^{b3} may be the same or different and represent a hydrocarbon group which may be substituted or a heterocyclic group which may be substituted, or R^{a3} and R^{b3} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group which may be substituted.

[0146] In the definition of aforementioned acyl sulfanyl group, among the groups represented by the formula ($\omega-1\text{C}$), those groups in which R^{a3} is a hydrocarbon group are referred to as "hydrocarbon-carbonyl-sulfanyl group", and those groups in which R^{a3} is a heterocyclic group are referred to as "heteroring-carbonyl-sulfanyl group".

[0147] Among the groups represented by the formula ($\omega-2\text{C}$), those groups in which R^{a3} is a hydrocarbon group are referred to as "hydrocarbon-oxy-carbonyl-sulfanyl group", and those groups in which R^{a3} is a heterocyclic group are referred to as "heteroring-oxy-carbonyl-sulfanyl group".

[0148] Among the groups represented by the formula ($\omega-3\text{C}$), those groups in which R^{a3} is a hydrocarbon group are referred to as "hydrocarbon-carbonyl-carbonyl-sulfanyl group", and those groups in which R^{a3} is a heterocyclic group are referred to as "heteroring-carbonyl-carbonyl-sulfanyl group".

[0149] Among the groups represented by the formula ($\omega-4\text{C}$), those groups in which R^{a3} is a hydrocarbon group are referred to as "hydrocarbon-oxy-carbonyl-carbonyl-sulfanyl group", and those groups in which R^{a3} is a heterocyclic group are referred to as "heteroring-oxy-carbonyl-carbonyl-sulfanyl group".

[0150] Among the groups represented by the formula ($\omega-5\text{C}$), those groups in which R^{a3} is a hydrocarbon group are referred to as "hydrocarbon-sulfanyl-carbonyl-sulfanyl group", and those groups in which R^{a3} is a heterocyclic group are referred to as "heteroring-sulfanyl-carbonyl-sulfanyl group".

[0151] Among the groups represented by the formula ($\omega-6\text{C}$), those groups in which R^{a3} is a hydrocarbon group are referred to as "hydrocarbon-thiocarbonyl-sulfanyl group", and those groups in which R^{a3} is a heterocyclic group are referred to as "heteroring-thiocarbonyl-sulfanyl group".

[0152] Among the groups represented by the formula ($\omega-7\text{C}$), those groups in which R^{a3} is a hydrocarbon group are referred to as "hydrocarbon-oxy-thiocarbonyl-sulfanyl group", and those groups in which R^{a3} is a heterocyclic group are referred to as "heteroring-oxy-thiocarbonyl-sulfanyl group".

[0153] Among the groups represented by the formula ($\omega-8\text{C}$), those groups in which R^{a3} is a hydrocarbon group are referred to as "hydrocarbon-sulfanyl-thiocarbonylsulfanyl group", and those groups in which R^{a3} is a heterocyclic group are referred to as "heteroring-sulfanyl-thiocarbonyl-sulfanyl group".

[0154] Among the groups represented by the formula ($\omega-9\text{C}$), those groups in which R^{a3} is a hydrocarbon group are referred to as "N-hydrocarbon-carbamoyl-sulfanyl group", and those groups in which R^{a3} is a heterocyclic group are referred to as "N-heteroring-carbamoyl-sulfanyl group".

[0155] Among the groups represented by the formula ($\omega-10\text{C}$), those groups in which both R^{a3} and R^{b3} are a hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-carbamoyl-sulfanyl group", those groups in which both R^{a3} and R^{b3} are heterocyclic groups are referred to as "N,N-di(heteroring)-carbamoyl-sulfanyl group", groups in which R^{a3} is a hydrocarbon group and R^{b3} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-carbamoyl-sulfanyl group", and those groups in which R^{a3} and R^{b3} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclicamino-carbonyl-sulfamoyl group".

[0156] Among the groups represented by the formula ($\omega-11\text{C}$), those groups in which R^{a3} is a hydrocarbon group are referred to as "N-hydrocarbon-thiocarbamoyl-sulfanyl group", and those groups in which R^{a3} is a heterocyclic group are referred to as "N-heteroring-thiocarbamoyl-sulfanyl group".

[0157] Among the groups represented by the formula ($\omega-12\text{C}$), those groups in which both R^{a3} and R^{b3} are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-thiocarbamoyl-sulfanyl group", those groups in which R^{a3} and R^{b3} are heterocyclic groups are referred to as "N,N-di(heteroring)-thiocarbamoyl-sulfanyl group", those groups in which R^{a3} is a hydrocarbon group and R^{b3} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-thiocarbamoyl-sulfanyl group", and those groups in which R^{a3} and R^{b3} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclicamino-thiocarbonyl-sulfamoyl group".

[0158] Among the groups represented by the formula (ω -13C), those groups in which R^{a3} is a hydrocarbon group are referred to as "N-hydrocarbon-sulfamoyl-sulfanyl group", and those groups in which R^{a3} is a heteroring group are referred to as "N-heterocyclic-sulfamoyl-sulfanyl group".

[0159] Among the groups represented by the formula (ω -14C), those groups in which both R^{a3} and R^{b3} are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-sulfamoyl-sulfanyl group", those groups in which both R^{a3} and R^{b3} are heterocyclic groups are referred to as "N,N-di(heteroring)-sulfamoyl-sulfanyl group", those groups in which R^{a3} is a hydrocarbon group and R^{b3} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-sulfamoyl-sulfanyl group", and those groups in which R^{a3} and R^{b3} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclicamino-sulfonyl-sulfanyl group".

[0160] Among the groups represented by the formula (ω -15C), those groups in which R^{a3} is a hydrocarbon group are referred to as "N-hydrocarbon-sulfinamoyl-sulfanyl group", and those groups in which R^{a3} is a heteroring group are referred to as "N-heterocyclic-sulfinamoyl-sulfanyl group".

[0161] Among the groups represented by the formula (ω -16C), those groups in which both R^{a3} and R^{b3} are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-sulfinamoyl-sulfanyl group", those groups in which both R^{a3} and R^{b3} are heterocyclic groups are referred to as "N,N-di(heteroring)-sulfinamoyl-sulfanyl group", those groups in which R^{a3} is a hydrocarbon group and R^{b3} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-sulfinamoyl-sulfanyl group", and those groups in which R^{a3} and R^{b3} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclicamino-sulfanyl-sulfanyl group".

[0162] Among the groups represented by the formula (ω -17C), those groups in which R^{a3} is a hydrocarbon group are referred to as "hydrocarbon-oxy-sulfonyl-sulfanyl group", and those groups in which R^{a3} is a heteroring group are referred to as "heterocyclic-oxy-sulfonyl-sulfanyl group".

[0163] Among the groups represented by the formula (ω -18C), those groups in which R^{a3} is a hydrocarbon group are referred to as "hydrocarbon-oxy-sulfinyl-sulfanyl group", and those groups in which R^{a3} is a heteroring group are referred to as "heterocyclic-oxy-sulfinyl-sulfanyl group".

[0164] Among the groups represented by the formula (ω -19C), those groups in which both R^{a3} and R^{b3} are hydrocarbon groups are referred to as "O,O'-di(hydrocarbon)-phosphono-sulfanyl group", those groups in which both R^{a3} and R^{b3} are heterocyclic groups are referred to as "O,O'-di(heteroring)-phosphono-sulfanyl group", and those groups in which R^{a3} is a hydrocarbon group and R^{b3} is a heteroring group are referred to as "O-hydrocarbon-O'-heterocyclic-phosphono-sulfanyl group". Among the groups represented by the formula (ω -20C), those groups in which R^{a3} is a hydrocarbon group are referred to as "hydrocarbon-sulfonyl-sulfanyl group", and those groups in which R^{a3} is a heterocyclic group are referred to as "heteroring-sulfonyl-sulfanyl group".

[0165] Among the groups represented by the formula (ω -21C), those groups in which R^{a3} is a hydrocarbon group are referred to as "hydrocarbon-sulfinyl-sulfanyl group", and those groups in which R^{a3} is a heteroring group are referred to as "heterocyclic-sulfinyl-sulfanyl group".

[0166] Examples of the hydrocarbon in the groups represented by the aforementioned formulas (ω -1C) through (ω -21C) include similar groups to the aforementioned hydrocarbon group. Examples of the hydrocarbon-carbonyl-sulfanyl groups represented by the formula (ω -1C) include, for example, an alkyl-carbonyl-sulfanyl group, an alkenyl-carbonyl-sulfanyl group, an alkynyl-carbonyl-sulfanyl group, a cycloalkyl-carbonyl-sulfanyl group, a cycloalkenyl-carbonyl-sulfanyl group, a cycloalkenedienyl-carbonyl-sulfanyl group, a cycloalkyl-alkyl-carbonyl-sulfanyl group which is an aliphatic hydrocarbon-carbonyl-sulfanyl group, an aryl-carbonyl-sulfanyl group, an aralkyl-carbonyl-sulfanyl group, a bridged cyclic hydrocarbon-carbonyl-sulfanyl group, a spiro cyclic hydrocarbon-carbonyl-sulfanyl group, and a terpene family hydrocarbon-carbonyl-sulfanyl group. In the following, groups represented by the formulas (ω -2C) through (ω -21C) are similar to those mentioned above.

[0167] Examples of the heteroring in the groups represented by the aforementioned formulas (ω -1C) through (ω -21C) include similar groups to the aforementioned heterocyclic group. Examples of the heteroring-carbonyl-sulfanyl group represented by the formula (ω -1C) include, for example, a monocyclic heteroaryl-carbonyl-sulfanyl group, a fused polycyclic heteroaryl-carbonyl-sulfanyl group, a monocyclic nonaromatic heteroring-carbonyl-sulfanyl group, and a fused polycyclic non-aromatic heteroring-carbonyl-sulfanyl group. In the following, groups represented by the formula (ω -2C) through (ω -21C) are similar to those groups mentioned above.

[0168] Examples of the cyclic amino in the groups represented by the aforementioned formulas (ω -10C) through (ω -16C) include similar groups to the aforementioned cyclic amino group.

[0169] The aforementioned acyl-sulfanyl group, hydrocarbon-sulfanyl group, and heterocyclic-sulfanyl group are generically referred to as "substituted sulfanyl group". These substituted sulfanyl group and sulfanyl group are generically referred to as "sulfanyl groups" which may be substituted.

[0170] Examples of the N-hydrocarbon-amino group include the group in which one hydrogen atom of amino group is substituted with a hydrocarbon group, and include, for example, an N-alkyl-amino group, an N-alkenyl-amino group, an N-alkynyl-amino group, an N-cycloalkyl-amino group, an N-cycloalkyl-alkyl-amino group, an N-aryl-amino group, and an N-aralkyl-amino group.

[0171] Examples of the N-alkyl-amino group include, for example, methylamino, ethylamino, n-propylamino, isopropylamino, n-butylamino, isobutylamino, sec-butylamino, tert-butylamino, n-pentylamino, isopentylamino, (2-methylbutyl)amino, (1-methylbutyl)amino, neopentylamino, (1,2-dimethylpropyl)amino, (1-ethylpropyl)amino, n-hexylamino, (4-methylpentyl)amino, (3-methylpentyl)amino, (2-methylpentyl)amino, (1-methylpentyl)amino, (3,3-dimethylbutyl)amino, (2,2-dimethylbutyl)amino, (1,1-dimethylbutyl)amino, (1,2-dimethylbutyl)amino, (1,3-dimethylbutyl)amino, (2,3-dimethylbutyl)amino, (2-ethylbutyl)amino, (1-ethylbutyl)amino, (1-ethyl-1-methylpropyl)amino, n-heptylamino, n-octylamino, n-nonylamino, n-decylamino, n-undecylamino, n-dodecylamino, n-tridecylamino, n-tetradecylamino, and n-pentadecylamino group, which are C₁ to C₁₅ straight chain or branched chain N-alkyl amino groups.

[0172] Examples of the N-alkenyl-amino group include, for example, vinyl amino, (prop-1-en-1-yl)amino, allylamino, isopropenylamino, (but-1-en-1-yl)amino, (but-2-en-1-yl)amino, (but-3-en-1-yl)amino, (2-methylprop-2-en-1-yl)amino, (1-methylprop-2-en-1-yl)amino, (pent-1-en-1-yl)amino, (pent-2-en-1-yl)amino, (pent-3-en-1-yl)amino, (pent-4-en-1-yl)amino, (3-methylbut-2-en-1-yl)amino, (3-methylbut-3-en-1-yl)amino, (hex-1-en-1-yl)amino, (hex-2-en-1-yl)amino, (hex-3-en-1-yl)amino, (hex-4-en-1-yl)amino, (hex-5-en-1-yl)amino, (4-methylpent-3-en-1-yl)amino, (4-methylpent-3-en-1-yl)amino, (hept-1-en-1-yl)amino, (hept-6-en-1-yl)amino, (oct-1-en-1-yl)amino, (oct-7-en-1-yl)amino, (non-1-en-1-yl)amino, (non-8-en-1-yl)amino, (dec-1-en-1-yl)amino, (dec-9-en-1-yl)amino, (undec-1-en-1-yl)amino, (undec-10-en-1-yl)amino, (dodec-1-en-1-yl)amino, (dodec-11-en-1-yl)amino, (tridec-1-en-1-yl)amino, (tridec-12-en-1-yl)amino, (tetradec-1-en-1-yl)amino, (tetradec-13-en-1-yl)amino, (pentadec-1-en-1-yl)amino, and (pentadec-14-en-1-yl)amino group, which are C₂ to C₁₅ straight chain or branched chain N-alkenyl amino groups.

[0173] Examples of the N-alkynyl-amino group include, for example, ethynylamino, (prop-1-yn-1-yl)amino, (prop-2-yn-1-yl)amino, (but-1-yn-1-yl)amino, (but-3-yn-1-yl)amino, (1-methylprop-2-yn-1-yl)amino, (pent-1-yn-1-yl)amino, (pent-4-yn-1-yl)amino, (hex-1-yn-1-yl)amino, (hex-5-yn-1-yl)amino, (hept-1-yn-1-yl)amino, (hept-6-yn-1-yl)amino, (oct-1-yn-1-yl)amino, (oct-7-yn-1-yl)amino, (non-1-yn-1-yl)amino, (non-8-yn-1-yl)amino, (dec-1-yn-1-yl)amino, (dec-9-yn-1-yl)amino, (undec-1-yn-1-yl)amino, (undec-10-yn-1-yl)amino, (dodec-1-yn-1-yl)amino, (dodec-11-yn-1-yl)amino, (tridec-1-yn-1-yl)amino, (tridec-12-yn-1-yl)amino, (tetradec-1-yn-1-yl)amino, (tetradec-13-yn-1-yl)amino, (pentadec-1-yn-1-yl)amino, and (pentadec-14-yn-1-yl)amino group, which are C₂ to C₁₅ straight chain or branched chain N-alkynyl-amino groups.

[0174] Examples of the N-cycloalkyl-amino group include, for example, cyclopropylamino, cyclobutylamino, cyclopentylamino, cyclohexylamino, cycloheptylamino, and cyclooctylamino group, which are C₃ to C₈ N-cycloalkyl-amino groups.

[0175] Examples of the N-cycloalkyl-alkyl-amino group include, for example, (cyclopropylmethyl)amino, (1-cyclopropylethyl)amino, (2-cyclopropylethyl)amino, (3-cyclopropylpropyl)amino, (4-cyclopropylbutyl)amino, (5-cyclopropylpentyl)amino, (6-cyclopropylhexyl)amino, (cyclobutylmethyl)amino, (cyclopentylmethyl)amino, (cyclobutylmethyl)amino, (cyclopentylmethyl)amino, (cyclohexylmethyl)amino, (2-cyclohexylethyl)amino, (3-cyclohexylpropyl)amino, (4-cyclohexylbutyl)amino, (cycloheptylmethyl)amino, (cyclooctylmethyl)amino, and (6-cyclooctylhexyl)amino, which are C₄ to C₁₄ N-cycloalkyl-alkyl-amino groups.

[0176] Examples of the N-aryl-amino group include, for example, phenylamino, 1-naphthylamino, 2-naphthylamino, anthrylamino, phenanthrylamino, and acenaphthylenylamino, which are C₆ to C₁₄ N-mono-aryl-amino groups.

[0177] Examples of the N-aralkyl-amino group include, for example, benzylamino, (1-naphthylmethyl)amino, (2-naphthylmethyl)amino, (anthracenylmethyl)amino, (phenanthrenylmethyl)amino, (acenaphthylenylmethyl)amino, (diphenylmethyl)amino, (1-phenethyl)amino, (2-phenethyl)amino, (1-(1-naphthyl)ethyl)amino, (1-(2-naphthyl)ethyl)amino, (2-(1-naphthyl)ethyl)amino, (2-(2-naphthyl)ethyl)amino, (3-phenylpropyl)amino, (3-(1-naphthyl)propyl)amino, (3-(2-naphthyl)propyl)amino, (4-phenylbutyl)amino, (4-(1-naphthyl)butyl)amino, (4-(2-naphthyl)butyl)amino, (5-phenylpentyl)amino, (5-(1-naphthyl)pentyl)amino, (5-(2-naphthyl)pentyl)amino, (6-phenylhexyl)amino, (6-(1-naphthyl)hexyl)amino, and (6-(2-naphthyl)hexyl)amino, which are C₇ to C₁₆ N-aralkyl-amino groups.



















[0178] Examples of the N,N-di(hydrocarbon)-amino group include the group in which 2 hydrogen atoms of amino group are substituted with hydrocarbon group, and include, for example, N,N-dimethylamino, N,N-diethylamino, N-ethyl-N-methylamino, N,N-di-n-propylamino, N,N-diisopropylamino, N-allyl-N-methylamino, N-(prop-2-yn-1-yl)-N-methylamino, N,N-dicyclohexylamino, N-cyclohexyl-N-methylamino, N-cyclohexylmethylamino-N-methylamino, N,N-diphenylamino, N-methyl-N-phenylamino, N,N-dibenzylamino, and N-benzyl-N-methylamino group.




[0179] Examples of the N-heteroring-amino group include the group in which one hydrogen atom of amino group is substituted with heterocyclic group, and include, for example, (3-pyrroliziny)amino, (4-piperidiny)amino, (2-tetrahydropyrany)amino, (3-indoliny)amino, (4-chromany)amino, (3-thienyl)amino, (3-pyridyl)amino, (3-quinolyl)amino, and (5-indolyl)amino.

[0180] Examples of the N-hydrocarbon-N-heteroring-amino group include the group in which 2 hydrogen atoms of amino group are substituted with hydrocarbon group and heterocyclic group respectively, and include, for example, N-methyl-N-(4-piperidiny)amino, N-(4-chromanyl)-N-methylamino, N-methyl-N-(3-thienyl)amino, N-methyl-N-(3-pyridyl)amino, N-methyl-N-(3-quinolyl)amino and the like.

[0181] Examples of the acyl-amino group include the group in which one hydrogen atom of the amino group is sub-

stituted with an acyl group, and include, for example, formylamino group, glyoxyloylamino group, and thioformylamino group, and groups represented by the following formulas :





wherein R^{a4} and R^{b4} may be the same or different and represent a hydrocarbon group which may be substituted or a heterocyclic group which may be substituted, or R^{a4} and R^{b4} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group which may be substituted.

[0182] In the definition of the aforementioned acyl-amino group, among the groups represented by the formula ($\omega-1\text{D}$), those groups in which R^{a4} is a hydrocarbon group are referred to as "hydrocarbon-carbonyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "heteroring-carbonyl-amino group".

[0183] Among the groups represented by the formula ($\omega-2\text{D}$), those groups in which R^{a4} is a hydrocarbon group are referred to as "hydrocarbon-oxy-carbonyl-amino group", and those groups in which R^{a4} is a heteroring group are referred to as "heterocyclic-oxy-carbonyl-amino group".

[0184] Among the groups represented by the formula ($\omega-3\text{D}$), those groups in which R^{a4} is a hydrocarbon group are referred to as "hydrocarbon-carbonyl-carbonyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "heteroring-carbonyl-carbonyl-amino group".

[0185] Among the groups represented by the formula ($\omega-4\text{D}$), those groups in which R^{a4} is a hydrocarbon group are referred to as "hydrocarbon-oxy-carbonyl-carbonyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "heteroring-oxy-carbonyl-carbonyl-amino group".

[0186] Among the groups represented by the formula ($\omega-5\text{D}$), those groups in which R^{a4} is a hydrocarbon group are referred to as "hydrocarbon-sulfanyl-carbonyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "heteroring-sulfanyl-carbonyl-amino group".

[0187] Among the groups represented by the formula ($\omega-6\text{D}$), those groups in which R^{a4} is a hydrocarbon group are referred to as "hydrocarbon-thiocarbonyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "heteroring-thiocarbonyl-amino group".

[0188] Among the groups represented by the formula ($\omega-7\text{D}$), those groups in which R^{a4} is a hydrocarbon group are referred to as "hydrocarbon-oxy-thiocarbonyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "heteroring-oxy-thiocarbonyl-amino group".

[0189] Among the groups represented by the formula ($\omega-8\text{D}$), those groups in which R^{a4} is a hydrocarbon group are referred to as "hydrocarbon-sulfanyl-thiocarbonyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "heteroring-sulfanyl-thiocarbonyl-amino group".

[0190] Among the groups represented by the formula ($\omega-9\text{D}$), those groups in which R^{a4} is a hydrocarbon group are referred to as "N-hydrocarbon-carbamoyl group", and those groups in which R^{a4} is a heterocyclic group are referred to as "N-heteroring-carbamoyl-amino group".

[0191] Among the groups represented by the formula ($\omega-10\text{D}$), those groups in which both R^{a4} and R^{b4} are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-carbamoyl-amino group", those groups in which both R^{a4} and R^{b4} are heterocyclic groups are referred to as "N,N-di(heteroring)-carbamoyl-amino group", those groups in which R^{a4} is a hydrocarbon group and R^{b4} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-carbamoyl-amino group", and those groups in which R^{a4} and R^{b4} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclicamino-carbonyl-amino group".

[0192] Among the groups represented by the formula ($\omega-11\text{D}$), those groups in which R^{a4} is a hydrocarbon group are referred to as "N-hydrocarbon-thiocarbamoyl-amino group", and those groups in which R^{a4} is a heteroring group are referred to as "N-heterocyclic-thiocarbamoyl-amino group".

[0193] Among the groups represented by the formula ($\omega-12\text{D}$), those groups in which both R^{a4} and R^{b4} are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-thiocarbamoyl-amino group", those groups in which both R^{a4} and R^{b4} are heterocyclic groups are referred to as "N,N-di(heteroring)-thiocarbamoyl-amino group", those groups in which R^{a4} is a hydrocarbon group and R^{b4} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-thiocarbamoyl-amino group", and those groups in which R^{a4} and R^{b4} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclicamino-thiocarbonyl-amino group".

[0194] Among the groups represented by the formula (ω -13D), those groups in which R^{a4} is a hydrocarbon group are referred to as "N-hydrocarbon-sulfamoyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "N-heteroring-sulfamoyl-amino group".

[0195] Among the groups represented by the formula (ω -14D), those groups in which both R^{a4} and R^{b4} are hydrocarbon groups are referred to as "di(hydrocarbon)-sulfamoyl-amino group", those groups in which both R^{a4} and R^{b4} are heterocyclic groups are referred to as "N,N-di(heteroring)-sulfamoyl-amino group", those groups in which R^{a4} is a hydrocarbon group and R^{b4} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-sulfamoyl-amino group", and those groups in which R^{a4} and R^{b4} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclicamino-sulfonyl-amino group".

[0196] Among the groups represented by the formula (ω -15D), those groups in which R^{a4} is a hydrocarbon group are referred to as "N-hydrocarbon-sulfinamoyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "N-heteroring-sulfinamoyl-amino group".

[0197] Among the groups represented by the formula (ω -16D), those groups in which both R^{a4} and R^{b4} are hydrocarbon groups are referred to as "N,N-di(hydrocarbon)-sulfinamoyl-amino group", those groups in which both R^{a4} and R^{b4} are heterocyclic groups are referred to as "N,N-di(heteroring)-sulfinamoyl-amino group", groups in which R^{a4} is a hydrocarbon group and R^{b4} is a heterocyclic group are referred to as "N-hydrocarbon-N-heteroring-sulfinamoyl-amino group", and those groups in which R^{a4} and R^{b4} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "cyclicamino-sulfinyl-amino group".

[0198] Among the groups represented by the formula (ω -17D), those groups in which R^{a4} is a hydrocarbon group are referred to as "hydrocarbon-oxy-sulfonyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "heteroring-oxy-sulfonyl-amino group".

[0199] Among the groups represented by the formula (ω -18D), those groups in which R^{a4} is a hydrocarbon group are referred to as "hydrocarbon-oxy-sulfinyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "heteroring-oxy-sulfinyl-amino group".

[0200] Among the groups represented by the formula (ω -19D), those groups in which both R^{a4} and R^{b4} are hydrocarbon groups are referred to as "O,O'-di(hydrocarbon)-phosphono-amino group", those groups in which both R^{a4} and R^{b4} are heterocyclic groups are referred to as "O,O'-di(heteroring)-phosphono-amino group", and those groups in which R^{a4} is a hydrocarbon group and R^{b4} is a heterocyclic group are referred to as "O-hydrocarbon-O'-heteroring-phosphono-amino group".

[0201] Among the groups represented by the formula (ω -20D), those groups in which R^{a4} is a hydrocarbon group are referred to as "hydrocarbon-sulfonyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "heteroring-sulfonyl-amino group".


[0202] Among the groups represented by the formula (ω -21D), those groups in which R^{a4} is a hydrocarbon group are referred to as "hydrocarbon-sulfinyl-amino group", and those groups in which R^{a4} is a heterocyclic group are referred to as "heteroring-sulfinyl-amino group".

[0203] Examples of the hydrocarbon in the groups represented by the aforementioned formulas (ω -1D) through (ω -21D), similar groups to the aforementioned hydrocarbon group. Examples of the hydrocarbon-carbonyl-amino groups represented by the formula (ω -1D) include, for example, an alkyl-carbonyl-amino group, an alkenyl-carbonyl-amino group, an alkynyl-carbonyl-amino group, a cycloalkyl-carbonyl-amino group, a cycloalkenyl-carbonyl-amino group, a cycloalkanediaryl-carbonyl-amino group, a cycloalkyl-alkyl-carbonyl-amino group which is an aliphatic hydrocarbon-carbonyl-amino groups, an aryl-carbonyl-amino group, an aralkyl-carbonyl-amino group, a bridged cyclic hydrocarbon-carbonyl-amino group, a spiro cyclic hydrocarbon-carbonyl-amino group, and a terpene family hydrocarbon-carbonyl-amino group. In the following, groups represented by the formulas (ω -2D) through (ω -21D) are similar to those explained above.

[0204] Examples of the heteroring in the groups represented by the aforementioned formulas (ω -1D) through (ω -21D) include similar groups to the aforementioned heterocyclic group. Examples of the heteroring-carbonyl-amino group represented by the formula (ω -1D) include, for example, a monocyclic heteroaryl-carbonyl-amino group, a fused polycyclic heteroaryl-carbonyl-amino group, a monocyclic nonaromatic heterocyclic-carbonyl-amino group, and a fused polycyclic nonaromatic heterocyclic-carbonyl-amino group. In the following, groups represented by the formulas (ω -2D) through (ω -21D) are similar to those groups mentioned above.

[0205] Examples of the cyclic amino in the groups represented by the aforementioned formulas (ω -10D) through (ω -16D) include similar groups to the aforementioned cyclic amino group.

[0206] The aforementioned di(acyl)-amino group include the group in which 2 hydrogen atoms of amino group are substituted with acyl groups in the definitions of the aforementioned substituents according to "which may be substituted". Examples include, for example, di(formyl)-amino group, di(glyoxyloyl)-amino group, and di(thioformyl)-amino group, and groups represented by the following formulas :



wherein R^{a5} and R^{b5} may be the same or different and represent hydrogen atom, a hydrocarbon group which may be substituted or a heterocyclic group which may be substituted, or R^{a5} and R^{b5} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group which may be substituted.

[0207] In the definition of aforementioned di(acyl)-amino group, among the groups represented by the formula ($\omega - 1 \text{ E}$), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(hydrocarbon-carbonyl)-amino group", and those groups in which R^{a5} is a heteroring group are referred to as "bis(heterocyclic-carbonyl)-amino group".

[0208] Among the groups represented by the formula ($\omega - 2 \text{ E}$), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(hydrocarbon-oxy-carbonyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(heteroring-oxy-carbonyl)-amino group".

[0209] Among the groups represented by the formula ($\omega - 3 \text{ E}$), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(hydrocarbon-carbonyl-carbonyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(heteroring-carbonyl-carbonyl)-amino group". Among the groups represented by the formula ($\omega - 4 \text{ E}$), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(hydrocarbon-oxy-carbonyl-carbonyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(heteroring-oxy-carbonyl-carbonyl)-amino group".

[0210] Among the groups represented by the formula (ω -5E), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(hydrocarbon-sulfanyl-carbonyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(heteroring-sulfanyl-carbonyl)-amino group".

[0211] Among the groups represented by the formula (ω -6E), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(hydrocarbon-thiocarbonyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(heteroring-thiocarbonyl)-amino group".

[0212] Among the groups represented by the formula (ω -7E), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(hydrocarbon-oxy-thiocarbonyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(ringoxy-thiocarbonyl)-amino group".

[0213] Among the groups represented by the formula (ω -8E), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(hydrocarbon-sulfanyl-thiocarbonyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(heteroring-sulfanyl-thiocarbonyl)-amino group".

[0214] Among the groups represented by the formula (ω -9E), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(N-hydrocarbon-carbamoyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(N-heteroring-carbamoyl)-amino group".

[0215] Among the groups represented by the formula (ω -10E), those groups in which both R^{a5} and R^{b5} are hydrocarbon groups are referred to as "bis[N,N-di(hydrocarbon)-carbamoyl]-amino group", those groups in which both R^{a5} and R^{b5} are heterocyclic groups are referred to as "bis[N,N-di(heteroring)-carbamoyl]-amino group", groups in which R^{a5} is a hydrocarbon group and R^{b5} is a heterocyclic group are referred to as "bis(N-hydrocarbon-N-heteroring-carbamoyl)-amino group", and those groups in which R^{a5} and R^{b5} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino groups are referred to as "bis(cyclicamino-carbonyl)amino group".

[0216] Among the groups represented by the formula (ω -11E), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(N-hydrocarbon-thiocarbamoyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(N-heteroring-thiocarbamoyl)-amino group".

[0217] Among the groups represented by the formula (ω -12E), those groups in which both R^{a5} and R^{b5} are hydrocarbon groups are referred to as "bis[N,N-di(hydrocarbon)-thiocarbamoyl]-amino group", those groups in which both R^{a5} and R^{b5} are heterocyclic groups are referred to as "bis[N,N-di(heteroring)-thiocarbamoyl]-amino group", those groups in which R^{a5} is a hydrocarbon group and R^{b5} is a heterocyclic group are referred to as "bis(N-hydrocarbon-N-heteroring-thiocarbamoyl)-amino group", and those groups in which R^{a5} and R^{b5} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "bis(cyclicamino-thiocarbonyl)-amino group".

[0218] Among the groups represented by the formula (ω -13E), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(N-hydrocarbon-sulfamoyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(N-heteroring-sulfamoyl)-amino group".

[0219] Among the groups represented by the formula (ω -14E), those groups in which both R^{a5} and R^{b5} are hydrocarbon groups are referred to as "bis[N,N-di(hydrocarbon)-sulfamoyl]-amino group", those groups in which both R^{a5} and R^{b5} are heterocyclic groups are referred to as "bis[N,N-di(heteroring)-sulfamoyl]-amino group", those groups in which R^{a5} is a hydrocarbon group and R^{b5} is a heterocyclic group are referred to as "bis(N-hydrocarbon-N-heteroring-sulfamoyl)-amino group", and those groups in which R^{a5} and R^{b5} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "bis(cyclicamino-sulfonyl)amino group".

[0220] Among the groups represented by the formula (ω -15E), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(N-hydrocarbon-sulfinamoyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(N-heteroring-sulfinamoyl)-amino group".

[0221] Among the groups represented by the formula (ω -16E), those groups in which R^{a5} and R^{b5} are hydrocarbon groups are referred to as "bis[N,N-di(hydrocarbon)-sulfinamoyl]-amino group", those groups in which R^{a5} and R^{b5} are heterocyclic groups are referred to as "bis[N,N-di(heteroring)-sulfinamoyl]-amino group", those groups in which R^{a5} is a hydrocarbon group and R^{b5} is a heterocyclic group are referred to as "bis(N-hydrocarbon-N-heteroring-sulfinamoyl)-amino group", and those groups in which R^{a5} and R^{b5} combine to each other, together with the nitrogen atom to which they bind, to form a cyclic amino group are referred to as "bis(cyclicamino-sulfinyl)amino group".

[0222] Among the groups represented by the formula (ω -17E), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(hydrocarbon-oxy-sulfonyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(heteroring-oxy-sulfonyl)-amino group".

[0223] Among the groups represented by the formula (ω -18E), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(hydrocarbon-oxy-sulfinyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(heteroring-oxy-sulfinyl)-amino group".

[0224] Among the groups represented by the formula (ω -19E), those groups in which both R^{a5} and R^{b5} are hydrocarbon groups are referred to as "bis[O,O'-di(hydrocarbon)-phosphono]-amino group", those groups in which both R^{a5} and R^{b5} are heterocyclic groups are referred to as "bis[O,O'-di(heteroring)-phosphono]-amino group", and those groups

in which R^{a5} is a hydrocarbon group and R^{b5} is a heterocyclic group are referred to as "bis(O-hydrocarbon-O'-heteroring-phosphono)-amino group".

[0225] Among the groups represented by the formula (ω -20E), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(hydrocarbon-sulfonyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(heteroring-sulfonyl)-amino group".

[0226] Among the groups represented by the formula (ω -21E), those groups in which R^{a5} is a hydrocarbon group are referred to as "bis(hydrocarbon-sulfinyl)-amino group", and those groups in which R^{a5} is a heterocyclic group are referred to as "bis(heteroring-sulfinyl)-amino group".

[0227] Examples of the hydrocarbon in the groups represented by the aforementioned formulas (ω -1E through (ω -21E) include similar groups to the aforementioned hydrocarbon group. Examples of the bis(hydrocarbon-carbonyl)-amino groups represented by the formula (ω -1E) include, a bis(alkyl-carbonyl)-amino group, a bis(alkenyl-carbonyl)-amino group, a bis(alkynyl-carbonyl)-amino group, a bis(cycloalkyl-carbonyl)-amino group, a bis(cycloalkenyl-carbonyl)-amino group, a bis(cycloalkanedienyl-carbonyl)-amino group, a bis(cycloalkyl-alkyl-carbonyl)-amino group which is a bis(aliphatic hydrocarbon-carbonyl)-amino group, a bis(aryl-carbonyl)-amino group, a bis(aralkyl-carbonyl)-amino group, a bis(bridged cyclic hydrocarbon-carbonyl)-amino group, a bis(spiro cyclic hydrocarbon-carbonyl)-amino group, and a bis(terpene family hydrocarbon-carbonyl)-amino group. In the following, groups represented by the formulas (ω -2E) through (ω -21E) are similar to those explained above.

[0228] Examples of the heteroring in the groups represented by the aforementioned formulas (ω -1E) through (ω -21E) include similar groups to the aforementioned heterocyclic group. Examples of the bis(heteroring-carbonyl)-amino group represented by the formula (ω -1E) include, for example, bis(monocyclic heteroaryl-carbonyl)-amino group, bis(fused polycyclic heteroaryl-carbonyl)-amino group, bis(monocyclic nonaromatic heterocyclic-carbonyl)-amino group, and bis(fused polycyclic nonaromatic heterocyclic-carbonyl)-amino group. In the following, groups represented by the formulas (ω -2E) through (ω -21E) are similar to those groups mentioned above.

[0229] Examples of the cyclic amino in the groups represented by the aforementioned formulas (ω -10E) through (ω -16E) include similar groups to the aforementioned cyclic amino group.

[0230] The aforementioned acyl-amino group and di(acyl)-amino group are generically referred to as "acyl substituted amino group". Furthermore, the aforementioned N-hydrocarbon-amino group, N,N-di(hydrocarbon)-amino group, N-heterocyclic-amino group, N-hydrocarbon-N-heterocyclic-amino group, cyclic amino group, acyl-amino group, and di(acyl)-amino group are generically referred to as "substituted amino group". These substituted amino group and amino group are generically referred to as "amino groups which may be substituted".


[0231] In the following, compounds represented by the aforementioned general formulas (I), (I-1), (I-2), (I-3), (I-4) are explained in details.

[0232] "Connecting group whose number of atoms of main chain is 2 to 4" in the definition of X means connecting groups wherein 2 to 4 atoms in a main chain link together between rings Z and E. The aforementioned "number of atoms of the main chain" is counted so as to minimize the number of connecting atoms existing between the rings Z and E, regardless of the presence or absence of hetero atom(s). For example, the number of atoms of 1,2-cyclopentylene is counted as 2, the number of atoms of 1,3-cyclopentylene is counted as 3, the number of atoms of 1,4-phenylene is counted as 4, the number of atoms of 2,6-pyridine-diyl is counted as 3.

[0233] The aforementioned "connecting group whose number of atoms of main chain is 2 to 4" is formed by one functional group selected from the following group of divalent group ζ -1, or formed by combining 2 to 4 functional groups of 1 to 4 kinds selected from the following divalent group ζ -2.




[Divalent group ζ -1] the groups of the following formulas :

[0234]



[Divalent group ζ -2] the groups of the following formulas :

[0235]







When 2 or more divalent groups combine, each group may be the same or different.

[0236] The aforementioned "connecting group whose number of atoms of a main chain is 2 to 4" is preferably a group selected from the following "connecting group α ".


[Connecting group α] the following formulas :

[0237]




wherein the bond at the left end binds to ring Z and the bond at the right end binds to E.

[0238] The group represented by the following formula is most preferred:



wherein, the bond at the left end binds to ring Z and the bond at the right end binds to E.


[0239] Examples of the substituent, according to "connecting group which may be substituted" in the definition of "a connecting group whose number of atoms of the main chain is 2 to 4", include similar groups to the substituents in the definition of the aforementioned "which may be substituted". A C₁ to C₆ alkyl group is preferred, and a methyl group is more preferred. The substituent may combine with a substituent of the ring E or Z, together with atoms to which they bind, to form a cyclic group which may be substituted. Examples include the compounds represented by the general formula (I) being those represented by the following formulas:




[0240] In the aforementioned general formula (I), examples of A include hydrogen atom and an acetyl group, and hydrogen atom is preferred.

[0241] Examples of the "arene" in "an arene which may be substituted" in the definition of ring Z include a monocyclic or fused heterocyclic aromatic hydrocarbon, and include, for example, benzene ring, naphthalene ring, anthracene ring, phenanthrene ring, and acenaphylene ring. C₆ to C₁₀ arenes such as benzene ring, naphthalene ring and the like are preferred, benzene ring and naphthalene ring are more preferred, and benzene ring is most preferred.

[0242] When ring Z is a benzene ring, substituents according to the definition of "which may be substituted in addition to the group represented by formula -O-A wherein A has the same meaning as that defined in the general formula (I) and the group represented by formula -X-E wherein each of X and E has the same meaning as that defined in the general formula (I)" are preferred to locate on the position of R_Z when the following partial formula (Iz-1) in the general formula containing ring Z



is a group represented by the following formula (Iz-2).



(I z - 2)

[0243] Examples of the "hetero arene" in "a hetero arene which may be substituted" in the definition of ring Z include a monocyclic or a fused polycyclic aromatic heterocyclic rings containing at least one of 1 to 3 kinds of heteroatoms selected from oxygen atom, sulfur atom and nitrogen atom and the like as ring-constituting atoms (ring forming atoms), and include, for example, furan ring, thiophene ring, pyrrole ring, oxazole ring, isoxazole ring, thiazole ring, isothiazole ring, imidazole ring, pyrazole ring, 1,2,3-oxadiazole ring, 1,2,3-thiadiazole ring, 1,2,3-triazole ring, pyridine ring, pyridazine ring, pyrimidine ring, pyrazine ring, 1,2,3-triazine ring, 1,2,4-triazine ring, 1H-azepine ring, 1,4-oxepine ring, 1,4-thiazepine ring, benzofuran ring, isobenzofuran ring, benzo[b]thiophene ring, benzo[c]thiophene ring, indole ring, 2H-isoindole ring, 1H-indazole ring, 2H-indazole ring, benzooxazole ring, 1,2-benzoisoxazole ring, 2,1-benzoisoxazole ring, benzothiazole ring, 1,2-benzisothiazole ring, 2,1-benzisothiazole ring, 1,2,3-benzooxadiazole ring, 2,1,3-benzooxadiazole ring, 1,2,3-benzothiadiazole ring, 2,1,3-benzothiadiazole ring, 1H-benzotriazole ring, 2H-benzotriazole ring, quinoline ring, isoquinoline ring, cinnoline ring, quinoxaline ring, quinoxaline ring, phthalazine ring, naphthyridine ring, 1H-1,5-benzodiazepine ring, carbazole ring, α -carboline ring, β -carboline ring, γ -carboline ring, acridine ring, phenoxazine ring, phenothiazine ring, phenazine ring, phenanthridine ring, phenanthroline ring, thianthrene ring, indolizine ring, and phenoxathiine ring, which are 5 to 14 membered monocyclic or fused polycyclic aromatic heterocyclic rings. 6 to 13 membered monocyclic or fused polycyclic aromatic heterocyclic rings are preferred, and pyridine ring, indole ring, quinoxaline ring, and carbazole ring are more preferred.

[0244] Examples of the substituent in the definition of "which may be substituted in addition to the group represented by formula -O-A wherein A has the same meaning as that defined in the general formula (I) and the group represented by formula -X-E wherein each of X and E has the same meaning as that defined in the general formula (I)" in the definition of ring Z include similar groups to the substituent explained for the definition "which may be substituted". When ring Z is "a benzene ring which may be substituted in addition to the group represented by formula -O-A wherein A has the same meaning as that defined in the general formula (I) and the group represented by formula -X-E wherein each of X and E has the same meaning as that defined in the general formula (I)", preferred examples of the substituents include halogen atoms, nitro group, cyano group, hydroxy group which may be substituted, amino group which may be substituted, hydrocarbon group which may be substituted, heterocyclic group which may be substituted, acyl group which may be substituted, ureido group which may be substituted, thiureido group which may be substituted, and diazenyl group which may be substituted, which are defined as those of substituent group γ -1z.

[0245] Examples of the "hydroxy group which may be substituted" in the definition of the substituent group γ -1z, and the "hydroxy group which may be substituted" in the definition of R^z include similar groups to the "hydroxy group which may be substituted" according to the definition of the aforementioned "which may be substituted", and examples of the substituent include similar groups to the substituents in the definition of the aforementioned "which may be substituted". Hydrocarbon-oxy group which may be substituted is preferred as the "hydroxy group which may be substituted", a C_1 to C_6 alkoxy group which may be substituted is more preferred, and methoxy group is further preferred.

[0246] Examples of the "amino group which may be substituted" in the definition of the substituent group γ -1z and the "amino group which may be substituted" in the definition of R^z include similar group to the "amino group which may be substituted" according to the definition of the aforementioned "which may be substituted", and examples of the substituent include similar groups to the substituent according to the definition of the aforementioned "which may be substituted". Di(hydrocarbon)-amino group and hydrocarbon-carbonyl-amino group are preferred as the "amino group which may be substituted", di(C_1 to C_6 alkyl)-amino group and C_6 to C_{10} aryl-carbonyl-amino group are more preferred, and dimethylamino group and benzoylamino group are further preferred.

[0247] Examples of the substituent in the definition of "hydrocarbon group which may be substituted" in the definition of the substituent group γ -1z, and the substituent in the definition of "hydrocarbon group which may be substituted" in the definition of R^z include similar groups to the substituents according to the definition of the aforementioned "which may be substituted". A C_1 to C_6 alkyl group which may be substituted, a C_1 to C_6 halogenated alkyl group which may be substituted, a C_2 to C_6 alkenyl group which may be substituted, a C_2 to C_6 alkynyl group which may be substituted, a C_6 to C_{10} aryl group which may be substituted, and a C_7 to C_{16} aralkyl group which may be substituted are preferred as the "hydrocarbon group which may be substituted", and methyl group, tert-butyl group, 1-hydroxyethyl group, 1-(methoxyimino)ethyl group, 1-((benzyloxy)imino)ethyl group, trifluoromethyl group, pentafluoroethyl group, phenyl group,

4-(trifluoromethyl)phenyl group, 4-fluorophenyl group, 2,4-difluorophenyl group, 2-phenylethen-1-yl group, 2,2-dicyanoethen-1-yl group, 2-cyano-2-(methoxycarbonyl)ethen-1-yl group, 2-carboxy-2-cyanoethen-1-yl group, ethynyl group, phenylethynyl group, (trimethylsilyl)ethynyl group, phenyl group, and 2-phenethyl group are more preferred.

[0248] Examples of the substituent in the definition of "heterocyclic group which may be substituted" in the definition of the substituent group γ -1z, and the substituent in the definition of "heterocyclic group which may be substituted" in the definition of R^2 include similar groups to the substituents according to the definition of the aforementioned "which may be substituted". A heteroaryl group which may be substituted is preferred as the "heterocyclic group which may be substituted", a 5 to 6-membered heteroaryl group which may be substituted is more preferred, and 2-thienyl group, 3-thienyl group, 1-pyrrolyl group, 2-methylthiazol-4-yl group, and 2-pyridyl group are further preferred.

[0249] Examples of the "acyl group which may be substituted" in the definition of the substituent group γ -1z and the "acyl group which may be substituted" in the definition of R^2 include similar groups exemplified in the aforementioned definition of "acyl group which may be substituted", and examples of the substituent include similar groups to the substituent explained for the definition "which may be substituted". Carbamoyl group which may be substituted, sulfamoyl group which may be substituted, a hydrocarbon-carbonyl group which may be substituted, a hydrocarbon-oxy-carbonyl group which may be substituted, a heteroring-carbonyl group which may be substituted, and a heteroring-sulfonyl group which may be substituted are preferred as the "acyl group which may be substituted", carbamoyl group which may be substituted, sulfamoyl group which may be substituted, a C_1 to C_6 alkyl-carbonyl group which may be substituted, a C_1 to C_6 alkoxy-carbonyl group which may be substituted, a 5-membered heteroaryl-sulfonyl group which may be substituted, and a 6-membered nonaromatic heterocyclic-sulfonyl group which may be substituted are more preferred, and [3,5-bis(trifluoromethyl)phenyl]carbamoyl group, dimethylcarbamoyl group, dimethylsulfamoyl group, acetyl group, isobutyryl group, methoxycarbonyl group, piperidinocarbonyl group, 4-benzylpiperidinocarbonyl group, and (pyrrol-1-yl)sulfonyl group are further preferred.

[0250] Examples of the substituent in the definition of "ureido group which may be substituted" in the definition of the substituent group γ -1z, and the substituent in the definition of "ureido group which may be substituted" in the definition of R^2 include similar groups to the substituent explained for the definition "which may be substituted". 3-Phenylureido group is preferred as the "ureido group which may be substituted".

[0251] Examples of the substituent in the definition of "thioureido group which may be substituted" in the definition of the substituent group γ -1z and the substituent in the definition of "thioureido group which may be substituted" in the definition of R^2 include similar groups to the substituent explained for the definition "which may be substituted". (3-Phenylthio)ureido group is preferred as the "thioureido group which may be substituted".

[0252] Examples of the substituent in the definition of "diazanyl group which may be substituted" in the definition of the substituent group γ -1z, and the substituent in the definition of "diazanyl group which may be substituted" in the definition of R^2 include similar groups to the substituent explained for the definition "which may be substituted". (4-Nitrophenyl)diazanyl group and {[(4-pyridin-2-yl)sulfamoyl]-phenyl}diazanyl group are preferred as the "diazanyl group which may be substituted".

[0253] Examples of R^2 include halogen atom, nitro group, cyano group, hydroxy group which may be substituted, amino group which may be substituted, hydrocarbon group which may be substituted, heterocyclic group which may be substituted, acyl group which may be substituted, ureido group which may be substituted, thioureido group which may be substituted, and diazenyl group which may be substituted, and halogen atom is most preferred.

[0254] Examples of the aryl group of "an aryl group which may be substituted" in the definition of E include similar groups to the aryl group in the definition of the aforementioned "hydrocarbon group", and C_6 to C_{10} aryl groups such as phenyl group, 1-naphthyl group, 2-naphthyl group and the like are preferred, and phenyl group is most preferred.

[0255] Examples of the substituent in the definition of "an aryl group which may be substituted" in the definition of E include similar groups to the substituent explained for the definition "which may be substituted".

[0256] Preferred embodiments of the phenyl group according to "an aryl group which may be substituted" in the definition of E are:

- (1) phenyl group substituted with two C_1 to C_6 halogenated alkyl groups wherein said phenyl group may be substituted in addition to the two C_1 to C_6 halogenated alkyl groups;
- (2) phenyl group substituted with one C_1 to C_6 halogenated alkyl group wherein said phenyl group may be substituted (except with a C_1 to C_6 halogenated alkyl group) in addition to the C_1 to C_6 halogenated alkyl group; and
- (3) phenyl group which may be substituted (except with a C_1 to C_6 halogenated alkyl group).

[0257] Examples of the " C_1 to C_6 halogenated alkyl group" in "phenyl group substituted with two C_1 to C_6 halogenated alkyl groups wherein said phenyl group may be substituted in addition to the two C_1 to C_6 halogenated alkyl groups" in the definition of E include similar groups to those exemplified in the aforementioned definition of " C_1 to C_6 halogenated alkyl group", and examples of the substituent in the definition of "said phenyl group may be substituted in addition to the two C_1 to C_6 halogenated alkyl groups" include similar groups to the substituent explained for the definition "which

may be substituted".

[0258] "Phenyl group substituted with two C₁ to C₆ halogenated alkyl groups" is preferred as the "phenyl group substituted with two C₁ to C₆ halogenated alkyl groups wherein said phenyl group may be substituted in addition to the two C₁ to C₆ halogenated alkyl groups". 3,5-Bis(trifluoromethyl)phenyl group and 2,5-bis(trifluoromethyl)phenyl group are preferred, and 3,5-bis(trifluoromethyl)phenyl group is most preferred.

[0259] Examples of the substituent in the definition of "phenyl group substituted with one C₁ to C₆ halogenated alkyl group wherein said phenyl group may be substituted (except with a C₁ to C₆ halogenated alkyl group) in addition to one C₁ to C₆ halogenated alkyl group" in the definition of E include similar groups to the substituent explained for the definition "which may be substituted" (a C₁ to C₆ halogenated alkyl group is excluded). Halogen atoms, nitro group, cyano group, hydroxy group which may be substituted, hydrocarbon group which may be substituted, heterocyclic group which may be substituted, sulfanyl group which may be substituted which are defined in substituent group γ-1e are preferred.

[0260] Examples of the "hydroxy group which may be substituted" in the definition of the substituent group γ-1e include similar groups to the "hydroxy group which may be substituted" in the definition of the aforementioned "which may be substituted", and examples of the substituent include similar groups to the substituent explained for the definition "which may be substituted". Hydrocarbon-oxy group which may be substituted is preferred as the "hydroxy group which may be substituted", C₁ to C₆ alkoxy group which may be substituted which is defined as substituent group γ-2e is more preferred, and methoxy group is further preferred.

[0261] Examples of the substituent in the definition of "hydrocarbon group which may be substituted" in the definition of the substituent group γ-1e, and the substituent in the definition of "hydrocarbon group which may be substituted" in the definition of R² include similar groups to the substituent explained for the definition "which may be substituted". A C₁ to C₆ alkyl group which may be substituted which is defined in substituent group γ-2e is preferred as the "hydrocarbon group which may be substituted", and methyl group is more preferred.

[0262] Examples of the substituent in the definition of "heterocyclic group which may be substituted" in the definition of the substituent group γ-1e include similar groups to the substituents explained for the definition "which may be substituted". A 5 to 6-membered non-aromatic heterocyclic group which may be substituted which is defined in substituent group γ-2e is preferred as the "heterocyclic group which may be substituted", and 1-pyrrolidinyl group and morpholino group are more preferred.

[0263] Examples of the "sulfanyl group which may be substituted" in the definition of the substituent group γ-1e include similar groups to the "sulfanyl group which may be substituted" according to the definition of the aforementioned "which may be substituted", and examples of the substituent include similar groups to the substituent explained for the definition "which may be substituted". A hydrocarbon-sulfanyl group which may be substituted is preferred as the "sulfanyl group which may be substituted", and a C₁ to C₆ alkyl-sulfanyl group which may be substituted, which is defined in substituent group γ-2e, is more preferred, and methylsulfanyl group is further preferred.

[0264] Examples of the "C₁ to C₆ halogenated alkyl group" in "phenyl group substituted with one C₁ to C₆ halogenated alkyl group wherein said phenyl group may be substituted (except with a C₁ to C₆ halogenated alkyl group) in addition to the C₁ to C₆ halogenated alkyl group" in the definition of E include similar groups to the aforementioned "C₁ to C₆ halogenated alkyl group". C₁ to C₆ alkyl groups substituted with one or more fluorine atoms are preferred, C₁ to C₆ alkyl groups substituted with three or more fluorine atoms are more preferred, and trifluoromethyl group is most preferred.

[0265] Examples of the "phenyl group substituted with one C₁ to C₆ halogenated alkyl group wherein said phenyl group may be substituted (except with a C₁ to C₆ halogenated alkyl group) in addition to the C₁ to C₆ halogenated alkyl group" in the definition of E include 2-(trifluoromethyl)phenyl group, 3-(trifluoromethyl)phenyl group, 4-(trifluoromethyl)phenyl group, 2-fluoro-3-(trifluoromethyl)phenyl group, 2-chloro-4-(trifluoromethyl)phenyl group, 2-fluoro-5-(trifluoromethyl)phenyl group, 2-chloro-5-(trifluoromethyl)phenyl group, 3-fluoro-5-(trifluoromethyl)phenyl group, 3-bromo-5-(trifluoromethyl)phenyl group, 4-chloro-2-(trifluoromethyl)phenyl group, 4-fluoro-3-(trifluoromethyl)phenyl group, 4-chloro-3-(trifluoromethyl)phenyl group, 2-nitro-5-(trifluoromethyl)phenyl group, 4-nitro-3-(trifluoromethyl)phenyl group, 4-cyano-3-(trifluoromethyl)phenyl group, 2-methyl-3-(trifluoromethyl)phenyl group, 2-methyl-5-(trifluoromethyl)phenyl group, 4-methyl-3-(trifluoromethyl)phenyl group, 2-methoxy-5-(trifluoromethyl)phenyl group, 3-methoxy-5-(trifluoromethyl)phenyl group, 4-methoxy-3-(trifluoromethyl)phenyl group, 2-(methylsulfanyl)-5-(trifluoromethyl)phenyl group, 2-(1-pyrrolidino)-5-(trifluoromethyl)phenyl group, and 2-morpholino-5-(trifluoromethyl)phenyl group. 2-Chloro-5-(trifluoromethyl)phenyl group, 4-chloro-3-(trifluoromethyl)phenyl group, 2-methoxy-5-(trifluoromethyl)phenyl group, and 3-methoxy-5-(trifluoromethyl)phenyl group are more preferred, and 2-chloro-5-(trifluoromethyl)phenyl group is most preferred.

[0266] Examples of the substituent in the definition of "phenyl group which may be substituted (except with a C₁ to C₆ halogenated alkyl group)" in the definition of E include similar groups to the substituent explained for the definition "which may be substituted". Halogen atoms, nitro group, hydroxy group which may be substituted, hydrocarbon group which may be substituted, and acyl group which may be substituted which are defined in substituent group γ-3e are

preferred.

[0267] Examples of the "hydroxy group which may be substituted" in the definition of the substituent group γ -3e include similar groups to the "hydroxy group which may be substituted" according to the definition of the aforementioned "which may be substituted", and examples of the substituent include similar groups to the substituent explained for the definition "which may be substituted". Unsubstituted hydroxy group and a hydrocarbon-oxy group which may be substituted are preferred as the "hydroxy group which may be substituted", and unsubstituted hydroxy group and C_1 to C_6 alkoxy group which may be substituted which are defined in substituent group γ -4e are more preferred, and unsubstituted hydroxy group and methoxy group are further preferred.

[0268] Examples of the substituent in the definition of "hydrocarbon group which may be substituted" in the definition of the substituent group γ -3e include similar groups to the substituent explained for the definition "which may be substituted". A C_1 to C_6 alkyl group which may be substituted, a C_6 to C_{10} aryl group which may be substituted, and a C_1 to C_6 alkylene group which may be substituted which are defined in substituent group γ -4e are preferred as the "hydrocarbon group which may be substituted", and methyl group, tert-butyl group and 1,1,4,4-tetramethylbutane-1,4-diyl group are more preferred.

[0269] Examples of the "acyl group which may be substituted" in the definition of the substituent group γ -3e include the similar group to those exemplified in the aforementioned definition of "acyl group which may be substituted", and examples of the substituent include similar groups to the substituents explained for the definition "which may be substituted". A hydrocarbon-carbonyl group which may be substituted and a hydrocarbon-oxy-carbonyl group which may be substituted are preferred as the "acyl group which may be substituted". A C_1 to C_6 alkyl-carbonyl group which may be substituted, a C_1 to C_6 alkoxy-carbonyl group which may be substituted which are defined in substituent group γ -4e are more preferred, and acetyl group and methoxycarbonyl group are further preferred.

[0270] Phenyl group, 3-chlorophenyl group, 4-chlorophenyl group, 2,5-dichlorophenyl group, 3,4-dichlorophenyl group, 3,5-difluorophenyl group, 3,5-dichlorophenyl group, 3,4,5-trichlorophenyl group, pentafluorophenyl group, 3,5-dinitrophenyl group, 3,5-dichloro-4-hydroxyphenyl group, 2,5-dimethoxyphenyl group, 3,5-dimethoxyphenyl group, 3,5-dimethylphenyl group, 2,5-bis[(1,1-dimethyl)ethyl]phenyl group, 3,5-bis[(1,1-dimethyl)ethyl]phenyl group, 5-(1,1-dimethyl)ethyl-2-methoxyphenyl group, 3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydronaphthalen-2-yl group, biphenyl-3-yl group, 4-methoxybiphenyl-3-yl group, 3-acetylphenyl group, and 3,5-bis(methoxycarbonyl)phenyl group are preferred as "phenyl group which may be substituted (except with a C_1 to C_6 halogenated alkyl group)" in the definition of E. 2,5-Bis[(1,1-dimethyl)ethyl]phenyl group, 3,5-bis[(1,1-dimethyl)ethyl]phenyl group, and 5-(1,1-dimethyl)ethyl-2-methoxyphenyl group are more preferred, and 2-chloro-5-(trifluoromethyl)phenyl group is most preferred.

[0271] Examples of the "heteroaryl group" in "heteroaryl group which may be substituted" in the definition of E include similar groups to the "monocyclic heteroaryl group" and "fused polycyclic heteroaryl group" in the definition of the aforementioned "heteroaryl group". A 5 to 13-membered heteroaryl group is preferred, and thienyl group, pyrazolyl group, oxazolyl group, thiazolyl group, thiadiazolyl group, pyridyl group, pyrimidinyl group, indolyl group, and carbazolyl group are more preferred, and thiazolyl group is most preferred.

[0272] Examples of the substituent in the definition of "heteroaryl group which may be substituted" in the above definition of E include similar groups to the substituent explained for the definition "which may be substituted".

[0273] Examples of the substituent in the definition of "thiazolyl group which may be substituted" in the above definition of E include similar groups to the substituent explained for the definition "which may be substituted". Halogen atoms, cyano group, hydrocarbon group which may be substituted, heterocyclic group which may be substituted, and acyl group which may be substituted which are defined as substituent group γ -5e are preferred.

[0274] Examples of the substituent in the definition of "hydrocarbon group which may be substituted" in the definition of the substituent group γ -5e include similar groups to the substituents explained for the definition "which may be substituted". A C_1 to C_6 alkyl group which may be substituted, a C_1 to C_6 halogenated alkyl group which may be substituted, a C_6 to C_{10} aryl group which may be substituted, and a C_7 to C_{16} aralkyl group which may be substituted which are defined in substituent group γ -6e are preferred as the "hydrocarbon group which may be substituted". Methyl group, ethyl group, isopropyl group, n-butyl group, tert-butyl group, carboxymethyl group, trifluoromethyl group, phenyl group, 4-fluorophenyl group, 3-(trifluoromethyl)phenyl group, pentafluorophenyl group, and benzyl group are preferred.

[0275] Examples of the substituent in the definition of "heterocyclic group which may be substituted" in the definition of the substituent group γ -5e include similar groups to the substituent explained for the definition "which may be substituted". A 6-membered non-aromatic heterocyclic group which may be substituted, which is defined in substituent group γ -6e, is preferred as the "heterocyclic group which may be substituted", and piperidino group, morpholino group, 4-methylpiperidin-1-yl group, and 4-phenylpiperidin-1-yl group are more preferred.

[0276] Examples of the "acyl group which may be substituted" in the definition of the substituent group γ -5e include similar groups to those exemplified in the aforementioned definition of "acyl group which may be substituted", and examples of the substituent include similar groups to the substituents explained for the definition "which may be substituted". A hydrocarbon-carbonyl group which may be substituted, a carbamoyl group which may be substituted, and a hydrocarbon-oxy-carbonyl group which may be substituted are preferred as the "acyl group which may be substituted".

Carbamoyl group which may be substituted, a C₁ to C₆ alkyl-carbonyl group which may be substituted, a C₆ to C₁₀ aryl-carbonyl group which may be substituted, a C₁ to C₆ alkoxy-carbonyl group which may be substituted which are defined in substituent group γ-6e are more preferred, and N-methylcarbamoyl group, N-ethylcarbamoyl group, N-isopropylcarbamoyl group, N-(2-phenethyl)carbamoyl group, acetyl group, pivaloyl group, benzoyl group, and ethoxycarbonyl group are further preferred.

[0277] 5-Bromo-4-[(1,1-dimethyl)ethyl]thiazol-2-yl group, 5-bromo-4-(trifluoromethyl)thiazol-2-yl group, 5-cyano-4-[(1,1-dimethyl)ethyl]thiazol-2-yl group, 4-[(1,1-dimethyl)ethyl]thiazol-2-yl group, 5-phenyl-4-(trifluoromethyl)thiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-ethylthiazol-2-yl group, 5-methyl-4-phenylthiazol-2-yl group, 4-isopropyl-5-phenylthiazol-2-yl group, 4-benzyl-5-phenylthiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl group, 5-acetyl-4-phenylthiazol-2-yl group, 5-benzoyl-4-phenylthiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-(ethoxycarbonyl)thiazol-2-yl group, 5-ethoxycarbonyl-4-(trifluoromethyl)thiazol-2-yl group, 5-ethoxycarbonyl-4-phenylthiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-piperidinethiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-morpholinethiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-(4-phenylpiperidin-1-yl)thiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-(4-methylpiperidin-1-yl)thiazol-2-yl group, 4,5-diphenylthiazol-2-yl group, 4-phenylthiazol-2-yl group, 4,5-dimethylthiazol-2-yl group, 2-thiazolyl group, 5-methylthiazol-2-yl group, 4-ethyl-5-phenylthiazol-2-yl group, 5-carboxymethyl-4-phenylthiazol-2-yl group, 5-methylcarbamoyl-4-phenylthiazol-2-yl group, 5-ethylcarbamoyl-4-phenylthiazol-2-yl group, 5-isopropylcarbamoyl-4-phenylthiazol-2-yl group, 5-(2-phenethyl)carbamoyl-4-phenylthiazol-2-yl group, 4-(n-butyl)-5-phenylthiazol-2-yl group, 4-methyl-5-[(3-trifluoromethyl)phenyl]thiazol-2-yl group, and 5-(4-fluorophenyl)-4-methylthiazol-2-yl group are preferred as "thiazolyl group which may be substituted" in the definition of E, and 4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl group is more preferred.

[0278] In the aforementioned general formula (I-1), Z¹ is 2-hydroxyphenyl group which may be substituted in the 5-position or 2-acetoxyphenyl group which may be substituted in the 5-position.

[0279] Examples of the "halogenated alkyl group" in the "phenyl group substituted with two C₁ to C₆ halogenated alkyl groups wherein said phenyl group may be substituted in addition to the two C₁ to C₆ halogenated alkyl groups" in the definition of E¹ include similar group to those exemplified in the aforementioned definition of "halogenated alkyl group". Examples of substituents according to the definition of "said phenyl group may be substituted in addition to the two C₁ to C₆ halogenated alkyl groups" include similar groups to the substituents explained for the definition "which may be substituted". As the "phenyl group substituted with two C₁ to C₆ halogenated alkyl groups wherein said phenyl group may be substituted in addition to the two C₁ to C₆ halogenated alkyl groups", an example is a phenyl group substituted with two C₁ to C₆ alkyl groups which is substituted with one or more fluorine atoms in which said phenyl group may further have substituents in addition to the two C₁ to C₆ alkyl groups substituted with one or more fluorine atoms. More preferred example is a phenyl group substituted with two C₁ to C₆ alkyl groups which is substituted with three or more fluorine atoms in which said phenyl group may further have substituents in addition to the two C₁ to C₆ halogenated alkyl groups which are substituted with one or more fluorine atoms, and a phenyl group substituted with the two C₁ to C₆ alkyl groups which are substituted with three or more fluorine atoms is further preferred. It is preferred that these two substituents are substituted in the 2-position and 5-position, or 3-position and 5-position on the phenyl group.

[0280] Trifluoromethyl group is most preferred as "C₁ to C₆ alkyl group which is substituted with three or more fluorine atoms" in the definition of R^{1e2}, R^{1e3}, and R^{1e5}.

[0281] As E¹, 3,5-bis(trifluoromethyl)phenyl group or 2,5-bis(trifluoromethyl)phenyl group is preferred, and 2,5-bis(trifluoromethyl)phenyl group is most preferred.

[0282] A¹ is hydrogen atom and an acetyl group, and hydrogen atom is preferred.

[0283] Examples of the "hydroxy group which may be substituted" in the definition of R^{1z} include similar groups to the "hydroxy group which may be substituted" in the definition of the aforementioned "which may be substituted", and examples of the substituent include similar groups to the substituent explained for the definition "which may be substituted". A hydrocarbon-oxy group which may be substituted is preferred as the "hydroxy group which may be substituted". A C₁ to C₆ alkoxy group which may be substituted is more preferred, and methoxy group is further preferred.

[0284] Examples of the "amino group which may be substituted" in the definition of R^{1z} include similar groups to the "amino group which may be substituted" in the definition of the aforementioned "which may be substituted", and examples of the substituent include similar groups to the substituent explained for the definition "which may be substituted". Di(hydrocarbon)-amino group and hydrocarbon-carbonyl-amino group are preferred as the "amino group which may be substituted", and di(C₁ to C₆ alkyl)-amino group and C₆ to C₁₀ aryl-amino group are more preferred, and dimethylamino group and benzoylamino group are further preferred.

[0285] Examples of the substituent in the definition of "hydrocarbon group which may be substituted" in the definition of R^{1z} include similar groups to the substituent explained for the definition "which may be substituted". A C₁ to C₆ alkyl group which may be substituted, a C₁ to C₆ halogenated alkyl group which may be substituted, a C₂ to C₆ alkenyl group which may be substituted, a C₂ to C₆ alkynyl group which may be substituted, a C₆ to C₁₀ aryl group which may be substituted, and a C₇ to C₁₆ aralkyl group which may be substituted are preferred as the "hydrocarbon group which

may be substituted", and methyl group, tert-butyl group, 1-hydroxyethyl group, 1-(methoxyimino)ethyl group, 1-((benzyloxy)imino)ethyl group, trifluoromethyl group, pentafluoroethyl group, phenyl group, 4-(trifluoromethyl)phenyl group, 4-fluorophenyl group, 2,4-difluorophenyl group, 2-phenylethen-1-yl group, 2,2-dicyanoethen-1-yl group, 2-cyano-2-(methoxycarbonyl)ethen-1-yl group, 2-carboxy-2-cyanoethen-1-yl group, ethynyl group, phenylethynyl group, (trimethylsilyl)ethynyl group, and 2-phenethyl group are more preferred.

[0286] Examples of the substituent in the definition of "heterocyclic group which may be substituted" in the definition of R^{1z} include similar groups to the substituent explained for the definition "which may be substituted". A heteroaryl group which may be substituted is preferred as the "heterocyclic group which may be substituted", and a 5 to 6-membered heteroaryl group which may be substituted is more preferred, and 2-thienyl group, 3-thienyl group, 1-pyrrolyl group, 2-methylthiazol-4-yl group, and 2-pyridyl group are further preferred.

[0287] Examples of the "acyl group which may be substituted" in the definition of R^{1z} include similar groups to those exemplified in the aforementioned definition of "acyl group which may be substituted", and examples of the substituent include similar groups to the substituents explained for the definition "which may be substituted". Carbamoyl group which may be substituted, sulfamoyl group which may be substituted, a hydrocarbon-carbonyl group which may be substituted, a hydrocarbon-oxy-carbonyl group which may be substituted, a heterocyclic-carbonyl group which may be substituted, and a heterocyclic-sulfonyl group which may be substituted are preferred as the "acyl group which may be substituted", and carbamoyl group which may be substituted, sulfamoyl group which may be substituted, a C_1 to C_6 alkyl-carbonyl group which may be substituted, a C_1 to C_6 alkoxy-carbonyl group which may be substituted, a 5-member heteroaryl-sulfonyl group which may be substituted, and a 6-membered nonaromatic heterocyclic-sulfonyl group which may be substituted are more preferred. 3,5-Bis(trifluoromethyl)phenyl]carbamoyl group, dimethylcarbamoyl group, dimethylsulfamoyl group, acetyl group, isobutyryl group, methoxycarbonyl group, piperidinocarbonyl group, 4-benzylpiperidinocarbonyl group, and (pyrrol-1-yl)sulfonyl group are further preferred.

[0288] Examples of the substituent in the definition of "ureido group which may be substituted" in the definition R^{1z} include similar groups to the substituents explained for the definition "which may be substituted". 3-Phenylureido group is preferred as the "ureido group which may be substituted".

[0289] Examples of the substituent in the definition of "thioureido group which may be substituted" in the definition of R^{1z} include similar groups to the substituents explained for the definition "which may be substituted". (3-Phenylthio)ureido group is preferred as the "thioureido group which may be substituted".

[0290] Examples of the substituent in the definition of "diazenyl group which may be substituted" in the definition of R^{1z} include similar groups to the substituents explained for the definition "which may be substituted". (4-Nitrophenyl)diazenyl group and (((4-pyridin-2-yl)sulfamoyl)phenyl)diazenyl group are preferred as the "diazenyl group which may be substituted".

[0291] Examples of R^{1z} include a halogen atom, nitro group cyano group, hydroxy group which may be substituted, amino group which may be substituted, a hydrocarbon group which may be substituted, a heterocyclic group which may be substituted, an acyl group which may be substituted, an ureido group which may be substituted, a thioureido group which may be substituted, a diazenyl group which may be substituted, a halogen atom, a C_1 to C_6 alkyl group which may be substituted, and a C_1 to C_6 halogenated alkyl group which may be substituted are preferred, and a halogen atom is most preferred.

[0292] Each compound defined by the aforementioned general formula (I-1) or a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof, except the following 6 compounds, is novel compound, and uses of the compounds according to the present invention relating to the chemical substances are not limited.

- N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide
- N-[3,5-bis(trifluoromethyl)phenyl]-5-chloro-2-hydroxybenzamide
- N-[3,5-bis(trifluoromethyl)phenyl]-5-bromo-2-hydroxybenzamide
- N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-iodobenzamide
- N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-nitrobenzamide
- 2-Hydroxy-N-[2,3,5-tris(trifluoromethyl)phenyl]benzamide

[0293] In the aforementioned general formula (1-2), Z^2 is 2-hydroxyphenyl group which may be substituted in the 5-position or 2-acetoxyphenyl group which may be substituted at the 5-position.

[0294] Examples of the "halogenated alkyl group" in "phenyl group wherein a C_1 to C_6 halogenated alkyl group is substituted in the 3-position or 5-position" in the definition of E^2 (wherein said phenyl group may further have one or more substituents (except when the substituent is C_1 to C_6 halogenated alkyl group) in addition to the C_1 to C_6 halogenated alkyl group in the 3-position or 5-position) include similar groups to those exemplified in the aforementioned definition of "halogenated alkyl group". A C_1 to C_6 alkyl group substituted with one or more fluorine atoms is preferred, and a C_1 to C_6 alkyl group substituted with three or more fluorine atoms is more preferred, and trifluoromethyl group is most preferred. Examples of the substituent in the definition of "phenyl group wherein a C_1 to C_6 halogenated alkyl

group is substituted in the 3-position or 5-position" (wherein said phenyl group may further have one or more substituents (except when the substituent is C₁ to C₆ halogenated alkyl group) in addition to the C₁ to C₆ halogenated alkyl group in the 3-position or 5-position) include similar groups to the substituents explained for the definition "which may be substituted". A halogen atom, nitro group, cyano group, a C₁ to C₆ alkyl group which may be substituted, a 5 to 6 membered nonaromatic heterocyclic group which may be substituted, a C₁ to C₆ alkoxy group which may be substituted, a C₁ to C₆ alkyl-sulfanyl group which may be substituted, which are defined in substituent group γ -7e, are preferred, and a halogen atom, nitro group, cyano group, methoxy group, methyl group, 1-pyrrolidinyl group, morpholino group, and methyl sulfanyl group are more preferred.

[0295] Preferred examples of the "phenyl group wherein a C₁ to C₆ halogenated alkyl groups is substituted in the 3-position or 5-position (wherein said phenyl group may further have one or more substituents (except when the substituent is a C₁ to C₆ halogenated alkyl) in addition to the C₁ to C₆ halogenated alkyl group in the 3-position or 5-position))" in the definition of E² include 3-(Trifluoromethyl)phenyl group, 2-fluoro-3-(trifluoromethyl)phenyl group, 2-fluoro-5-(trifluoromethyl)phenyl group, 2-chloro-5-(trifluoromethyl)phenyl group, 3-fluoro-5-(trifluoromethyl)phenyl group, 3-bromo-5-(trifluoromethyl)phenyl group, 4-chloro-2-(trifluoromethyl)phenyl group, 4-fluoro-3-(trifluoromethyl)phenyl group, 4-chloro-3-(trifluoromethyl)phenyl group, 2-nitro-5-(trifluoromethyl)phenyl group, 4-nitro-3-(trifluoromethyl)phenyl group, 4-cyano-3-(trifluoromethyl)phenyl group, 2-methyl-3-(trifluoromethyl)phenyl group, 2-methyl-5-(trifluoromethyl)phenyl group, 4-methyl-3-(trifluoromethyl)phenyl group, 2-methoxy-5-(trifluoromethyl)phenyl group, 3-methoxy-5-(trifluoromethyl)phenyl group, 4-methoxy-3-(trifluoromethyl)phenyl group, 2-(methylsulfanyl)-5-(trifluoromethyl)phenyl group, 2-(1-pyrrolidino)-5-(trifluoromethyl)phenyl group, and 2-morpholino-5-(trifluoromethyl)phenyl group.

[0296] A² is hydrogen atom and an acetyl group, and hydrogen atom is preferred.

[0297] Examples of R^{2z} include a halogen atom, a C₁ to C₆ alkyl group which may be substituted, and a C₁ to C₆ halogenated alkyl group which may be substituted, and preferred examples include a halogen atom, methyl group, tert-butyl group, trifluoromethyl group, and pentafluoromethyl group.

[0298] Each compound defined by the aforementioned general formula (1-2) or a pharmacologically acceptable salt thereof, or a hydrate thereof or a solvate thereof, except the following 15 compounds, is novel, and uses of the compounds according to the present invention relating to chemical substances are not limited.

- 5-chloro-2-hydroxy-N-[3-(trifluoromethyl)phenyl]benzamide
- 5-bromo-2-hydroxy-N-[3-(trifluoromethyl)phenyl]benzamide
- 2-hydroxy-5-iodo-N-[3-(trifluoromethyl)phenyl]benzamide
- 5-chloro-N-[4-chloro-3-(trifluoromethyl)phenyl]-2-hydroxybenzamide
- 5-chloro-N-[5-chloro-3-(trifluoromethyl)phenyl]-2-hydroxybenzamide
- 5-chloro-2-hydroxy-N-[4-nitro-3-(trifluoromethyl)phenyl]benzamide
- 5-fluoro-2-hydroxy-N-[2-(2,2,2-trifluoroethoxy)-5-(trifluoromethyl)phenyl]benzamide
- 5-fluoro-2-hydroxy-N-[2-(6,6,6-trifluorohexyloxy)-5-(trifluoromethyl)phenyl]benzamide
- 5-chloro-2-hydroxy-N-[3-(trifluoromethyl)-4-([4-(trifluoromethyl)sulfanyl]phenoxy)phenyl]benzamide
- N-[4-(benzothiazol-2-yl)sulfanyl-3-(trifluoromethyl)phenyl]-5-chloro-2-hydroxybenzamide
- 5-chloro-N-[2-(4-chlorophenoxy)-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide
- 5-chloro-2-hydroxy-N-[2-(4-methylphenoxy)-5-(trifluoromethyl)phenyl]benzamide
- 5-chloro-N-[2-(4-chlorophenyl)sulfanyl-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide
- 5-chloro-2-hydroxy-N-[2-(1-naphthyloxy)-5-(trifluoromethyl)phenyl]benzamide
- 5-chloro-2-hydroxy-N-[2-(2-naphthyloxy)-5-(trifluoromethyl)phenyl]benzamide

[0299] In the aforementioned general formula(1-3), Z³ is 2-hydroxyphenyl group which may be substituted in the 5-position and 2-acetoxyphenyl group which may be substituted in the 5-position.

[0300] Examples of the substituent in the definition of "hydrocarbon group which may be substituted" in the definition of R^{3e2} and R^{3e3} include similar groups to the substituents explained for the definition "which may be substituted". A C₁ to C₆ alkyl group is preferred as the "hydrocarbon group which may be substituted", and tert-butyl group is most preferred.

[0301] Examples of the substituent in the definition of "hydroxy group which may be substituted" in the definition of R^{3e2} and R^{3e3} include similar groups to the substituents explained for the definition "which may be substituted". A C₁ to C₆ alkoxy group is preferred as the "hydroxy group which may be substituted", and methoxy group is most preferred.

[0302] Examples of the substituent in the definition of "C₂ to C₆ hydrocarbon group which may be substituted" in the definition of R^{3e5} include similar groups to the substituents explained for the definition "which may be substituted". A C₂ to C₆ alkyl group is preferred as the "C₂ to C₆ alkyl group which may be substituted", and tert-butyl group is most preferred.

[0303] As E³, 2,5-bis[(1,1-dimethyl)ethyl]phenyl group, 3,5-bis[(1,1-dimethyl)ethyl]-phenyl group and 5-(1,1-dimethyl)phenyl group are preferred.

thyl)ethyl-2-methoxyphenyl group are preferred.

[0304] A³ is hydrogen atom and an acetyl group, and hydrogen atom is preferred.

[0305] Examples of R^{3z} include halogen atom, C₁ to C₆ alkyl group and C₁ to C₆ halogenated alkyl group, and halogen atom, methyl group, tert-butyl group, trifluoromethyl group, and pentafluoroethyl group are preferred.

[0306] Each compound defined by the aforementioned general formula (I-3) or a pharmacologically acceptable salt thereof, or a hydrate thereof and a solvate thereof is novel, and uses of the compounds according to the present invention relating to chemical substances are not limited.

[0307] In the aforementioned general formula (I-4), Z⁴ includes 2-hydroxyphenyl group which may be substituted in the 5-position and 2-acetoxyphenyl group which may be substituted in the 5-position.

[0308] Examples of the substituent in the definition of "hydrocarbon group which may be substituted" in the definition of R^{4e4} include similar groups to the substituent explained for the definition "which may be substituted". A C₁ to C₆ alkyl group which may be substituted, a C₁ to C₆ halogenated alkyl group which may be substituted, and a C₆ to C₁₀ aryl group which may be substituted are preferred as the "hydrocarbon group which may be substituted", and methyl group, isopropyl group, tert-butyl group, phenyl group, and pentafluorophenyl are more preferred.

[0309] Examples of the substituent in the definition of "acyl group which may be substituted" in the definition of R^{4e5} include similar groups to the substituents explained for the definition "which may be substituted". A C₁ to C₆ alkyl-carbonyl group which may be substituted, a C₆ to C₁₀ aryl-carbonyl group which may be substituted, and a C₁ to C₆ alkoxy-carbonyl group which may be substituted are preferred as the "acyl group which may be substituted", and acetyl group, pivaloyl group and benzoyl group are more preferred.

[0310] Examples of the substituent in the definition of "heterocyclic group which may be substituted" in the definition of R^{4e5} include similar groups to the substituents explained for the definition "which may be substituted". A 6-membered nonaromatic heterocyclic group which may be substituted is preferred as the "heterocyclic group which may be substituted", and piperidino group, morpholino group, 4-methylpiperazin-1-yl group, 4-phenylpiperazin-1-yl group are more preferred.

[0311] As E⁴, 5-bromo-4-[(1,1-dimethyl)ethyl]thiazol-2-yl group, 5-bromo-4-(trifluoromethyl)thiazol-2-yl group, 5-cyano-4-[(1,1-dimethyl)ethyl]thiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl group, 5-acetyl-4-phenylthiazol-2-yl group, 5-benzoyl-4-phenylthiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-(ethoxycarbonyl)thiazol-2-yl group, 5-ethoxycarbonyl-4-(trifluoromethyl)thiazol-2-yl group, 5-ethoxycarbonyl-4-phenylthiazol-2-yl group, 5-ethoxycarbonyl-4-(pentafluorophenyl)thiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-piperidinethiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-morpholinethiazol-2-yl group, 4-(1,1-dimethyl)ethyl-5-(4-methylpiperidin-1-yl)thiazol-2-yl group, and 4-(1,1-dimethyl)ethyl-5-(4-phenylpiperidin-1-yl)thiazol-2-yl group are preferred, and 4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl group is most preferred.

[0312] A⁴ includes hydrogen atom and an acetyl group, and hydrogen atom is preferred.

[0313] Examples of R^{4z} include a halogen atom, a C₆ to C₁₀ aryl group, and a 5-membered heteroaryl group, and a halogen atom, phenyl group, 4-fluorophenyl group, 2,4-difluorophenyl group, 4-(trifluoromethyl)phenyl group, 1-pyrrolyl group, and 2-thienyl group are preferred.


[0314] Each compound defined by the aforementioned general formula (I-4) or a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof is novel, and uses of the compounds according to the present invention relating to chemical substances are not limited.

[0315] The compounds represented by the aforementioned general formulas (I), (I-1), (I-2), (I-3), and (I-4) may form salts. Examples of pharmacologically acceptable salts include, when acidic groups exist, metal salts such as lithium salt, sodium salt, potassium salt, magnesium salt, calcium salts, or ammonium salts such as ammonium salt, methylammonium salt, dimethylammonium salt, trimethylammonium salt, dicyclohexylammonium salt, and when basic groups exist, mineral acid salts such as hydrochloride, oxalate, hydrosulfate, nitrate, phosphate, or organic acid salts such as methane sulfonate, benzene sulfonate, para-toluene sulfonate, acetate, propionate, tartrate, fumarate, maleate, malate, oxalate, succinate, citrate, benzoate, mandelate, cinnamate, lactate. Salts may sometimes be formed with amino acids such as glycine. As active ingredients of the medicament of the present invention, pharmacologically acceptable salts may also be suitably used.




























[0316] The compounds or salts thereof represented by the aforementioned general formulas (I), (I-1), (I-2), (I-3), and (I-4) may exist as hydrates or solvates. As active ingredients of the medicament of the present invention, any of the aforementioned substances may be used. Furthermore, the compounds represented by the aforementioned general formulas (I), (I-1), (I-2), (I-3), and (I-4) may sometimes have one or more asymmetric carbons, and may exist as steric isomers such as optically active substance and diastereomer. As active ingredients of the medicament of the present invention, pure forms of stereoisomers, arbitrary mixture of enantiomers or diastereomers, and racemates may be used. When the compounds represented by the general formulas (I), (I-1), (I-2), (I-3), and (I-4) have olefinic double bonds, the configuration may be in either E or Z, and as active ingredients of the medicament of the present invention, geometrical isomer in either of the configurations or a mixture thereof may be used.

[0317] Examples of the compounds as preferred active ingredients of the medicaments of the present invention are






shown below. However, the active ingredients of the medicaments of the present invention are not limited to the compound set out below. In the table, Me represents methyl group, and Et represents ethyl group.




Compound Number		X	E
1			
2			
3			
4			
5			
6			















7			
8			
9			
10			
11			
12			
13			
14			
15			


EP 1 352 650 A1


16			
17			




Compound Number		E
18		
19		
20		
21		
22		

2 3		
2 4		
2 5		
2 6		
2 7		
2 8		
2 9		




















Compound Number		E
3 1		
3 2		
3 3		
3 4		



























Compound Number		E
3 5		
3 6		
3 7		
3 8		
3 9		
4 0		













5	4 1		
10	4 2		
15	4 3		
20	4 4		
25	4 5		
30	4 6		
35	4 7		
40			
45			
50			
55			





























Compound Number		E
4 8		
4 9		
5 0		
5 1		
5 2		
5 3		















5 4		
5 5		
5 6		
5 7		
5 8		
5 9		
6 0		











5 10	6 1		
15	6 2		
20 25	6 3		
30 35	6 4		
40 45	6 5		















6 6		
6 7		
6 8		
6 9		
7 0		
7 1		


5 7 2		
10 7 3		
15 7 4		
20 7 5		
25 7 6		
30 7 7		

7 8		
7 9		
8 0		
8 1		
8 2		
8 3		
8 4		



















8 5		
8 6		
8 7		
8 8		
8 9		
9 0		
9 1		

















<p>9 2</p>		
<p>9 3</p>		
<p>9 4</p>		
<p>9 5</p>		
<p>9 6</p>		

















9 7		
9 8		
9 9		
1 0 0		
1 0 1		
1 0 2		
1 0 3		












Compound Number		E
104		
105		
106		
107		
108		
109		

5 110		
10 111		
15 112		
20 113		
25 114		
30 115		
35 116		
40 117		
45 118		



















1 1 9		
1 2 0		
1 2 1		
1 2 2		
1 2 3		
1 2 4		
1 2 5		
1 2 6		

















5 1 2 7		
10 1 2 8		
15 1 2 9		
20 1 3 0		
25 1 3 1		
30 1 3 2		
35 1 3 3		
40 1 3 4		















1 3 5		
1 3 6		
1 3 7		
1 3 8		

























Compound Number		E
1 3 9		
1 4 0		
1 4 1		
1 4 2		
1 4 3		
1 4 4		


5	1 4 5		
10	1 4 6		
15	1 4 7		
20	1 4 8		
25	1 4 9		
30	1 5 0		
35	1 5 1		
40	1 5 2		
45	1 5 3		
50			
55			

5 154		
10 155		
15 156		
20 157		
25 158		
30 159		
35 160		
40 161		

















5	1 6 2		
10	1 6 3		
15	1 6 4		
20	1 6 5		
25	1 6 6		
30	1 6 7		
35	1 6 8		















169		
170		
171		
172		
173		
174		
175		















5 10 15	1 7 6		
20 25 30	1 7 7		
35 40 45	1 7 8		
50 55	1 7 9		





























Compound Number		E
180		
181		
182		
183		
184		
185		







5	186		
10	187		
15	188		
20	189		
25	190		
30	191		
35	192		
40	193		
45			
50			
55			

194		
195		
196		
197		
198		
199		
200		

201		
202		
203		
204		
205		
206		
207		

208		
209		
210		
211		
212		
213		
214		

2 1 5		
2 1 6		
2 1 7		
2 1 8		
2 1 9		
2 2 0		


2 2 1		
2 2 2		
2 2 3		

[0318] The compounds represented by the general formula (I) can be prepared, for example, by methods shown below.

< Method 1 >

[0319] The compounds represented by the general formula (I), wherein X is -CONH- (the hydrogen atom on the nitrogen may be substituted) can be prepared, for example, by a method described in the reaction scheme 1.

Reaction Scheme 1



wherein A^{101} represents a hydrogen atom or protecting groups of hydroxy group (preferably, an alkyl group such as methyl group and the like; an aralkyl group such as benzyl group and the like; an acetyl group, an alkoxyalkyl group such as methoxymethyl group and the like; a substituted silyl group such as trimethylsilyl group or the like), R^{101} represents a hydrogen atom, a C_1 to C_6 alkyl group or the like, E^{101} represents E or precursor of E in the definition of the general formula (I), G represents a hydroxy group, halogen atoms (preferably, a chlorine atom), a hydrocarbon-oxy group (preferably, an aryl-oxy group which may be substituted by halogen atom), an acyl-oxy group, an imido-oxy group or the like.

(First Step)

[0320] The amide (3) can be prepared by dehydrocondensation of the carboxylic acid derivative (1) and the amine (2). This reaction is carried out at a reaction temperature of from 0°C to 180°C, without solvent or in an aprotic solvent, in the existence of an acid halogenating agent or a dehydrocondensating agent, and in the existence of or nonexistence of a base.

[0321] As the halogenating agent reagent, examples include, for example, thionyl chloride, thionyl bromide, sulfuryl chloride, phosphorus oxychloride, phosphorus trichloride, phosphorus pentachloride or the like, when A¹⁰¹ is hydrogen atom, phosphorus trichloride is preferable, and when A¹⁰¹ is acetyl group or the like, phosphorus oxychloride is preferable. As the dehydrocondensating agent, examples include, for example, N,N'-dicyclohexylcarbodiimide, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, diphenylphosphoryl azide or the like. As the base, examples include inorganic bases such as sodium carbonate, potassium carbonate, sodium hydrocarbonate or the like, or organic bases such as pyridine, triethylamine, N,N'-diethylaniline or the like. As the aprotic solvent, examples include dichloromethane, dichloroethane, chloroform, tetrahydrofuran, 1,4-dioxane, benzene, toluene, monochlorobenzene, o-dichlorobenzene, N,N'-dimethylformamide, N-methylpyrrolidone or the like, when the reaction is carried out in the presence of the acid halogenating agent, particularly, toluene, monochlorobenzene, o-dichlorobenzene are preferable.

[0322] Furthermore, a target compound can be prepared, for example, by a method or similar method described in J. Med. Chem., 1998, 41, 2939, that the acid chloride is prepared and isolated from carboxylic acid in advance, then it is made to react with an amine having E¹⁰¹.

(Second Process)


[0323] When the amide (3) has a protecting group and/or has a favorable substituent for functional group modification, for example, an amino group and its protector or precursor; a carboxy group and its protector or precursor; a hydroxy group and its protector or precursor, the final target compound (4) can be prepared by a reaction for deprotection and/or functional group modification in this step. Various well-known methods can be used for the reaction. For the reaction of deprotection and functional group modification, for example, methods described in "Protective Groups in Organic Syntheses" P.G.M. Wuts, T.Green, Eds., Third version, 1999, Wiley, John & Sons, "Handbook of Reagents for Organic Synthesis" L.A. Paquette, Ed., 4 Volumes, 1999, Wiley, John & Sons can be used, and for reaction of functional group modification, for example, methods described in "Palladium Reagents in Organic Syntheses" R.F. Heck, 1985, Academic Press, "Palladium Reagents and Catalysts: Innovations in Organic Synthesis" J. Tsuji, 1999, Wiley, John & Sons, or the like can be used.

[0324] Aforementioned methods are applicable by combining raw materials properly even in the case where X is other connecting group, for example, -SO₂NH-, -NHCO-, -NHSO₂-, -CONHCH₂-, -CONHCH₂CH₂-; wherein the hydrogen atom on the said connecting group may be substituted.

< Method 2 >

[0325] The compounds represented by the general formula (I), wherein X is -CH₂NH-, for example, can be prepared by a method described in the reaction scheme 2.

Reaction Scheme 2



(wherein each of A and E has the same meaning as that defined in the general formula (I))

[0326] First, the imine derivative of the formula (7) (wherein the definition of R¹~R⁴ and B the same as those in the general formula (I)) can be prepared by dehydrocondensation of the aldehyde (5) and the amine (6). This reaction is carried out at a reaction temperature of from 0°C to 100°C in a solvent, in the existence of or nonexistence of a dehydrating agent. As the dehydrating agent, examples include anhydrous magnesium sulfate, molecular sieves or the like.

As the solvent, examples include nonreactive solvent, and tetrahydrofuran, 1,4-dioxane, methanol, ethanol or the like are preferable.


[0327] Aforementioned methods are applicable by combining raw materials properly even in the case where X is other connecting group, for example, $-\text{CONHN}=\text{CH}-$; the hydrogen atom on the said connecting group may be substituted.

[0328] Next, the target compound (8) can be prepared by reduction of the imine derivative (7). This reaction is carried out at the reaction temperature of from 0°C to 100°C in a solvent, in the existence of a reducing agent. As the reducing agent, examples include sodium borohydride, lithium borohydride or the like. As the solvent, examples include nonreactive solvent, and tetrahydrofuran, 1,4-dioxane, methanol, ethanol or the like are preferable. Moreover, this reaction can be carried out by a method of catalytic hydrogenation also. As the catalyst, examples include palladium carbon, platinum carbon, palladium hydroxide, palladium black or the like. As solvent, examples include nonreactive solvent, and tetrahydrofuran, 1,4-dioxane, methanol, ethanol or the like are preferable. The reaction is carried out at the reaction temperature of from 0°C to 200°C , and the hydrogen pressure is at normal pressure or applied pressure.

< Method 3 >

[0329] The compounds represented by the general formula (I), wherein X is $-\text{CH}=\text{CH}-$ (the hydrogen atom on the said connecting group may be substituted), can be prepared by a method described in the reaction scheme 3.

Reaction Scheme 3




wherein each of A and E has the same meaning as that defined in the general formula (I), W represents O,O'-di-hydrocarbon-phosphono group or triarylphosphonium group.

[0330] The target compound (11) can be prepared by dehydrocondensation of the aldehyde (9) and the phosphorus compound (10). This reaction is carried out in a solvent at a reaction temperature of from 0°C to the boiling point of the solvent, in the existence of a base. As the base, examples include inorganic base such as sodium carbonate, potassium carbonate, sodium hydrogencarbonate or the like, or organic base such as pyridine, triethylamine, N,N'-diethylaniline or the like. Examples include nonreactive solvent, and tetrahydrofuran, 1,4-dioxane, methanol, ethanol, water or the like are preferable.

< Method 4 >

[0331] The compounds represented by the general formula (I), wherein X is $-\text{COCH}=\text{CH}-$ and $-\text{COCH}_2\text{CH}_2-$ (the hydrogen atom on the said connecting group may be substituted), can be prepared by the method, for example, described in the reaction scheme 4.

Reaction Scheme 4



15 [0332] First, the target compound enone (14) can be prepared by dehydrocondensation of the ketone (12) and the aldehyde (13). This reaction is carried out in a solvent at the a reaction temperature of from 0°C to the boiling point of the solvent in the existence of a base. As the base, examples include inorganic base such as sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium hydrogencarbonate or the like, or organic base such as pyridine, triethylamine, N,N'-diethylamine or the like. Examples include nonreactive solvent, and tetrahydrofuran, 1,4-dioxane, methanol, ethanol, water or the like are preferable.

20 [0333] Next, the target compound (15) can be prepared by reduction of the enone (14). This reaction is carried out at the reaction temperature of from 0°C to 100°C in solvent, in the existence of a reducing agent. As the reducing agent, examples include sodium borohydride, lithium borohydride or the like. As the solvent, examples include nonreactive solvent, and tetrahydrofuran, 1,4-dioxane, methanol, ethanol or the like are preferable. Moreover, this reaction is carried out by a method of catalytic hydrogenation also. As the catalyst, examples include palladium carbon, platinum carbon, palladium hydroxide, palladium black or the like. As solvent, examples include nonreactive solvent, and tetrahydrofuran, 1,4-dioxane, methanol, ethanol or the like are preferable. The reaction is carried out at the reaction temperature of from 0°C to 200°C, and the hydrogen pressure is at normal pressure or applied pressure

25 [0334] In the examples of the specification, preparation methods of typical compounds included in the general formula (I) are explained in details. Therefore, those skilled in the art can prepare any compound included in the general formula (I) by referring to the explanations of the aforementioned general preparation methods and of specific preparation methods of the examples, selecting appropriate reaction raw materials, reaction reagents, and reaction conditions, and by adding appropriate modification and alteration of these methods, if necessary.

30 [0335] The compounds represented by the general formula (I) have inhibitory action against NF- κ B activation and inhibitory action against the production and release of inflammatory cytokines, and are useful as active ingredients of pharmaceutical compositions such as NF- κ B inhibitor and inflammatory cytokine release inhibitor. The aforementioned medicament can be suitably used as an expression inhibitor of genes of one or more substances selected from a group comprising tumor necrosis factor (TNF), interleukin-1, interleukin-2, interleukin-6, interleukin-8, granulocyte colony-stimulating factor, interferon β , cell adhesion factor ICAM-1, VCAM-1, and ELAM-1, nitricoxide synthetase, major histocompatibility antigen family class I, major histocompatibility antigen family class II, β 2-microglobulin, immunoglobulin light chain, serum amyloid A, angiotensinogen, complement B, complement C4, c-myc, transcript derived from HIV gene, transcript derived from HTLV gene, transcript derived from simian virus 40 gene, transcript derived from cytomegalovirus gene, and transcript derived from adenovirus gene. Moreover, the medicament of the present invention is useful for preventive and/or therapeutic treatment of diseases caused by NF- κ B activation and inflammatory cytokine overproduction.

45 [0336] More specifically, the medicament of the present invention may be used for preventive and/or therapeutic treatment of the following diseases wherein NF- κ B activation and/or inflammatory cytokine is believed to be involved, for example, autoimmune diseases such as chronic rheumatism, osteoarthritis, systematic lupus erythematosus, systematic scleroderma, polymyositis, Sjogren's syndrome, vasculitis syndrome, antiphospholipid syndrome, Still's disease, Behcet's disease, periarteritis nodosa, ulcerative colitis, Crohn's disease, active chronic hepatitis, glomerulonephritis, and chronic nephritis, chronic pancreatitis, gout, atherosclerosis, multiple sclerosis, arteriosclerosis, endothelial hypertrophy, psoriasis, psoriatic arthritis, contact dermatitis, atopic dermatitis, allergic disease such as pollinosis, asthma, bronchitis, interstitial pneumonia, lung disease involving granuloma, chronic obstructive lung disease, chronic pulmonary thromboembolism, inflammatory colitis, insulin resistance, obesity, diabetes and its complications (nephropathy, retinopathy, neurosis, hyperinsulinemia, arteriosclerosis, hypertension, peripheral vessel obstruction, etc.) diseases involving abnormal vascular proliferation such as hyperlipemia, retinopathy, and pneumonia, Alzheimer's disease, encephalomyelitis, acute hepatitis, chronic hepatitis, drug induced toxic hepatopathy, alcoholic hepatitis, viral hepatitis, icterus, cirrhosis, hepatic insufficiency, atrial myxoma, Caslemann's syndrome, mesangial nephritis, kidney

cancer, lung cancer, liver cancer, breast cancer, uterine cancer, pancreatic cancer, other solid cancer, sarcoma, osteosarcoma, metastatic invasion of cancer, carcination of inflammatory focus, cancerous cachexia, metastasis of cancer, leukemia such as acute myeloblastic leukemia, multiple myeloma, Lennert's lymphoma, malignant lymphoma, development of carcinostatic resistance of cancer, carcination of foci such as viral hepatitis and cirrhosis, carcination from polyp of colon, brain tumor, nervous tumor, endotoxic shock, sepsis, cytomegaloviral pneumonia, cytomegaloviral retinopathy, adenoviral cold, adenoviral pool fever, adenoviral ophthalmia, conjunctivitis, AIDS, uveitis, diseases or complications provoked by infections of other bacteria, viruses, and mycetes, complications after surgery such as generalized inflammatory symptoms, restenosis after percutaneous tubal coronary artery plastic surgery, reperfusion disorders after vascular occlusion opening such as ischemia reperfusion disorders, organ transplantation rejection and reperfusion disorders of heart, liver, kidney or the like, itch, anorexia, malaise, chronic fatigue syndrome or the like. Furthermore, inflammatory cytokine and NF- κ B are involved in differentiation and activation of osteoclast, and consequently, the medicament of the present invention is also useful for preventive and/or therapeutic treatment of metabolic bone diseases or the like such as osteoporosis and osteocarcinomic pain or the like. The medicament may also be used for prevention of deterioration of an organ during organ conservation before transplantation.

[0337] As the active ingredient of the medicament of the present invention, 1 or more kinds of substances selected from the group consisting of the compound represented by the general formula (I) and a pharmacologically acceptable salt thereof, and a hydrate thereof and a solvate thereof may be used. The aforementioned substance, per se, may be administered as the medicament of the present invention, however, preferably, the medicament of the present invention is provided in the form of a pharmaceutical composition comprising the aforementioned substance which is an active ingredient together with one or more pharmacologically acceptable pharmaceutical additives. In the aforementioned pharmaceutical compositions, a ratio of the active ingredient to the pharmaceutical additives is 1 weight % to 90 weight %.

[0338] The pharmaceutical compositions of the present invention may be administered as pharmaceutical compositions for oral administration, for example, granules, subilized granules, powders, hard capsules, soft capsules, syrup, emulsion, suspension, or solution, or may be administered as pharmaceutical compositions for parenteral administration, for example, injections for intravenous administration, intramuscular administration, or subcutaneous administration, drops, suppositories, percutaneous absorbent, transmucosal absorption preparations, nasal drops, ear drops, instillation, and inhalants. Preparations made as pharmaceutical compositions in a form of powder may be dissolved when necessary and used as injections or drip infusions.

[0339] For preparation of pharmaceutical compositions, solid or liquid pharmaceutical additives may be used. Pharmaceutical additives may either be organic or inorganic. When an oral solid preparation is prepared, an excipient is added to the active ingredient, and further binders, disintegrator, lubricant, colorant, corrigent are added, if necessary, preparations in the forms of tablets, coating tablets, granules, powders, capsules and the like may be manufactured by common procedures. Examples of the excipient include lactose, sucrose, saccharose, glucose, corn starch, starch, talc, sorbit, crystal cellulose, dextrin, kaolin, calcium carbonate, and silicon dioxide. Examples of the binder include, for example, polyvinyl alcohol, polyvinyl ether, ethyl cellulose, methyl cellulose, gum Arabic, tragacanth, gelatine, shellac, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, calcium citrate, dextrin, and pectin. Examples of the lubricant include, for example, magnesium stearate, talc, polyethylene glycol, silica, and hydrogenated vegetable oil. As the coloring agent, any material can be used which are approved to be added to ordinary pharmaceuticals. As the corrigent, cocoa powder, menthol, aromatic acid, peppermint oil, d-borneol, cinnamon powder and the like can be used. These tablets and granules may be applied with sugarcoating, gelatine coating, or an appropriate coating, if necessary. Preservatives, antioxidant and the like may be added, if required.

[0340] For liquid preparations for oral administration such as emulsions, syrups, suspensions, and solutions, ordinary used inactive diluents, for example, water or vegetable oil may be used. For these preparations, besides inactive diluents, adjuvants such as wetting agents, suspending aids, sweating agents, flavoring agents, coloring agents or preservatives may be blended. After a liquid preparation is manufactured, the preparation may be filled in capsules made of an absorbable substance such as gelatin. Examples of solvents or suspending agents used for the preparations of parenteral administration such as injections or suppositories include, for example, water, propylene glycol, polyethylene glycol, benzyl alcohol, ethyl oleate, and lecithin. Examples of base materials used for preparation of suppositories include, for example, cacao butter, emulsified cacao butter, lauric fat, and witepsol. Methods for preparation of the aforementioned preparations are not limited, and any method ordinarily used in the art may be used.

[0341] When the composition are prepared in the form of injections, carriers such as, for example, diluents including water, ethanol, macrogol, propyleneglycol, citric acid, acetic acid, phosphoric acid, lactic acid, sodium lactate, sulfuric acid and sodium hydroxide, pH modifiers and buffer solutions including sodium citrate, sodium acetate and sodium phosphate, stabilizers such as sodium pyrosulfite, ethylenediaminetetraacetic acid, thioglycolic acid and thiolactate may be used. For the preparation, a sufficient amount of a salt, glucose, mannitol or glycerin may be blended in the preparation to manufacture an isotonic solution, and an ordinary solubilizer, a soothing agent, or a topical anesthetic may be used.

[0342] When the preparation in the form of an ointment such as a paste, a cream, and a gel is manufactured, an ordinarily used base material, a stabilizer, a wetting agent, and a preservative may be blended, if necessary, and may be prepared by mixing the components by a common method. As the base material, for example, white petrolatum, polyethylene, paraffin, glycerin, cellulose derivatives, polyethylene glycol, silicon, and bentonite may be used. As the preservative, paraoxy methyl benzoate, paraoxy ethyl benzoate, paraoxy propyl benzoate and the like may be used. When the preparation in the form of a patch is manufactured, the aforementioned ointment, cream gel, or paste and the like may be applied by a common method to an ordinary support. As the support, fabric made of cotton, rayon, and synthetic fibers or nonwoven fabric, and a film or a foam sheet such as made of soft vinyl chloride, polyethylene, and polyurethane and the like may be preferably used.

[0343] A dose of the medicament of the present invention is not particularly limited. For oral administration, a dose may generally be 0.01 to 5,000 mg per day for an adult as the weight of the compound of the present invention. It is preferred to increase or decrease the above dose appropriately depending on the age, pathological conditions, and symptoms of a patient. The above dose may be administered once a day or 2 to 3 times a day as divided portions with proper intervals, or intermittent administration for every several days may be acceptable. When the medicament is used as an injection, the dose may be 0.001 to 100 mg per day for an adult as the weight of the compound of the present invention.

Examples

[0344] The present invention will be explained more specifically with reference to the following examples. However the scope of the present invention is not limited to the following examples. And the commercially available compounds, which were purchased and used for the examinations, are contained in these examples. As for such compounds, the suppliers of the reagents and the catalog code numbers are shown.

Example 1: Preparation of N-[[3,5-bis(trifluoromethyl)phenyl]methyl]-5-bromo-2-hydroxybenzamide (Compound No. 1).

[0345] Under argon atmosphere, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (it is abbreviated as WSC · HCl hereafter; 192mg, 1mmol) was added to a mixture of 5-bromosalicylic acid (217mg, 1mmol), 3,5-bis(trifluoromethyl)benzylamine (243mg, 1mmol), 4-dimethylaminopyridine (12mg, 0.1mmol) and tetrahydrofuran (10mL), and the mixture was stirred at room temperature for 1 hour. The reaction mixture was poured into diluted hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous magnesium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel (n-hexane:ethyl acetate=4:1) to give the title compound (244.8mg, 55.4%) as a white solid.

¹H-NMR(DMSO-d₆): δ 4.69(2H, d, J=5.7Hz), 6.93(1H, d, J=8.7Hz), 7.56(1 dd, J=8.7, 2.4Hz), 8.02(1H, d, J=2.4Hz), 8.06(3H, s), 9.41(1H, t, J=5.7Hz), 12.13(1H, s).

Example 2: 5-Bromo-2-hydroxy-N-(2-phenethyl)benzamide (Compound No. 2).

(1) 2-Acetoxy-N-(2-phenethyl)benzamide.

[0346] o-Acetylsalicyloyl chloride (0.20g, 1.00mmol) was dissolved in benzene (8mL). Phenethylamine (0.12g, 1.00mmol) and pyridine (0.3mL) were added, and the mixture was stirred at room temperature for 2 hours. The reaction mixture was poured into diluted hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel (n-hexane:ethyl acetate=2:1→1:1) to give the title compound (155.5mg, 54.9%) as a white crystal.

¹H-NMR(CDCl₃): δ 2.09(3H, s), 2.92(2H, t, J=6.8Hz), 3.71(2H, q, J=6.8Hz), 6.32(1H, brs), 7.07(1H, dd, J=8.4, 1.2Hz), 7.23-7.35(6H, m), 7.44(1H, ddd, J=8.0, 7.6, 1.6Hz), 7.73(1H, dd, J=7.6, 1.6Hz).

(2) 2-Hydroxy-N-(2-phenethyl)benzamide.

[0347] Methanol (5mL) and 2 N sodium hydroxide (0.1mL) were added to 2-acetoxy-N-(2-phenethyl)benzamide (155.5mg), and the mixture was stirred at room temperature for 30 minutes. The reaction mixture was poured into diluted hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was crystallized (dichloromethane/hexane) to give the title compound (106.9mg, 80.7%) as a white solid.

¹H-NMR(DMSO-d₆): δ 2.86(2H, t, J=7.6Hz), 3.52(1H, q, J=7.6Hz), 6.84-6.88(2H, m), 7.18-7.31(5H, m), 7.37(1H, ddd,

EP 1 352 650 A1

J=8.4, 7.2, 1.6Hz), 7.80(1H, dd, J=8.4, 1.6Hz), 8.84(1H, s), 12.51(1H, s).

(3) 5-Bromo-2-hydroxy-N-(2-phenethyl)benzamide.

[0348] Carbon tetrachloride(5mL), iron powder(0.03g) and bromine(25 μ L, 0.48mmol) were added to 2-hydroxy-N-(2-phenethyl)benzamide(79.6mg, 0.33mmol), and the mixture was stirred at room temperature for 1 hour. The reaction mixture was poured into aqueous sodium hydrogen sulfite and extracted with ethyl acetate. After the organic layer was washed with brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel(n-hexane:ethyl acetate=5:1) to give the title compound (62mg, 58.7%) as white powder.

¹H-NMR(DMSO-d₆): δ 2.85(2H, t, J=7.6Hz), 3.52(1H, q, J=7.6Hz), 6.87(1H, d, J=8.8Hz), 7.18-7.31(5H, m), 7.52(1H, dd, J=8.8, 2.4Hz), 8.01(1H, d, J=2.4Hz), 8.90(1H, s), 12.51(1H, s).

Example 3: 5-Bromo-2-hydroxy-N-[5-(morpholinocarbonyl)indan-2-yl]benzamide (Comopund No. 3).

[0349] WSC \cdot HCl(96mg, 0.5mmol) was added to a solution of 5-bromosalicylic acid(109mg, 0.5mmol), 2-amino-5-(morpholino)carbonylindan(refer to Chem. Pharm. Bull., 2000, 48, 131.; 141mg, 0.5mmol) and triethylamine (70 μ L, 0.5mmol) in dichloromethane(5mL), and the mixture was stirred at 40°C for 1.5 hours. After cooling, the reaction mixture was diluted with ethyl acetate, washed with 2 N hydrochloric acid, water and brine one after another, dried over anhydrous magnesium sulfate, concentrated, and the residue was purified by column chromatography on silica gel(dichloromethane:methanol=19:1) to give the title compound(26mg, 11.9%) as a white crystal.

¹H-NMR(CDCl₃): δ 2.66(1H, dd, J=16.2, 7.2Hz), 2.82(1H, dd, J=16.2, 7.2Hz), 3.16-3.25(2H, m), 3.43-3.86(8H, m), 4.79-4.92(1H, m), 6.88(1H, d, J=8.7Hz), 7.14-7.15(3H, m), 7.46(1H, dd, J=8.7, 2.4Hz), 7.74(1H, d, J=7.8Hz), 7.84(1H, d, J=2.4Hz).

Example 4:

3-(4-Chlorophenyl)-1-(2,6-dihydroxyphenyl)-3-(4-hydroxyphenyl)propan-1-one (Comopund No. 4).

[0350] This compound is a commercially available compound.

Supplier: Apin Chemicals.

Catalog code number: N 0100D.

Example 5: 1-(5-Chloro-2-hydroxyphenyl)-3-(4-methoxyphenyl)propan-1-one (Comopund No. 5).

[0351] This compound is a commercially available compound.

Supplier: Specs.

Catalog code number: AI-233/31581024.

Example 6: 1-(2-Hydroxy-4-methoxyphenyl)-3-(2-methoxyphenyl)propen-1-one (Comopund No. 6).

[0352] This compound is a commercially available compound.

Supplier: Maybridge.

Catalog code number: RJC 00106.

Example 7: 3-(3,4-Dihydro-2H-benzo[b][1,4]dioxepin-7-yl)-1-(2-hydroxy-5-methylphenyl)propen-1-one (Comopund No. 7).

[0353] This compound is a commercially available compound.

Supplier: Maybridge.

Catalog code number: BTB 13230.

Example 8: 3-(3,4-Dihydro-2H-benzo[b][1,4]dioxepin-7-yl)-1-[2-hydroxy-4-(methoxymethyl)phenyl]propen-1-one (Comopund No. 8).

[0354] This compound is a commercially available compound.

Supplier: Maybridge.

Catalog code number: BTB 114482.

EP 1 352 650 A1

Example 9: 4-Chloro-2-((4-chlorophenyl)ethen-2-yl)phenol (Comopund No. 9).

[0355] 5-Chlorosalicylaldehyde(313mg, 2mmol) and 4-chlorobenzyltriphenylphosphonium chloride(847mg, 2mmol) were dissolved in N,N-dimethylformamide(20mL). Potassium carbonate(1.382g, 10mmol dissolved in water(10mL was added, and the mixture was refluxed for 5 hours. After cooling, the reaction mixture was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous magnesium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel(n-hexane:ethyl acetate=3:1) to give the title compound(44.6mg, 8.4%) as a light gray solid.

¹H-NMR(CDCl₃): δ 5.04(1H, s), 6.74(1H, d, J=9.0Hz), 7.05(1H, d, J=16.5Hz), 7.10(1H, dd, J=8.4, 2.4Hz), 7.26(1H, d, J=16.5Hz), 7.33(2H, d, J=8.4Hz), 7.45(2H, d, J=8.4Hz), 7.49(1H, d, J=2.4Hz).

Example 10: 5-Bromo-N-(3,5-dichloro)phenyl-2-hydroxybenzenesulfonamide (Comopund No. 10).

[0356] (1) 5-Bromo-N-(3,5-dichloro)phenyl-2-methoxybenzenesulfonamide. 5-Bromo-2-methoxybenzenesulfonyl chloride(857mg, 3mmol) was dissolved in dichloromethane(6mL). A solution of 3,5-dichloroaniline(510mg, 3.15mmol) and pyridine(261mg, 3.3mmol) in dichloromethane(2mL) was added dropwise under ice cooling and argon atmosphere, and the mixture was stirred at room temperature for 6 hours. After the reaction mixture was diluted with dichloromethane, washed with 2 N hydrochloric acid, water and brine one after another, dried over anhydrous magnesium sulfate, the solvent was evaporated under reduced pressure. The obtained residue was crystallized from n-hexane-ethyl acetate to give 5-bromo-2-methoxy-N-(3,5-dichloro)benzenesulfonamide (900mg, 73.0%) as a white crystal.

¹H-NMR(DMSO-d₆): δ 4.03(3H, s), 6.92(1H, d, J=9.0Hz), 7.01(2H, d, J=1.8Hz), 7.07-7.08(1H, m), 7.24(1H, brs), 7.63(1H, dd, J=8.7, 2.4Hz), 7.99(1H, d, J=2.4Hz).

(2) 5-Bromo-N-(3,5-dichloro)phenyl-2-hydroxybenzenesulfonamide.

[0357] A mixture of the white crystal of 5-Bromo-N-(3,5-dichloro)phenyl-2-methoxybenzenesulfonamide(206mg, 0.5mmol), lithium iodide(134mg, 1mmol) and 2,4,6-collidine(5mL) was refluxed for 30 minutes under argon atmosphere. After cooling to room temperature, the reaction mixture was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine one after another, dried over anhydrous magnesium sulfate, the solvent was evaporated under reduced pressure. The obtained residue was crystallized from n-hexane-ethyl acetate to give the title compound(90mg, 45.3%) as a white crystal.

mp 158-159°C.

¹H-NMR(DMSO-d₆, δ): 6.92(1H, d, J=8.7Hz), 7.11(2H, d, J=2.1Hz), 7.21-7.22(1H, m), 7.62(1H, dd, J=8.7, 2.7Hz), 7.80(1H, d, J=2.4Hz), 10.70(1H, br), 11.37(1H, br).

Example 11: 3,5-Bis(trifluoromethyl)-N-(2-hydroxyphenyl)benzamide (Comopund No. 11).

[0358] 2-Aminophenol(120mg, 1.1mmol) was dissolved in dichloromethane(5mL). A solution of 3,5-bis(trifluoromethyl)benzoyl chloride(300mg, 1.1mmol) in dichloromethane(3mL) and pyridine(0.5mL) was added dropwise under ice cooling and argon atmosphere, and the mixture was stirred at room temperature for 1 hour. The reaction mixture was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine one after another, dried over anhydrous magnesium sulfate, the solvent was evaporated under reduced pressure. The obtained residue was dissolved in ethanol(5mL), added dropwise 2 N sodium hydroxide(0.1mL, 0.2mmol), and stirred at room temperature for 30 minutes. The reaction mixture was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine one after another, dried over anhydrous sodium sulfate, the solvent was evaporated under reduced pressure. The obtained residue was purified by column chromatography on silica gel(n-hexane:ethyl acetate=4:1) to give the title compound(288mg, 73.6%) as a light pink crystal.

mp 183°C (dec.).

¹H-NMR(DMSO-d₆, δ): 6.83(1H, td, J=8.0, 1.2Hz), 6.93(1H, dd, J=8.0, 1.2Hz), 7.08(1H, td, J=8.0, 1.6Hz), 7.50(1H, d, J=8.0Hz), 8.35(2H, s), 9.61(1H, s), 10.15(1H, s).

Example 12: N-(5-Chloro-2-hydroxyphenyl)-3,5-dichlorobenzamide (Comopund No. 12).

[0359] 2-Amino-4-chlorophenol(316mg, 2.2mmol) and triethylamine(243mg, 2.4mmol) were dissolved in dichloromethane(5mL). A solution of 3,5-dichlorobenzoyl chloride(419mg, 2mmol) in dichloromethane(2mL) was added dropwise under ice cooling and argon atmosphere, and the mixture was stirred at room temperature for 15 hours. After the reaction mixture was diluted with ethyl acetate, washed with water and brine one after another, dried over anhydrous

magnesium sulfate, the solvent was evaporated under reduced pressure. The obtained residue was purified by column chromatography on silica gel(n-hexane:ethyl acetate=3:1) to give a light brown solid. The solid was suspended and washed with n-hexane-ethyl acetate under heating at reflux to give the title compound(205mg, 32.4%) as a white crystal. mp 251-252°C.

¹H-NMR(DMSO-d₆): δ 6.93(1H, d, J=9.0Hz), 7.11(1H, dd, J=8.7, 2.7Hz), 7.67(2H, d, J=2.7Hz), 7.86-7.87(1H, m), 7.97(1H, d, J=1.8Hz), 9.85(1H, s), 10.03(1H, s).

Example 13: N-(5-Chloro-2-hydroxyphenyl)-3,5-dichlorobenzenesulfonamide (Comopund No. 13).

[0360] 2-Amino-4-chlorophenol(287mg, 2mmol) and 3,5-dichlorobenzenesulfonyl chloride(540mg, 2.2mmol) were dissolved in dichloromethane(4mL). Pyridine(1mL) was added dropwise under ice cooling and argon atmosphere, and the mixture was stirred at room temperature for 1 hour. The reaction mixture was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine one after another, dried over anhydrous magnesium sulfate, the solvent was evaporated under reduced pressure. The obtained residue was purified by column chromatography on silica gel(n-hexane:ethyl acetate=3:1→ 1:1) to give a reddish brown solid. The solid was crystallized from n-hexane-ethyl acetate to give the title compound(445mg, 63.1%) as a slight dark brown crystal. mp 190-191°C.

¹H-NMR(DMSO-d₆): δ 7.68(1H, d, J=9.0Hz), 7.08(1H, dd, J=8.7, 2.7Hz), 7.17(1H, d, J=2.4Hz), 7.70(2H, d, J=1.8Hz), 7.95-7.96(1H, m), 10.00(1H, s), 10.06(1H, s).

Example 14: N-[(5-Bromo-2-hydroxyphenyl)methyl]-3,5-dichloroaniline (Comopund No. 14).

(1) 4-Bromo-2-[(3,5-diphenylimino)methyl]phenol.

[0361] A mixture of 5-bromosalicylaldehyde(1.01g, 5mmol), 3,5-dichloroaniline(810mg, 5mmol) and ethanol(25mL) was refluxed for 1 hour under argon atmosphere. After the reaction mixture was cooled to room temperature, the separated crystal was filtered to give 3,5-dichloro-N-(5-bromo-2-hydroxybenzylidene)aniline(1.52g, 88.2%) as an orange crystal.

mp 161-163°C. ¹H-NMR(CDCl₃, δ): 6.94(1H, d, J=9.0Hz), 7.16(2H, d, J=1.8Hz), 7.30-7.31(1H, m), 7.47-7.53(2H, m), 8.51(1H, s).

(2) N-[(5-Bromo-2-hydroxyphenyl)methyl]-3,5-dichloroaniline

[0362] 3,5-Dichloro-N-(5-bromo-2-hydroxybenzylidene)aniline(1.04g, 3mmol) was dissolved in tetrahydrofuran(12mL) and ethanol(6mL). Sodium borohydride(113mg, 3mmol) was added under ice cooling and argon atmosphere, and the mixture was stirred at room temperature for 12 hours. Acetone(10mL) was added to the reaction mixture, and the residue obtained by concentration under reduced pressure was added water and extracted with dichloromethane. After the dichloromethane layer was washed with water and brine one after another, dried over anhydrous magnesium sulfate, the solvent was evaporated under reduced pressure. The obtained residue was purified by column chromatography on silica gel(n-hexane:ethyl acetate=4:1) to give a light yellow viscous material. This was crystallized by n-hexane to give the title compound(971mg, 93.3%) as a white crystal.

mp 125-126°C. ¹H-NMR(CDCl₃, δ): δ 4.31(2H, s), 6.64(2H, d, J=1.8Hz), 6.74-6.77(1H, m), 6.84-6.85(1H, m), 7.30-7.34(2H, m).

Example 15: 5-Chloro-2-hydroxybenzoic acid (2,4-dihydroxybenzylidene)hydrazide (Comopund No. 15).

[0363] This compound is a commercially available compound.

Supplier: Sigma-Aldrich.

Catalog code number: S3203-5.

Example 16: N-[3,5-Bis(trifluoromethyl)phenyl]-5-chloro-2-hydroxy-N-methylbenzamide (Comopund No. 16).

[0364] A mixture of 5-chlorosalicylic acid(173mg, 1mmol), 3,5-bis(trifluoromethyl)-N-methylaniline(243mg, 1mmol), phosphorus trichloride(44μ 1, 0.5mmol) and monochlorobenzene(5mL) was refluxed for 3 hours under argon atmosphere. After the reaction mixture was cooled to room temperature, n-hexane(50mL) was added, and the separated crude crystal was filtered and dissolved in ethyl acetate(50mL). After the ethyl acetate solution was washed with water and brine one after another, dried over anhydrous sodium sulfate, the solvent was evaporated under reduced pressure. The obtained residue was purified by column chromatography on silica gel(n-hexane:ethyl acetate=2:1) to give the-

EP 1 352 650 A1

title compound (75mg, 18.9%) as a white crystal.

¹H-NMR(CDCl₃): δ 6.59(1H, d, J=2.4Hz), 6.94(1H, d, J=9.0Hz), 7.21(1H, dd, J=9.0, 2.7Hz), 7.58(2H, s), 7.80(1H, s), 10.00(1H, brs).

Example 17: 1-(5-Bromo-2-hydroxy)benzoyl-7-(trifluoromethyl)-1,2,3,4-tetrahydroquinoline (Comopund No. 17).

[0365] Using 5-bromosalicylic acid and 7-(trifluoromethyl)-1,2,3,4-tetrahydroquinoline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 42.0%.

¹H-NMR(CDCl₃): δ 2.08(2H, m), 2.92(2H, t, J=6.6Hz), 3.95(2H, t, J=6.6Hz), 6.91-6.94(2H, m), 7.14(1H, s), 7.32-7.35(2H, m), 7.40(1H, dd, J=8.7, 2.4Hz), 10.06(1H, s).

Example 18: N-(3,5-Dichlorophenyl)-2-hydroxy-1-naphthamide (Comopund No. 18).

[0366] Using 2-hydroxynaphthalene-1-carboxylic acid and 3,5-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 51.2%.

mp 246-248°C.

¹H-NMR(DMSO-d₆): δ 7.26(1H, d, J=9.3Hz), 7.31-7.37(2H, m), 7.44-7.50(1H, m), 7.65-7.68(1H, m), 7.85-7.90(4H, m), 10.23(1H, s), 10.74(1H, s).

Example 19: N-(3,5-Dichlorophenyl)-3-hydroxy-2-naphthamide (Comopund No. 19).

[0367] Using 3-hydroxynaphthalene-2-carboxylic acid and 3,5-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

mp 254-255°C.

¹H-NMR(DMSO-d₆): δ 7.34-7.39(3H, m), 7.49-7.54(1H, m), 7.76-7.79(1H, m), 7.89(2H, d, J=1.8Hz), 7.92(1H, m), 8.39(1H, s), 10.75(1H, s), 11.01(1H, s).

Example 20: N-(3,5-Dimethoxyphenyl)-3-hydroxy-2-naphthamide (Comopund No. 20).

[0368] This compound is a commercially available compound.

Supplier: Sigma-Aldrich.

Catalog code number: S01361-8.

Example 21: N-[3,5-Bis(trifluoromethyl)phenyl]-1-hydroxy-2-naphthamide (Comopund No. 21).

[0369] Using 1-hydroxynaphthalene-2-carboxylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 65.5%.

¹H-NMR(DMSO-d₆): δ 7.51(1H, d, J=9.0Hz), 7.60(1H, td, J=7.8, 0.9Hz), 7.70(1H, td, J=7.8, 0.9Hz), 7.89(1H, s), 7.93(1H, d, J=8.4Hz), 8.09(1H, d, J=9.0Hz), 8.33(1H, d, J=8.7Hz), 8.51(2H, s), 10.92(1H, s), 13.36(1H, s).

Example 22: (((1-Hydroxynaphthalen-2-yl)carbonyl)amino)benzenesulfonyl fluoride (Comopund No. 22).

[0370] This compound is a commercially available compound.

Supplier: Sigma-Aldrich.

Catalog code number: S58026-0.

Example 23: 4-([4-(2,5-Dichlorophenyl)azo-1-hydroxynaphthalen-2-yl]carbonyl)amino)benzenesulfonyl fluoride (Comopund No. 23).

[0371] This compound is a commercially available compound.

Supplier: Sigma-Aldrich.

Catalog code number: S63263-5.

EP 1 352 650 A1

Example 24: N-[3,5-Bis(trifluoromethyl)phenyl]-5-chloro-2-hydroxypyridine-3-carboxamide (Comopund No. 24).

[0372] 5-Chloro-2-hydroxynicotinic acid(174mg, 1mmol), 3,5-bis(trifluoromethyl)aniline(275mg, 1.2mmol) and pyridine(316mg, 4mmol) were dissolved in tetrahydrofuran(20mL) and dichloromethane(10mL). Phosphorus oxychloride (0.112ml, 1.2mmol) was added, and the mixture was stirred at room temperature for 2 hours. The reaction mixture was poured into ethyl acetate(100mL) and 0.2 N hydrochloric acid(100mL), filtered through celite after stirring for 30 minutes, and the water layer of the filtrate was extracted with ethyl acetate. After the combined ethyl acetate layer was washed with water and brine one after another, dried over anhydrous magnesium sulfate, the solvent was evaporated under reduced pressure. The residue was purified by column chromatography on silica gel(n-hexane:ethyl acetate=2:1→ 1:1) to give a light yellow solid. This was suspended and washed with ethanol under heating at reflux to give the title compound(183mg, 47.6%) as a white crystal.

mp >270°C.

¹H-NMR(DMSO-d₆): δ 7.83(1H, s), 8.15(1H, d, J=3.3Hz), 8.36(1H, d, J=3.0Hz), 8.40(2H, s), 12.43(1H, s).

Example 25: N-[2-Chloro-5-(trifluoromethyl)phenyl]-5-chloro-2-hydroxynicotinamide (Comopund No. 25).

[0373] Using 5-chloro-2-hydroxynicotinic acid and 2-chloro-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 24 gave the title compound.

Yield: 42.9%.

¹H-NMR(DMSO-d₆): δ 7.52(1H, dd, J=8.4, 2.1Hz), 7.81(1H, d, J=8.4Hz), 8.16(1H, s), 8.39(1H, d, J=2.7Hz), 8.96(1H, d, J=2.1Hz), 12.76(1H, s), 13.23(1H, s).

Example 26: N-(3,5-Bis[(1,1-dimethyl)ethyl]phenyl)-5-chloro-2-hydroxynicotinamide (Comopund No. 26).

[0374] Using 5-chloro-2-hydroxynicotinic acid and 3,5-bis[(1,1-dimethyl)ethyl]aniline as the raw materials, the same operation as the example 24 gave the title compound.

Yield: 59.1%.

¹H-NMR(DMSO-d₆): δ 1.29(18H, s), 7.18(1H, t, J=1.8Hz), 7.52(2H, d, J=1.8Hz), 8.07(1H, d, J=2.4Hz), 8.35(1H, d, J=3.3Hz), 11.92(1H, s), 13.10(1H, s).

Example 27: N-[3,5-Bis(trifluoromethyl)phenyl]-3-hydroxypyridine-2-carboxamide (Comopund No. 27).

[0375] Using 3-hydroxypyridine-2-carboxylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 24 gave the title compound.

Yield: 45.0%.

¹H-NMR(CDCl₃): δ 7.40(1H, dd, J=8.4, 1.8Hz), 7.46(1H, dd, J=8.4, 4.2Hz), 7.68(1H, s), 8.16(1H, dd, J=4.2, 1.2Hz), 8.25(2H, s), 10.24(1H, s), 11.42(1H, s).

Example 28: N-[3,5-Bis(trifluoromethyl)phenyl]-6-chloro-2-hydroxyindole-3-carboxamide (Comopund No. 28).

[0376] Under argon atmosphere, 3,5-bis(trifluoromethyl)isocyanate(255mg, 1.0mmol) was dissolved in tetrahydrofuran(5mL). A solution of 6-chloro-oxindole(184mg, 1.1mmol) in tetrahydrofuran(5ml) and triethylamine(0.3mL) were added, and the mixture was stirred at room temperature for 4 hours. The reaction mixture was poured into diluted hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous magnesium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel(n-hexane:ethyl acetate=4:1) to give the title compound(172.2mg, 40.7%) as a pink solid.

¹H-NMR(DMSO-d₆): δ 3.97(2H, s), 7.29(1H, dd, J=8.1, 2.1Hz), 7.41(1H, d, J=8.1Hz), 7.88(1H, s), 8.04(1H, d, J=2.1Hz), 8.38(2H, s), 10.93(1H, s).

Example 29: N-[3,5-Bis(trifluoromethyl)phenyl]-3-hydroxyquinoxaline-2-carboxamide (Comopund No. 29).

[0377] Using 3-hydroxyquinoxaline-2-carboxylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 2.7%.

¹H-NMR(DMSO-d₆): δ 7.40-7.45(2H, m), 7.69(1H, td, J=8.4, 1.5Hz), 7.90-7.93(2H, m), 8.41(2H, s), 11.64(1H, s), 13.02(1H, s).

EP 1 352 650 A1

Example 30: N-(4-Chlorophenyl)-2-hydroxy-9H-carbazole-3-carboxamide (Comopund No. 30).

[0378] This compound is a commercially available compound.

Supplier: Sigma-Aldrich.

Catalog code number: S83846-2.

Example 31: 2-Hydroxy-N-(1-naphthyl)benzamide (Comopund No. 31).

[0379] This compound is a commercially available compound.

Supplier: Maybridge.

Catalog code number: RDR 01818.

Example 32: 5-Chloro-2-hydroxy-N-(1-naphthyl)benzamide (Comopund No. 32).

[0380] Using 5-chlorosalicylic acid and 1-naphthylamine as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 65.0%.

¹H-NMR(DMSO-d₆): δ 7.09(1H, d, J=8.7Hz), 7.51-7.61(4H, m), 7.85(1H, d, J=8.4Hz), 7.96(1H, d, J=7.5Hz), 7.99-8.05(2H, m), 8.13(1 d, J=2.7Hz), 10.88(1H, s), 12.31(1H, s).

Example 33: 5-Chloro-2-hydroxy-N-(4-methoxynaphthalen-2-yl)benzamide (Comopund No. 33).

[0381] Using 5-chlorosalicylic acid and 4-methoxy-1-naphthylamine as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 84.3%.

¹H-NMR(DMSO-d₆): δ 3.99(3H, s), 7.05(1H, d, J=9.0Hz), 7.30(1H, d, J=1.5Hz), 7.39-7.45(1H, m), 7.48-7.54(2H, m), 7.83(1H, d, J=7.8Hz), 8.00(1H, s), 8.02(1H, d, J=2.4Hz), 8.09(1H, d, J=7.8Hz), 10.54(1H, s), 11.88(1H, s).

Example 34: 2-Acetoxy-5-Chloro-N-(4-methoxynaphthalen-2-yl)benzamide (Comopund No. 34).

[0382] Using 2-acetoxy-5-chlorobenzoic acid and 4-methoxy-1-naphthylamine as the raw materials, the same operation as the example 24 gave the title compound.

(2-Acetoxy-5-chlorobenzoic acid: refer to Eur. J. Med. Chem., 1996, 31, 861.)

Yield: 39.9% red solid.

¹H-NMR(DMSO-d₆): δ 2.23(3H, s), 3.96(3H, s), 7.23(1H, d, J=1.2Hz), 7.34(1H, d, J=8.7Hz), 7.40(1H, dt, J=8.1, 1.2Hz), 7.50(1H, dt, J=8.1, 1.5Hz), 7.67(1H, dd, J=8.7, 2.7Hz), 7.81(1H, d, J=8.7Hz), 8.72(1H, d, J=3.0Hz), 8.02(1H, s), 8.08(1H, d, J=8.7Hz), 10.58(1H, s).

Example 35: 2-(5-Chloro-2-hydroxybenzoyl)amino-4,5,6,7-tetrahydrobenzo[b]thiophene-3-carboxylic acid ethyl ester (Comopund No. 35).

[0383] Using 5-chlorosalicylic acid and 2-amino-4,5,6,7-tetrahydrobenzo[b]thiophene-3-carboxylic acid ethyl ester as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 49.6%.

¹H-NMR(DMSO-d₆): δ 1.32(3H, t, J=7.2Hz), 1.74(4H, br), 2.63(2H, br), 2.75(2H, br), 4.30(2H, q, J=7.2Hz), 7.05(1H, d, J=9.0Hz), 7.50(1H, dd, J=8.7, 3.0Hz), 7.92(1H, d, J=3.0Hz), 12.23(1H, s), 13.07(1H, s).

Example 36: 5-Bromo-2-hydroxy-N-(5-phenylpyrazol-3-yl)benzamide (Comopund No. 36).

[0384] Using 5-bromosalicylic acid and 3-amino-5-phenylpyrazole as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 9.2%.

¹H-NMR(DMSO-d₆): δ 6.98(1H, d, J=8.8Hz), 7.01(1H, s), 7.35(1H, t, J=7.6Hz), 7.46(2H, t, J=7.6Hz), 7.58(1H, dd, J=8.8, 2.8Hz), 7.74-7.76(2H, m), 8.19(1H, s), 10.86(1H, s), 12.09(1H, s), 13.00(1H, brs).

EP 1 352 650 A1

Example 37: 5-Bromo-N-(4,5-diethyloxazol-2-yl)-2-hydroxybenzamide (Comopund No. 37).

(1) 2-Amino-4,5-diethyloxazole.

[0385] Propionin(1.03g, 8.87mmol) was dissolved in ethanol(15mL). Cyanamide(0.75g, 17.7mmol) and sodium ethoxide(1.21g, 17.7mmol) were added, and the mixture was stirred at room temperature for 3.5 hours. The reaction mixture was poured into water and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel(dichloromethane:methanol=9:1) to give the title compound(369.2mg, 29.7%) as a yellow amorphous.

¹H-NMR(DMSO-d₆): δ 1.04(3H, t, J=7.5Hz), 1.06(3H, t, J=7.5Hz), 2.20(2H, q, J=7.5Hz), 2.43(2H, q, J=7.5Hz), 6.15(2H, s).

(2) 2-Acetoxy-5-bromo-N-(4,5-diethyloxazol-2-yl)benzamide.

[0386] Using 2-acetoxy-5-bromobenzoic acid and 2-amino-4,5-diethyloxazole as the raw materials, the same operation as the example 24 gave the title compound.

(2-Acetoxy-5-bromobenzoic acid: refer to Eur. J. Med. Chem., 1996, 31, 861.)

Yield: 22.0%.

¹H-NMR(CDCl₃): δ 1.22(3H, t, J=7.5Hz), 1.23(3H, t, J=7.5Hz), 2.48(2H, q, J=7.5Hz), 2.57(2H, q, J=7.5Hz), 6.96(1H, d, J=8.7Hz), 7.58(1H, dd, J=8.7, 2.7Hz), 8.32(1H, s), 11.40(1H, br).

(3) 5-Bromo-N-(4,5-diethyloxazol-2-yl)-2-hydroxybenzamide.

[0387] Using 2-acetoxy-5-bromo-N-(4,5-diethyloxazol-2-yl)benzamide as the raw material, the same operation as the example 2(2) gave the title compound.

Yield: 70.2%.

¹H-NMR(CDCl₃): δ 1.25(3H, t, J=7.5Hz), 1.26(3H, t, J=7.5Hz), 2.52(2H, q, J=7.5Hz), 2.60(2H, q, J=7.5Hz), 6.84(1H, d, J=8.7Hz), 7.43(1H, dd, J=8.7, 3.0Hz), 8.17(1H, d, J=3.0Hz), 11.35(1H, br), 12.83(1H, br).

Example 38: 5-Bromo-N-(4,5-diphenyloxazol-2-yl)-2-hydroxybenzamide (Comopund No. 38).

[0388] Using 5-bromosalicylic acid and 2-amino-4,5-diphenyloxazole as the raw materials, the same operation as the example 16 gave the title compound.

(2-Amino-4,5-diphenyloxazole: refer to Zh. Org. Khim., 1980, 16, 2185.)

Yield: 32.6%.

mp 188-189°C.

¹H-NMR(DMSO-d₆, δ): 6.98(1 d, J=8.7Hz), 7.40-7.49(6H, m), 7.53-7.56(2H, m), 7.59-7.63(3H, m), 8.01(1H, d, J=2.4Hz), 11.80(2H, brs).

Example 39: 5-Bromo-N-[4,5-bis(furan-2-yl)oxazol-2-yl]-2-hydroxybenzamide (Comopund No. 39).

(1) 2-Amino-4,5-bis(furan-2-yl)oxazole.

[0389] Furoin(0.50g, 2.60mmol) was dissolved in ethanol(15mL). Cyanamide(218.8mg, 5.20mmol) and sodium ethoxide(530.8mg, 7.80mmol) were added, and the mixture was stirred at room temperature for 2 hours. The reaction mixture was poured into water and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel(hexane:ethyl acetate=1:1→1:2) to give the title compound (175.0mg, 31.1%) as a dark brown crystal.

¹H-NMR(DMSO-d₆): δ 6.59(1H, dd, J=3.3, 2.1Hz), 6.62(1H, dd, J=3.3, 2.1Hz), 6.73(1H, dd, J=3.3, 0.6Hz), 6.80(1H, dd, J=3.3, 0.9Hz), 7.05(2H, s), 7.75-7.76(2H, m).

(2) 5-Bromo-N-[4,5-bis(furan-2-yl)oxazol-2-yl]-2-hydroxybenzamide.

[0390] Using 5-bromosalicylic acid and 2-amino-4,5-bis(furan-2-yl)oxazole as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 12.9%.

EP 1 352 650 A1

¹H-NMR(DMSO-d₆): δ 6.65(1H, dd, J=3.6, 1.8Hz), 6.68(1H, dd, J=3.6, 1.8Hz), 6.75(1H, d, J=8.7Hz), 6.92(1H, dd, J=3.6, 0.9Hz), 6.93(1H, d, J=3.3Hz), 7.37(1H, dd, J=8.7, 2.7Hz), 7.80(1H, dd, J=1.8, 0.9Hz), 7.84(1H, dd, J=1.8, 0.9Hz), 7.92(1H, d, J=3.0Hz), 14.88(2H, br).

5 Example 40: 2-Hydroxy-N-5-[(trifluoromethyl)-1,3,4-thiadiazol-2-yl]benzamide (Comopund No. 40).

(1) 2-Acetoxy-N-5-[(trifluoromethyl)-1,3,4-thiadiazol-2-yl]benzamide.

10 [0391] Using o-acetylsalicyloyl chloride and 2-amino-5-(trifluoromethyl)-1,3,4-thiadiazole as the raw materials, the same operation as the example 2(1) gave the title compound.

Yield: 51.1%.

¹H-NMR(DMSO-d₆): δ 2.23(3H, s), 7.32(1H, dd, J=8.0, 1.2Hz), 7.45(1H, td, J=7.6, 1.2Hz), 7.69(1H, td, J=8.0, 2.0Hz), 7.87(1H, dd, J=8.0, 2.0Hz), 13.75(1H, brs).

15 (2) 2-Hydroxy-N-5-[(trifluoromethyl)-1,3,4-thiadiazol-2-yl]benzamide.

[0392] Using 2-acetoxy-N-5-[(trifluoromethyl)-1,3,4-thiadiazol-2-yl]benzamide as the raw material, the same operation as the example 2(2) gave the title compound.

Yield: 92.9%.

20 ¹H-NMR(DMSO-d₆): δ 7.00(1H, td, J=8.0, 0.8Hz), 7.06(1H, d, J=8.4Hz), 7.51(1H, ddd, J=8.4, 7.6, 2.0Hz), 7.92(1H, dd, J=8.0, 1.6Hz), 12.16(1H, br).

Example 41: 5-Bromo-2-hydroxy-N-[5-(trifluoromethyl)-1,3,4-thiadiazol-2-yl]benzamide (Comopund No. 41).

25 [0393] Using 5-bromosalicylic acid and 2-amino-5-(trifluoromethyl)-1,3,4-thiadiazole as the raw materials, the same operation as the example 16 gave the title compound. Yield: 80.2%.

¹H-NMR(DMSO-d₆): δ 7.01(1H, d, J=9.0Hz) 7.63(1H, dd, J=8.7, 2.7Hz), 7.97(1H, d, J=2.4Hz).

Example 42: 5-Chloro-N-(2-chloropyridin-4-yl)-2-hydroxybenzamide (Comopund No. 42).

30

[0394] Using 5-chlorosalicylic acid and 4-amino-2-chloropyridine as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 12.2%.

35 ¹H-NMR(DMSO-d₆): δ 7.04(1H, d, J=9.0Hz), 7.49(1H, dd, J=9.0, 3.0Hz), 7.54(1H, d, J=8.4Hz), 7.88(1H, d, J=2.7Hz), 8.21(1H, dd, J=8.7, 2.7Hz), 8.74(1H, d, J=2.7Hz), 10.62(1H, s), 11.57(1H, s).

Example 43: 5-Chloro-N-(6-chloro-4-methoxypyrimidin-2-yl)-2-hydroxybenzamide (Comopund No. 43).

40 [0395] Using 5-chlorosalicylic acid and 2-amino-6-chloro-4-methoxypyrimidine as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 2.2%, white solid.

¹H-NMR(DMSO-d₆): δ 3.86(3H, s), 6.85(1H, s), 7.01(1H, d, J=9.0Hz) 7.47(1H, dd, J=9.0, 3.0Hz), 7.81(1H, d, J=3.0Hz), 11.08(1H s), 11.65(1H, s).

45 Example 44: 2-Aceoxy-5-chloro-N-(indol-2-yl)benzamide (Comopund No. 44).

[0396] Using 2-acetoxy-5-chlorosalicylic acid and 2-aminoindole as the raw materials, the same operation as the example 24 gave the title compound.

Yield: 13.3%.

50 ¹H-NMR(DMSO-d₆): δ 2.20(3H, s), 6.41(1H, t, J=2.1Hz), 7.27-7.36(4H, m), 7.63(1H, dd, J=8.7, 2.7Hz), 7.74(1H, d, J=2.7Hz), 7.93(1H, s), 10.21(1H, s), 11.04(1H, s).

Example 45: 7-[(2-Acetoxybenzoyl)aminolindole-3-carboxylic acid ethyl ester (Comopund No. 45).

55 [0397] This compound is a commercially available compound.

Supplier: Peakdale.

Catalog code number: PFC-0448.

EP 1 352 650 A1

Example 46: 5-Chloro-2-hydroxy-N-(quinolin-3-yl)benzamide (Comopund No. 46).

[0398] Using 5-chlorosalicylic acid and 2-aminoquinoline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 4.3%.

¹H-NMR(DMSO-d₆): δ 7.07(1H, d, J=8.7Hz), 7.51(1H, dd, J=9.0, 3.0Hz), 7.61(1H, dt, J=7.8, 1.2Hz), 7.70(1H, dt, J=7.8, 1.5Hz), 7.98(2H, d, J=3.0Hz), 8.01(1H, s), 8.82(1H, d, J=2.4Hz), 10.80(1H, s), 11.74(1H, s).

Example 47: N-(9-Ethylcarbazol-3-yl)-5-chloro-2-hydroxybenzamide (Comopund No. 47).

[0399] Using 5-chlorosalicylic acid and 3-amino-9-ethylcarbazole as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 64.6%.

¹H-NMR(DMSO-d₆): δ 1.33(3H, t, J=7.0Hz), 4.46(2H, q, J=7.0Hz), 7.04(1H, d, J=9.0Hz), 7.21(1H, t, J=7.3Hz), 7.45-7.52(2H, m), 7.64-7.65(2H, m), 7.70(1H, d, J=8.4, 1.9Hz), 8.11-8.15(2H, m), 8.49(1H, d, J=1.9Hz), 10.55(1H, s), 12.22(1H, s).

Example 48: 2-Acetoxy-[3,5-bis(trifluoromethyl)phenyl]benzamide (Comopund No. 95).

[0400] Using o-acetylsalicyloyl chloride and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 2(1) gave the title compound.

Yield: 84.2%.

¹H-NMR(DMSO-d₆): δ 2.36(3H, s), 7.19(1H, dd, J=8.0, 1.2Hz), 7.39(1H, td, J=7.6, 1.2Hz), 7.57(1H, ddd, J=8.0, 7.6, 1.6Hz), 7.65(1H, s), 7.83(1H, dd, J=8.0, 1.6Hz), 8.11(2H, s), 8.31(1H, s).

Example 49: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 48).

[0401] Using 2-acetoxy-[3,5-bis(trifluoromethyl)phenyl]benzamide as the raw material, the same operation as the example 2(2) gave the title compound.

Yield: 45.1%.

¹H-NMR(DMSO-d₆): δ 6.96-7.02(2H, m), 7.45(1H, ddd, J=9.0, 7.2, 1.6Hz), 7.81(1H, s), 7.87(1H, dd, J=8.0, 1.6Hz), 8.46(2H, s), 10.80(1H, s), 11.26(1H, s).

Example 50: N-[3,5-Bis(trifluoromethyl)phenyl]-5-fluoro-2-hydroxybenzamide (Comopund No. 49).

[0402] Using 5-fluorosalicic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 58.7%.

¹H-NMR(DMSO-d₆): δ 7.04(1H, ddd, J=9.0, 4.5, 1.2Hz), 7.30-7.37(1H, m), 7.66(1H, ddd, J=9.0, 3.3, 1.2Hz), 7.84(1H, s), 8.46(2H, s), 10.85(1H, s), 11.21(1H, brs).

Example 51: N-[3,5-Bis(trifluoromethyl)phenyl]-5-chloro-2-hydroxybenzamide (Comopund No. 50).

[0403] Using 5-chlorosalicylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 85.5%.

¹H-NMR(DMSO-d₆): δ 7.05(1H, d, J=8.7Hz), 7.49(1H, dd, J=8.7, 2.7Hz), 7.85(1H, s), 7.87(1H, d, J=2.7Hz), 8.45(2H, s), 10.85(1H, s), 11.39(1H, s).

Example 52: N-[3,5-Bis(trifluoromethyl)phenyl]-5-bromo-2-hydroxybenzamide (Comopund No. 51).

[0404] Using 5-bromosalicylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 88.5%.

¹H-NMR(DMSO-d₆): δ 6.98(1H, d, J=8.8Hz), 7.59(1H, dd, J=8.8, 2.8Hz), 7.83(1H, s), 7.98(1H, d, J=2.8Hz), 8.43(2H, s), 10.82(1H, s), 11.37(1H, s).

EP 1 352 650 A1

Example 53: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-iodobenzamide (Comopund No. 52).

[0405] Using 5-iodosalicylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 62.2%.

¹H-NMR(DMSO-d₆): δ 6.86(1H, d, J=8.4Hz), 7.74(1H, dd, J=8.7, 2.4Hz), 7.84(1H, s), 8.13(1H, d, J=2.1Hz), 8.84(2H, s), 10.82(1H, s), 11.41(1H, s).

Example 54: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-nitrobenzamide (Comopund No. 53).

[0406] Using 5-nitrosalicylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 57.2%.

¹H-NMR(DMSO-d₆): δ 7.18(1H, d, J=9.0Hz), 7.86(1H, s), 8.31(1H, dd, J=9.0, 3.0Hz), 8.45(2H, s), 8.70(1H, d, J=3.0Hz), 11.12(1H, s).

Example 55: N-[3,5-Bis(trifluoromethyl)phenyl]-5-cyano-2-hydroxybenzamide (Comopund No. 54).

[0407] Using 5-cyanosalicylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 16.6%.

¹H-NMR(DMSO-d₆): δ 7.15(1H, d, J=8.7Hz), 7.85(1H, s), 7.86(1H, dd, J=8.7, 2.1Hz), 8.22(1H, d, J=2.4Hz), 8.43(2H, s), 10.93(1H, s), 12.00(1H, brs).

Example 56: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-methylbenzamide (Comopund No. 55).

[0408] Using 5-methylsalicylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 54.9%.

¹H-NMR(DMSO-d₆): δ 6.92(1H, d, J=8.7Hz), 7.28(1H, dd, J=8.7, 1.8Hz), 7.71(1H, d, J=1.8Hz), 7.82(1H, s), 8.47(2H, s), 10.80(1H, s), 11.14(1H, s).

Example 57: N-[3,5-Bis(trifluoromethyl)phenyl]-5-(1,1-dimethyl)ethyl-2-hydroxybenzamide (Comopund No. 56).

[0409] Using 5-[(1,1-dimethyl)ethyl]salicylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 53.8%.

¹H-NMR(DMSO-d₆): δ 1.30(9H, s), 6.96(1H, d, J=8.7Hz), 7.50(1H, dd, J=8.7, 2.4Hz), 7.82(1H, d, J=2.4Hz), 7.83(1H, s), 8.46(2H, s), 10.80(1H, s), 11.12(1H, s).

Example 58: 5-Acetyl-N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 78).

(1) 5-Acetyl-2-benzyloxybenzoic acid methyl ester.

[0410] A mixture of 5-acetylsalicylic acid methyl ester(13.59g, 70mmol), benzyl bromide(17.96g, 105mmol), potassium carbonate(19.35g, 140mmol) and methylethylketone(350mL) was refluxed for 8 hours. After cooling, the solvent was evaporated under reduced pressure. 2 N hydrochloric acid was added to the residue, and it was extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine, dried over anhydrous magnesium sulfate and concentrated, the residue was recrystallized from isopropyl ether to give the title compound(14.20g, 71.4%) as a white solid.

¹H-NMR(CDCl₃): δ 2.58(3H, s), 3.93(3H, s), 5.27(2H, s), 7.07(1H, d, J=8.7Hz), 7.26-7.43(3H, m), 7.47-7.50(2H, m), 8.07(1H, dd, J=8.7, 2.4Hz), 8.44(1H, d, J=2.4Hz).

(2) 5-Acetyl-2-benzyloxybenzoic acid.

[0411] 5-Acetyl-2-benzyloxybenzoic acid methyl ester(5.69g, 20mmol) was dissolved in a mixed solvent of methanol (20mL) and tetrahydrofuran(20mL). 2 N sodium hydroxide(11mL) was added dropwise, and the mixture was stirred for 8 hours. The solvent was evaporated under reduced pressure. 2 N hydrochloric acid was added to the residue, and it

EP 1 352 650 A1

was extracted with dichloromethane. After the dichloromethane layer was washed with water and brine, dried over anhydrous magnesium sulfate and concentrated, the residue was washed with isopropyl ether to give the title compound (4.92g, 91.0%) as a white solid.

¹H-NMR(DMSO-d₆): δ 2.55(3H, s), 5.32(2H, s), 7.30-7.43(4H, m), 7.49-7.52(2H, m), 8.09(1H, dd, J=9.0, 2.7Hz), 8.22(1H, d, J=2.4Hz).

(3) 5-Acetyl-2-benzyloxy-N-[3,5-bis(trifluoromethyl)phenyl]benzamide.

[0412] Using 5-acetyl-2-benzyloxybenzoic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 24 gave the title compound(5.47g, 63.1%) as a slight yellowish green solid.

¹H-NMR(DMSO-d₆): δ 2.57(3H, s), 7.11(1H, d, J=8.7Hz), 7.86(1H, s), 8.05(1H, dd, J=8.4, 2.1Hz), 8.44(1H, d, J=2.1Hz), 8.47(2H, s), 10.96(1H, s), 11.97(1H, brs).

(4) Preparation of 5-acetyl-N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide.

[0413] Ethanol(6mL) and tetrahydrofuran(72mL) were added to 5-acetyl-2-benzyloxy N-[3,5-bis(trifluoromethyl)phenyl]benzamide(602mg, 1.25mmol) and 5% palladium-carbon(60mg), and the mixture was hydrogenated at room temperature for 30 minutes. After the insoluble matter was filtered off, the solvent was evaporated under reduced pressure and the residue was recrystallized from n-hexane-ethyl acetate to give the title compound(230mg, 47.0%) as a white solid. ¹H-NMR(DMSO-d₆): δ 2.59(3H, s), 5.35(2H, s), 7.32-7.36(3H, m), 7.43(1H, d, J=8.7Hz), 7.52-7.55(2H, m), 7.82(1H, s), 8.16(1H, dd, J=8.7, 2.4Hz), 8.25(1H, d, J=2.4Hz), 8.31(2H, s), 10.89(1H, s).

Example 59: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(1-hydroxyethyl)benzamide (Comopund No. 57).

[0414] 5-Acetyl-N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide(50.5mg, 0.13mmol) was suspended in ethanol(2mL). Sodium borohydride(23.6mg, 0.62mmol) was added, and the mixture was stirred at room temperature for 12 hours. The reaction mixture was poured into diluted hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was washed with isopropyl ether/n-hexane under suspension to give the title compound (39.7mg, 78.3%) as white powder.

¹H-NMR(DMSO-d₆): δ 1.34(3H, d, J=6.3Hz), 4.71(1H, q, J=6.3Hz), 5.18(1H, brs), 6.97(1H, d, J=8.4Hz), 7.44(1H, dd, J=8.4, 2.1Hz), 7.84(1H, s), 7.86(1H, d, J=2.1Hz), 8.48(2H, s), 10.85(1H, s), 11.32(1H, s).

Example 60: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-[(1-methoxyimino)ethyl]benzamide (Comopund No. 58).

[0415] 5-Acetyl-N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide(100.0mg, 0.26mmol) was dissolved in ethanol(3mL). Pyridine(45μl, 0.56mmol) and O-methylhydroxylamine hydrochloride(25.8mg, 0.31mmol) were added, and the mixture was refluxed for 1 hour. After cooling, the reaction mixture was poured into diluted hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel (hexane:ethyl acetate=4:1) to give the title compound(102.1mg, 95.3%) as a white crystal.

¹H-NMR(DMSO-d₆): δ 2.19(3H, s), 3.91(3H, s), 7.05(1H, d, J=8.7Hz), 7.77(1H, dd, J=8.7, 2.4Hz), 7.85(1H, s), 8.09(1H, d, J=2.4Hz), 8.47(2H, s), 10.87(1H, s), 11.48(1H, s).

Example 61: 5-[(1-Benzyloxymino)ethyl]-N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 59).

[0416] Using 5-acetyl-N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide and O-benzylhydroxylamine hydrochloride as the raw materials, the same operation as the example 60 gave the title compound.

Yield: 79.9%.

¹H-NMR(DMSO-d₆): δ 2.24(3H, s), 5.20(2H, s), 7.04(1H, d, J=8.7Hz), 7.29-7.47(5H,m), 7.76(1H, dd, J=8.7, 2.4Hz), 7.85(1H, s), 8.07(1H, d, J=2.1Hz), 8.46(2H, s), 10.87(1H, s), 11.47(1H, s).

Example 62: N-[3,5-Bis(trifluoromethyl)phenyl]-5-(2,2-dicyanoethen-1-yl)-2-hydroxybenzamide (Comopund No. 60).

(1) 5-(2,2-Dicyanoethen-1-yl)-2-hydroxybenzoic acid.

[0417] Malononitrile(132mg, 2mmol) was dissolved in ethanol(6mL), and 5-formylsalicylic acid (332mg, 2mmol) was added. After cooling with ice bath, benzylamine(0.1mL) was added and the mixture was stirred at room temperature.

EP 1 352 650 A1

for 2 hours. The separated yellow crystal was filtered and recrystallized (ethanol) to give the title compound(139.9mg, 32.7%) as a light yellow solid.

¹H-NMR(DMSO-d₆): δ 7.12(1H, d, J=8.7Hz), 8.09(1H, dd, J=8.7, 2.4Hz), 8.41(1H, s), 8.50(1H, d, J=2.4Hz).

(2) N-[3,5-Bis(trifluoromethyl)phenyl]-5-(2,2-dicyanoethen-1-yl)-2-hydroxybenzamide.

[0418] Using 5-(2,2-dicyanoethen-1-yl)-2-hydroxybenzoic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 9-1%.

¹H-NMR(DMSO-d₆): δ 7.13(1H, d, J=9.0Hz), 7.83(1H, s), 8.04(1H, dd, J=9.0, 2.4Hz), 8.36(1H, s), 8.38(1H, d, J=2.4Hz), 8.43(2H, s), 11.43(1H, s).

Example 63: 3-((N-[3,5-Bis(trifluoromethyl)phenyl]carbamoyl)-4-hydroxyphenyl)-2-cyanoacrylic acid methyl ester (Comopund No. 62).

(1) 5-[(2-Cyano-2-methoxycarbonyl)ethen-1-yl]-2-hydroxybenzoic acid.

[0419] Triethylamine(0.2mL) was added to a mixture of 5-formylsalicylic acid(332mg, 2mmol). Cyanoacetic acid methyl ester(198mg, 2mmol) and acetic acid(6mL), and the mixture was refluxed for 5 hours. After cooling, the reaction mixture was poured into water, and the separated crystal was filtered and recrystallized (n-hexane) to give the title compound(327.7mg, 66.3%) as a light yellow solid.

¹H-NMR(DMSO-d₆): δ 3.85(3H, s), 7.15(1H, d, J=8.7Hz), 8.20(1H, dd, J=8.7, 2.4Hz), 8.37(1H, s), 8.66(1H, d, J=2.4Hz).

(2) 3-((N-[3,5-Bis(trifluoromethyl)phenyl]carbamoyl)-4-hydroxyphenyl)-2-cyanoacrylic acid methyl ester.

[0420] Using 5-[(2-cyano-2-methoxycarbonyl)ethen-1-yl]-2-hydroxybenzoic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 66.3%.

¹H-NMR(DMSO-d₆): δ 3.85(3H, s), 7.15(1H, d, J=8.7Hz), 8.20(1H, dd, J=8.7, 2.4Hz), 8.37(1H, s), 8.66(1H, d, J=2.4Hz).

Example 64: 3-((N-[3,5-Bis(trifluoromethyl)phenyl]carbamoyl)-4-hydroxyphenyl)-2-cyanoacrylic acid (Comopund No. 61).

[0421] 3-((N-[3,5-Bis(trifluoromethyl)phenyl]carbamoyl)-4-hydroxyphenyl)-2-cyanoacrylic acid methyl ester(50mg, 0.11mmol) was dissolved in ethanol(5mL). 2 N sodium hydroxide(0.11ml, 0.22mmol) was added, and the mixture was stirred at room temperature for 3 hours. The reaction mixture was poured into diluted hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with brine, dried over anhydrous magnesium sulfate, the residue obtained by evaporation under reduced pressure was recrystallized (ethyl acetate) to give the title compound(13.5mg, 30.4%) as a light yellow solid.

¹H-NMR(DMSO-d₆): δ 7.12(1H, d, J=8.4Hz), 7.84(1H, s), 7.94(1H, dd, J=8.4, 2.1Hz), 8.38(1H, d, J=2.1Hz), 8.45(2H, s), 9.87(1H, s), 11.41(1H, s).

Example 65: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(2-phenylethen-1-yl)benzamide (Comopund No. 63).

[0422] A mixture of N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-iodobenzamide(475mg, 1mmol), styrene(130mg, 1.25mmol), palladium acetate(4.5mg, 0.02mmol), tris(ortho-tolyl)phosphine(12.2mg, 0.04mmol), diisopropylamine(388mg, 3mmol) and N,N-dimethylformamide(2mL) was refluxed for 8 hours. After cooling, water was added to the reaction mixture, and it was extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine, dried over anhydrous magnesium sulfate and concentrated, the residue was purified by column chromatography on silica gel(hexane-isopropyl ether:2/1→1/1) to give the title compound(173mg, 38.3%) as a pale yellow solid.

¹H-NMR(DMSO-d₆): δ 7.04(1H, d, J=8.4Hz), 7.20-7.29(3H, m), 7.38(2H, t, J=7.5Hz), 7.59(2H, d, J=7.5Hz), 7.72(1H, dd, J=8.4, 2.1Hz), 7.86(1H, s), 8.07(1H, d, J=2.1Hz), 8.49(2H, s), 10.89(1H, s), 11.33(1H, brs).

Example 66: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-[(trimethylsilyl)ethynyl]benzamide (Comopund No. 66).

[0423] N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-iodobenzamide(950mg, 2mmol) and trimethylsilylacetylene(246mg, 2.5mmol) were dissolved in triethylamine(2mL) and N,N-dimethylformamide(4mL). Tetrakis(triphenylphosphine)palladium(23mg, 0.02mmol) and cuprous iodide(4mg, 0.02mmol) were added under argon atmosphere, and the

EP 1 352 650 A1

mixture was stirred at 40°C for 2 hours. After cooling to room temperature, the reaction mixture was poured into ethyl acetate(100ml and 1 N citric acid(100mL stirred, and filtered through celite. After the ethyl acetate layer was washed with water and brine one after another, dried over anhydrous magnesium sulfate, the solvent was evaporated under reduced pressure. The obtained residue was purified by column chromatography on silica gel(n-hexane:ethyl acetate=19:1) to give a light orange solid. This was crystallized by n-hexane to give the title compound(286mg, 32.1%) as a white crystal.

¹H-NMR(DMSO-d₆): δ 0.23(9H, s), 7.00(1H, d, J=8.7Hz), 7.54(1H, dd, J=8.7, 2.4Hz), 7.85(1H, s), 7.98(1H, d, J=2.1Hz), 8.46(2H, s), 10.86(1H, s), 11.69(1H, s).

Example 67: N-[3,5-Bis(trifluoromethyl)phenyl]-5-ethynyl-2-hydroxybenzamide (Comopund No. 64).

[0424] N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-[(trimethylsilyl)ethynyl]benzamide(233mg, 0.5mmol) was dissolved in methanol(1mL). 2 N sodium hydroxide(1mL) was added, and the mixture was stirred at room temperature for 1 hour. The reaction mixture was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine one after another, dried over anhydrous magnesium sulfate, the solvent was evaporated under reduced pressure. The obtained residue was crystallized from ethanol-water to give the title compound(67mg, 35.9%) as a light gray crystal.

¹H-NMR(DMSO-d₆): δ 4.11(1H, s), 7.02(1H, d, J=8.4Hz), 7.55(1H, dd, J=8.4, 2.1Hz), 7.85(1H, s), 7.98(1H, d, J=2.1Hz), 8.46(2H, s), 10.86(1H, s), 11.62(1H, s).

Example 68: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(phenylethynyl)benzamide (Comopund No. 65).

[0425] Using N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-iodobenzamide and phenylacetylene as the raw materials, the same operation as the example 66 gave the title compound.

¹H-NMR(DMSO-d₆): δ 7.06(1H, d, J=8.4Hz), 7.42-7.46(3H, m), 7.53-7.57(2H, m), 7.64(1H, dd, J=8.7, 2.1Hz), 7.86(1H, s), 8.06(1H, d, J=2.1Hz), 8.48(2H, s), 10.94(1H, s), 11.64(1H, brs).

Example 69: N-[3,5-Bis(trifluoromethyl)phenyl]-4-hydroxybiphenyl-3-carboxamide (Comopund No. 67).

[0426] N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-iodobenzamide(200mg, 0.42mmol) was dissolved in 1,2-dimethoxyethane(3mL). Tetrakis(triphenylphosphine)palladium(16mg, 0.0014mmol) was added under argon atmosphere, and the mixture was stirred at room temperature for 5 minutes. Then dihydroxyphenylborane(57mg, 0.47mmol) and 1M sodium carbonate(1.3mL) were added and refluxed for 2 hours. After cooling to room temperature, the reaction mixture was poured into diluted hydrochloric acid and extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine one after another, dried over anhydrous sodium sulfate, the solvent was evaporated under reduced pressure. The obtained residue was purified by column chromatography on silica gel(n-hexane:ethyl acetate=6:1→3:1) to give the title compound(109mg, 61.1%) as a white crystal.

¹H-NMR(DMSO-d₆): δ 7.12(1H, d, J=8.7Hz), 7.33-7.38(1H, m), 7.48(2H, t, J=7.5Hz), 7.67-7.70(2H, m), 7.79(1H, dd, J=8.4, 2.4Hz), 7.87(1H, s), 8.17(1H, d, J=2.4Hz), 8.49(2H, s), 10.92(1H, s), 11.41(1H, s).

Example 70: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(2-phenethyl)benzamide (Comopund No. 68).

[0427] Using N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-(phenylethynyl)benzamide as the raw material, the same operation as the example 58(4) gave the title compound.

Yield: 86.2%.

¹H-NMR(DMSO-d₆): δ 2.88(4H, s), 6.93(1H, d, J=8.1Hz), 7.15-7.34(6H, m), 7.76(1H, d, J=2.4Hz), 7.84(1H, s), 8.47(2H, s), 10.79(1H, s), 11.15(1H, s).

Example 71: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(trifluoromethyl)benzamide (Comopund No. 69).

[0428] Using 2-hydroxy-5-(trifluoromethyl)benzoic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound. (2-Hydroxy-5-(trifluoromethyl)benzoic acid: refer to Chem. Pharm. Bull., 1996, 44, 734.)

Yield: 44.7%.

¹H-NMR(CDCl₃, δ): 7.17(1H, d, J=9.0Hz), 7.72-7.75(2H, m), 7.86(1H, s), 8.17(2H, s), 8.35(1H, s), 11.88(1H, s).

Example 72: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(pentafluoroethyl)benzamide (Comopund No.70).

[0429] Using 2-hydroxy-5-(pentafluoroethyl)benzoic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound. (2-Hydroxy-5-(pentafluoroethyl)benzoic acid: refer to Chem. Pharm. Bull., 1996, 44, 734.)

Yield: 65.7%.

¹H-NMR(CDCl₃, δ): 7.19(1H, d, J=9.0Hz), 7.70(1H, dd, J=8.7, 2.1Hz), 7.81(1H, d, J=2.1Hz), 8.17(2H, s), 8.37(1H, s), 11.92(1H, s).

Example 73: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(pyrrol-1-yl)benzamide (Comopund No. 71).

[0430] Using 2-hydroxy-5-(pyrrol-1-yl)benzoic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 57.8%.

¹H-NMR(DMSO-d₆, δ): 6.27(2H, dd, J=2.4, 1.8Hz), 7.10(1H, d, J=9.0Hz), 7.29(2H, dd, J=2.4, 1.8Hz), 7.66(1H, dd, J=9.0, 2.7Hz), 7.86(1H, s), 7.98(1H, d, J=2.4Hz), 8.47(2H, s), 10.89(1H, s), 11.24(1H, s).

Example 74: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(thiophen-2-yl)benzamide (Comopund No. 72).

[0431] Using N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-iodobenzamide and 2-thiopheneboronic acid as the raw materials, the same operation as the example 69 gave the title compound.

¹H-NMR(DMSO-d₆, δ): 7.08(1H, d, J=8.4Hz), 7.14(1H, dd, J=5.4, 3.6Hz), 7.45(1H, dd, J=3.6, 1.2Hz), 7.51(1H, dd, J=5.1, 0.9Hz), 7.75(1H, dd, J=8.4, 2.4Hz), 7.59(1H, s), 8.08(1H, d, J=2.4Hz), 8.48(2H, s), 10.91(1H, s), 11.38(1H, s).

Example 75: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(thiophen-3-yl)benzamide (Comopund No. 73).

[0432] Using N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-iodobenzamide and 3-thiopheneboronic acid as the raw materials, the same operation as the example 69 gave the title compound.

Yield: 38.7%.

¹H-NMR(DMSO-d₆, δ): 7.06(1H, d, J=8.7Hz), 7.57(1H, dd, J=4.8, 1.5Hz), 7.66(1H, dd, J=4.8, 3.0Hz), 7.81-7.84(2H, m), 7.86(1H, s), 8.18(1H, d, J=2.1Hz), 8.49(2H, s), 10.90(1H, s), 11.33(1H, s).

Example 76: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(2-methylthiazol-4-yl)benzamide (Comopund No. 75).

(1) 2-Benzyloxy-5-(2-bromoacetyl)-N-[3,5-bis(trifluoromethyl)phenyl]benzamide.

[0433] 5-Acetyl-2-benzyloxy-N-[3,5-bis(trifluoromethyl)phenyl]benzamide(4.81g, 10mmol) was dissolved in THF (30ml). Phenyltrimethylammonium bromide(3.75g, 10mmol) was added, and the mixture was stirred at room temperature for 12 hours. The reaction mixture was poured into water and extracted with ethyl acetate. After the organic layer was washed with aqueous sodium hydrogen sulfite, water, and brine, dried over anhydrous magnesium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel(hexane:ethyl acetate=4:1), and recrystallized(ethyl acetate/hexane) to give the title compound(2.39g, 42.7%) as a white solid.

¹H-NMR(DMSO-d₆, δ): 4.91(2H, s), 5.36(2H, s), 7.32-7.35(3H, m), 7.47(1H, d, J=9.0Hz), 7.52-7.56(2H, m), 7.82(1H, s), 8.21(1H, dd, J=8.7, 2.4Hz), 8.29(1H, d, J=2.4Hz), 8.31(2H, s), 10.91(1H, s).

(2) 2-Benzyloxy-N-[3,5-bis(trifluoromethyl)phenyl]-5-(2-methylthiazol-4-yl)benzamide.

[0434] A mixture of 2-benzyloxy-5-(2-bromoacetyl)-N-[3,5-bis(trifluoromethyl)phenyl]benzamide(280mg, 0.5mmol), thioacetamide(41mg, 0.55mmol), sodium hydrogen carbonate(50mg, 0.6mmol) and ethanol(15mL) was refluxed for 1 hour. The reaction mixture was poured into water, neutralized by sodium hydrogen carbonate, and extracted with ethyl acetate. After the organic layer was washed with water and saturated brine, dried over anhydrous magnesium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel(hexane:ethyl acetate=4:1) to give the title compound(181mg, 67.5%) as a white solid.

¹H-NMR(DMSO-d₆, δ): 2.72(3H, s), 5.29(2H, s), 7.33-7.36(3H, m), 7.40(1H, d, J=9.0Hz), 7.54-7.57(2H, m), 7.81(1H, s), 7.94(1H, s), 8.12(1H, dd, J=8.7, 2.1Hz), 8.27(1H, d, J=2.1Hz), 8.31(2H, s), 10.86(1H, s).

EP 1 352 650 A1

(3) N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(2-methylthiazol-4-yl)benzamide.

[0435] 2-Benzyloxy-N-[3,5-bis(trifluoromethyl)phenyl]-5-(2-methylthiazol-4-yl)benzamide (160mg, 0.3mmol) and 10% Pd-C (240mg) were dissolved in ethanol (10ml) and stirred for 3.5 hours under hydrogen atmosphere. The reaction mixture was filtered and the filtrate was evaporated under reduced pressure to give the title compound (103.4mg, 79.2%) as a white solid.

¹H-NMR(DMSO-d₆): δ 2.72(3H, s), 7.08(1H, d, J=8.7Hz), 7.83(1H, s), 7.85(1H, s), 8.01(1H, dd, J=8.7, 2.4Hz), 8.42(1H, d, J=2.1Hz), 8.50(2H, s), 10.96(1H, s), 11.40(1H, s).

Example 77: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(imidazo[1,2-a]pyridin-2-yl)benzamide (Compound No. 75).

[0436] A mixture of 2-benzyloxy-5-(2-bromoacetyl)-N-[3,5-bis(trifluoromethyl)phenyl]benzamide (280mg, 0.5mmol), 2-aminopyridine (51.8mg, 0.55mmol), sodium hydrogen carbonate (50mg, 0.6mmol) and ethanol (10mL) was refluxed for 2 hours. After cooling, the reaction mixture was poured into aqueous sodium hydrogen carbonate and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous magnesium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel (n-hexane:ethyl acetate=1:2) to give the title compound (130.3mg, 45.9%) as a white solid. Then, a mixture of this solid (108mg, 0.19mmol), 10% Pd-C (11mg), ethanol (8mL) and ethyl acetate (8mL) was stirred for 7 hours under hydrogen atmosphere. The reaction mixture was filtered and the residue obtained by evaporation of the filtrate under reduced pressure was purified by chromatography on silica gel (n-hexane:ethyl acetate=1:3) to give the title compound (18.3mg, 20.2%) as a white solid.

¹H-NMR(DMSO-d₆): δ 6.90(1H, dt, J=6.6, 0.9Hz), 7.10(1H, d, J=8.7Hz), 7.25(1H, m), 7.57(1H, d, J=9.0Hz), 7.86(1H, s), 8.04(1H, dd, J=8.7, 2.1Hz), 8.35(1H, s), 8.48-8.56(4H, m), 11.00(1H, s), 11.41(1H, s).

Example 78: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(pyridin-2-yl)benzamide (Compound No. 76).

(1) N-[3,5-Bis(trifluoromethyl)phenyl]-5-iodo-2-methoxymethoxybenzamide.

[0437] A mixture of N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-iodobenzamide (4.75g, 10mmol), chloromethyl methyl ether (1.14ml, 15ml), potassium carbonate (2.76g, 20mmol) and acetone (50mL) was refluxed for 8 hours. The reaction mixture was poured into diluted hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous magnesium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel (hexane:ethyl acetate=3:1), and recrystallized (n-hexane/ethyl acetate) to give the title compound (3.96g, 76.3%) as a white solid.

¹H-NMR(DMSO-d₆): δ 3.38(3H, s), 5.28(2H, s), 7.12(1H, d, J=9.0Hz), 7.81(1H, s), 7.82(1H, dd, J=8.7, 2.4Hz), 7.88(1H, d, J=2.4Hz), 8.40(2H, s), 10.87(1H, s).

(2) N-[3,5-Bis(trifluoromethyl)phenyl]-2-methoxymethoxy-5-(pyridin-2-yl)benzamide.

[0438] N-[3,5-Bis(trifluoromethyl)phenyl]-5-iodo-2-methoxymethoxybenzamide (0.20g, 0.39mmol) was dissolved in N,N-dimethylformamide (8ml). Tri-n-butyl(2-pyridyl)tin (0.13ml, 0.41mmol) and dichlorobis(triphenylphosphine)palladium (32.1mg, 0.05mmol) were added, and the mixture was stirred at 100°C for 1.5 hours. After cooling, the reaction mixture was poured into water and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel (n-hexane:ethyl acetate=2:1→1:1) to give the title compound (37.9mg, 20.8%) as a white powder.

¹H-NMR(CDCl₃): δ 3.64(3H, s), 5.53(2H, s), 7.23-7.28(1H, m), 7.36(1H, d, J=8.7Hz), 7.65(1H, s), 7.77-7.84(2H, m), 8.20(2H, s), 8.31(1H, dd, J=8.7, 2.4Hz), 8.68-8.70(1H, m), 8.83(1H, d, J=2.4Hz), 10.12(1H, s).

(3) N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(pyridin-2-yl)benzamide.

[0439] Methanol (3ml) and concentrated hydrochloric acid (0.5ml) were added to N-[3,5-bis(trifluoromethyl)phenyl]-2-methoxymethoxy-5-(pyridin-2-yl)benzamide (37.9 mg, 0.08mmol), and the mixture was refluxed for 2 hours. After cooling, the reaction mixture was poured into saturated aqueous sodium hydrogen carbonate and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel (n-hexane:ethyl acetate=2:1) to give the title compound (16.2mg, 47.2%) as a white powder.

EP 1 352 650 A1

¹H-NMR(DMSO-d₆): δ 7.13(1H, d, J=8.4Hz), 7.73(1H, ddd, J=7.5, 6.3, 1.2Hz), 7.86-7.91(2H, m), 7.97(1H, d, J=7.8Hz), 8.20(1H, dd, J=8.7, 2.1Hz), 8.50(2H, s), 8.59(1H, d, J=2.4Hz), 8.64-8.66(1H, m), 10.97(1H, s), 11.53(1H, s).

Example 79: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-methoxybenzamide (Comopund No. 77).

[0440] Using 5-methoxysalicylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 56.8%.

¹H-NMR(DMSO-d₆): δ 3.77(3H, s), 6.97(1H, d, J=9.0Hz), 7.10(1H, dd, J=9.0 3.0Hz), 7.43(1H, d, J=3.0Hz), 7.84(1H, s), 8.47(2H, s), 10.84(1H, s), 10.91(1H, s).

Example 80: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-isobutyrylbenzamide (Comopund No. 79).

(1) 5-Acetyl-2-methoxybenzoic acid methyl ester.

[0441] A mixture of 5-acetylsalicylic acid methyl ester(5.00g, 25.7mmol), sodium carbonate(7.10g, 51.4mmol) and N,N-dimethylformamide(25mL) was cooled with ice bath. Methyl iodide(2.5mL, 40.1mmol) was added, and the mixture was stirred at room temperature for 3 hours. The reaction mixture was poured into water, neutralized by hydrochloric acid, and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was washed under suspension(isopropyl ether/n-hexane) to give the title compound(5.17g, 96.5%) as a white crystal.

¹H-NMR(CDCl₃): δ 2.59(3H, s), 3.92(3H, s), 3.99(3H, s), 7.04(1H, d, J=8.7Hz), 8.12(1H, dd, J=8.7, 2.4Hz), 8.41(1H, d, J=2.4Hz).

(2) 5-Isobutyryl-2-methoxybenzoic acid methyl ester.

[0442] A mixture of 5-acetyl-2-methoxybenzoic acid methyl ester(0.50g, 2.40mmol), potassium tert-butoxide(0.81g, 7.22mmol) and tetrahydrofuran(10mL) was cooled with ice bath. Methyl iodide(0.5mL, 8.03mmol) was added, and the mixture was stirred at room temperature for 1 hour. The reaction mixture was poured into water, neutralized by hydrochloric acid, and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel(n-hexane:ethyl acetate=3:1→2:1) to give the title compound(143.1mg, 25.2%) as a light yellow oil.

¹H-NMR(CDCl₃): δ 1.22(6H, d, J=6.9Hz), 3.52(1H, m), 3.92(3H, s), 3.98(3H, s), 7.05(1H, d, J=8.7Hz), 8.13(1H, dd, J=8.7, 2.4Hz), 8.42(1H, d, J=2.4Hz).

(3) 5-Isobutyryl-2-methoxybenzoic acid.

[0443] 5-Isobutyryl-2-methoxybenzoic acid methyl ester(143.1mg, 0.60mmol) was dissolved in methanol(5mL). 2 N aqueous sodium hydroxide(1ml) was added, and the mixture was refluxed for 1 hour. After cooling, the reaction mixture was poured into 2 N hydrochloric acid and extracted with ethyl acetate. The organic layer was washed with water and brine, dried over anhydrous sodium sulfate, and evaporated under reduced pressure to give the title compound(134mg, yield:quantitative) as a white crystal.

¹H-NMR(CDCl₃): δ 1.22(6H, d, J=6.9Hz), 3.59(1H, m), 4.15(3H, s), 7.16(1H, d, J=8.7Hz), 8.24(1H, dd, J=8.7, 2.4Hz), 8.73(1H, d, J=2.1Hz).

(4) 5-Butyryl-N-[3,5-bis(trifluoromethyl)phenyl]-2-methoxybenzamide.

[0444] Using 5-isobutyryl-2-methoxybenzoic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

¹H-NMR(CDCl₃): δ 1.23(6H, d, J=6.9Hz), 3.64(1H, m), 4.20(3H, s), 7.18(1H, d, J=8.7Hz), 7.65(1H, s), 8.19(2H, s), 8.22(1H, dd, J=8.7, 2.1Hz), 8.88(1H, d, J=2.1Hz), 9.98(1H, s).

(5) N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-isobutyrylbenzamide.

[0445] A mixture of 5-butyryl-N-[3,5-bis(trifluoromethyl)phenyl]-2-methoxybenzamide(143.4mg, 0.33mmol), 2,4,6-collidine(3ml) and lithium iodide(53.1mg, 0.40mmol) was refluxed for 1 hour. After cooling, the reaction mixture was poured into 2N hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by

EP 1 352 650 A1

chromatography on silica gel(n-hexane:ethyl acetate=3:1) and crystallized(ethyl acetate/isopropyl ether) to give the title compound(90.3mg, 65.3%) as a white crystal.

¹H-NMR(DMSO-d₆): δ 1.12(6H, d, J=6.9Hz), 3.66(1H, m), 7.12(1H, d, J=8.4Hz), 7.85(1H, s), 8.07(1H, dd, J=8.4, 2.4Hz), 8.45(1H, d, J=2.4Hz), 8.47(2H, s), 10.93(1H, s), 11.95(1H, brs).

Example 81: N-[3,5-Bis(trifluoromethyl)phenyl]-4-hydroxyisophthalamic acid methyl ester (Comopund No. 81).

[0446] Using 4-hydroxyisophthalic acid 1-methyl ester and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 91.5%.

¹H-NMR(DMSO-d₆): δ 3.85(3H, s), 7.12(1H, d, J=8.4Hz), 7.86(1H, s), 8.02(1H, dd, J=8.7, 2.4Hz), 8.46-8.47(3H, m), 10.96(1H, s), 12.03(1H, brs).

Example 82: N-[3,5-Bis(trifluoromethyl)phenyl]-4-hydroxyisophthalamic acid (Comopund No. 80).

[0447] N-[3,5-Bis(trifluoromethyl)phenyl]-4-hydroxyisophthalamic acid methyl ester(2.85g, 7mmol) was suspended in a mixed solvent of methanol(14mL) and tetrahydrofuran(14mL). 2 N aqueous sodium hydroxide(14mL) was added, and the mixture was refluxed for 2 hours. After cooling, the reaction mixture was added 2 N hydrochloric acid(20ml) and the separated solid was filtered, washed with water, dried to give the title compound (2.68g, 97.4%) as a white crystal.

¹H-NMR(DMSO-d₆): δ 7.10(1H, d, J=8.7Hz), 7.82(1H, s), 7.86(1H, s), 8.01(1H, dd, J=8.7, 2.4Hz), 8.47(2H, s), 8.48(1H, d, J=2.4Hz), 10.97(1H, s), 11.98(1H, brs).

Example 83: N¹,N³-Bis[3,5-bis(trifluoromethyl)phenyl]-4-hydroxyisophthalamide (Comopund No. 82).

[0448] Using 4-hydroxyisophthalic acid(182mg, 1mmol), 3,5-bis(trifluoromethyl)-aniline(687mg, 3mmol), phosphorus trichloride(87 μl; 1mmol) and toluene(10mL) the same operation as the example 16 gave the title compound(151mg, 25.0%) as a white crystal.

¹H-NMR(DMSO-d₆): δ 7.18(1H, d, J=8.7Hz), 7.82(1H, s), 7.86(1H, s), 8.11(1H, dd, J=8.7, 2.4Hz), 8.50(2H, s), 8.54(2H, s), 8.56(1H, d, J=2.4Hz), 10.79(1H, s), 10.99(1H, s), 11.84(1H, brs).

Example 84: N³-[3,5-Bis(trifluoromethyl)phenyl]-4-hydroxy-N¹, N¹-dimethylisophthalamide (Comopund No. 83).

(1) 4-Benzyloxy-N-[3,5-bis(trifluoromethyl)phenyl]isophthalamic acid methyl ester.

[0449] Sodium hydride(60%; 1.04g, 26mmol) was washed with n-hexane, suspended in N,N-dimethylformamide (100mL). A solution of N-[3,5-bis(trifluoromethyl)-phenyl]-4-hydroxyisophthalamic acid methyl ester(8.15g, 20mmol) in N,N-dimethylformamide(100mL) was added dropwise under cooling with ice bath. After the addition was finished, the mixture was stirred at room temperature for 1 hour. A solution of benzyl bromide(4.45g, 26mmol) in N,N-dimethylformamide(10mL) was added, and the mixture was stirred at 60°C for 3 hours. After cooling, the reaction mixture was poured into ice and water, and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous magnesium sulfate, the residue obtained by evaporation under reduced pressure was recrystallized(ethyl acetate/n-hexane) to give the title compound(5.38g, 54.1%) as a white solid.

¹H-NMR(DMSO-d₆): δ 3.87(3H, s), 5.33(2H, s), 7.33-7.36(3H, m), 7.46(1H, d, J=8.7Hz), 7.53-7.56(2H, m), 7.82(1H, s), 8.15(1H, dd, J=8.7, 2.1Hz), 8.25(1H, d, J=2.1Hz), 8.28(2H, s), 10.87(1H, s).

(2) 4-Benzyloxy-N-[3,5-bis(trifluoromethyl)phenyl]isophthalamic acid.

[0450] Using 4-benzyloxy-N-[3,5-bis(trifluoromethyl)phenyl]isophthalamic acid methyl ester as the raw material, the same operation as the example 82 gave the title compound.

Yield: 79.7%.

¹H-NMR(DMSO-d₆): δ 5.32(2H, s), 7.32-7.34(3H, m), 7.43(1H, d, J=8.7Hz), 7.52-7.56(2H, m), 7.81(1H, s), 8.12(1H, dd, J=8.7, 2.1Hz), 8.22(1H, d, J=2.1Hz), 8.28(2H, s), 10.85(1H, s), 13.81(1H, brs).

(3) 4-Benzyloxy-N³-[3,5-bis(trifluoromethyl)phenyl]-N¹,N¹-dimethylisophthalamide.

[0451] WSC · HCl(95mg, 0.50mmol) was added to a solution of 4-benzyl-N-[3,5-bis(trifluoromethyl)phenyl]isophthalamic acid(242mg, 0.50mmol), dimethylamine hydrochloride(41mg, 0.50mmol) and triethylamine(51mg, 0.50mmol) in

EP 1 352 650 A1

tetrahydrofuran(5mL) under ice cooling, and the mixture was stirred at room temperature for 3 hours. The reaction mixture was poured into water and extracted with ethyl acetate. After the organic layer was washed with diluted hydrochloric acid, water and brine, dried over anhydrous magnesium sulfate, the residue obtained by evaporating the solvent under reduced pressure was purified by chromatography on silica gel(hexane:ethyl acetate=1:4) to give the title compound(165mg, 64.9%) as a white solid.

¹H-NMR(DMSO-d₆): δ 2.99(6H, s), 5.29(2H, s), 7.32-7.38(4H, m), 7.52-7.56(2H, m), 7.64(1H, dd, J=8.7, 2.1Hz), 7.73(1H, d, J=2.1Hz), 7.80(1H, s), 8.28(2H, s), 10.83(1H, s).

(4) N³-[3,5-bis(trifluoromethyl)phenyl]-4-hydroxy-N¹,N¹-dimethylisophthalamide.

[0452] A solution of 4-benzyloxy-N³-[3,5-bis(trifluoromethyl)phenyl]-N¹,N¹-dimethylisophthalamide(141mg, 0.28mmol) and 5% Pd-C(14mg) in the mixture of ethanol(5ml) and ethyl acetate(5ml) was stirred at room temperature for 1 hour under hydrogen atmosphere. The reaction mixture was filtered and the filtrate was evaporated under reduced pressure to give the title compound(106mg, 91.2%) as a white solid.

¹H-NMR(DMSO-d₆): δ 2.98(6H, s), 7.02(1H, d, J=8.7Hz), 7.52(1H, dd, J=8.7, 2.1Hz), 7.84(1H, s), 7.95(1H, d, J=2.1Hz), 8.46(2H, s), 11.10(1H, brs), 11.63(1H, brs).

Example 85: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(piperidine-1-carbonyl)benzamide.

(1) 2-Benzyloxy-N-[3,5-bis(trifluoromethyl)phenyl]-5-(piperidine-1-carbonyl)benzamide.

[0453] Using 4-benzyl-N-[3,5-bis(trifluoromethyl)phenyl]isophthalamic acid and piperidine as the raw materials, the same operation as the example 84(3) gave the title compound.

Yield: 56.4%.

¹H-NMR(CDCl₃): δ 1.53-1.70(6H, m), 3.44(2H, brs), 3.70(2H, brs), 5.26(2H, s), 7.24(1H, d, J=8.7Hz), 7.26(1H, s), 7.52-7.58(5H, m), 7.66(2H, s), 7.74(1H, dd, J=8.7, 2.4Hz), 8.37(1H, d, J=2.1Hz), 10.27(1H, s).

(2) N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(piperidine-1-carbonyl)benzamide.

[0454] Using 2-benzyloxy-N-[3,5-bis(trifluoromethyl)phenyl]-5-(piperidine-1-carbonyl)benzamide as the raw material, the same operation as the example 84(4) gave the title compound.

Yield: 96.3%, white solid.

¹H-NMR(DMSO-d₆): δ 1.51(4H, brs), 1.60-1.65(2H, m), 3.47(4H, brs), 7.04(1H, d, J=8.4Hz), 7.48(1H, dd, J=8.4, 2.1Hz), 7.85(1H, s), 7.92(1H, d, J=2.1Hz), 8.46(2H, s), 10.99(1H, s), 11.64(1H, brs).

Example 86: 5-(4-Benzylpiperidine-1-carbonyl)-N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide.

(1) 2-Benzyl-5-(4-benzylpiperidine-1-carbonyl)-N-[3,5-bis(trifluoromethyl)phenyl]benzamide.

[0455] Using 4-benzyl-N-[3,5-bis(trifluoromethyl)phenyl]isophthalamic acid and 4-benzylpiperidine as the raw materials, the same operation as the example 84(3) gave the title compound.

Yield: 76.7%.

¹H-NMR(CD₃OD): δ 1.18-1.38(2H, m), 1.67(1H, brs), 1.74(1H, brs), 1.84-1.93(1H, m), 2.60(2H, d, J=7.2Hz), 2.83(1H, brs), 3.10(1H, brs), 3.78(1H, brs), 4.59(1H, brs), 5.34(2H, s), 7.15-7.18(3H, m), 7.24-7.28(2H, m), 7.40-7.46(4H, m), 7.57-7.63(3H, m), 7.65(1H, dd, J=8.7, 2.4Hz), 7.96(2H, s), 8.05(1H, d, J=2.1Hz).

(2) N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(4-benzylpiperidine-1-carbonyl)benzamide.

[0456] Using 2-benzyl-5-(4-benzylpiperidine-1-carbonyl)-N-[3,5-bis(trifluoromethyl)phenyl]benzamide as the raw material, the same operation as the example 84(4) gave the title compound.

Yield: 54.3%, white solid.

¹H-NMR(DMSO-d₆): δ 1.08-1.22(2H, m), 1.59-1.62(2H, m), 1.77-1.80(1H, m), 2.50-2.55(2H, m), 2.87(2H, brs), 3.75(1H, br), 4.39(1H, br), 7.06(1H, d, J=8.4Hz), 7.17-7.20(3H, m), 7.28(2H, t, J=7.2Hz), 7.49(1H, dd, J=8.4, 2.1Hz), 7.84(1H, s), 7.93(1H, d, J=2.1Hz), 8.47(2H, s), 10.89(1H, s), 11.65(1H, s).

EP 1 352 650 A1

Example 87: N-[3,5-Bis(trifluoromethyl)phenyl]-5-dimethylsufamoyl-2-hydroxybenzamide.

(1) 2-Methoxy-5-sulfamoylbenzoic acid.

[0457] Methyl 2-methoxy-5-sulfamoylbenzoate(4.91g, 20mmol) was dissolved in methanol(30mL). 2 N aqueous sodium hydroxide(30mL, 60mmol) was added, and the mixture was stirred at room temperature for 1 hour. The reaction mixture was poured into 2 N hydrochloric acid, and the separated solid was filtered to give the title compound(4.55g, 98.3%) as a white solid.

¹H-NMR(DMSO-d₆): δ 3.89(3H, s), 7.30(1H, d, J=8.7Hz), 7.32(2H, s), 7.92(1H, dd, J=8.7, 2.7Hz), 8.09(1H, d, J=2.7Hz), 13.03(1H, br).

(2) N-[3,5-Bis(trifluoromethyl)phenyl]-2-methoxy-5-sufamoylbenzamide.

[0458] Using 2-methoxy-5-sulfamoylbenzoic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 24 gave the title compound.

Yield: 24.2%.

¹H-NMR(DMSO-d₆): δ 3.97(3H, s), 7.38(2H, s), 7.39(1H, d, J=8.7Hz), 7.85(1H, s), 7.96(1H, dd, J=8.7, 2.4Hz), 8.06(1H, d, J=2.4Hz), 8.43(2H, s), 10.87(1H, s).

(3) N-[3,5-Bis(trifluoromethyl)phenyl]-5-dimethylsufamoyl-2-methoxybenzamide.

[0459] A suspension of N-[3,5-bis(trifluoromethyl)phenyl]-2-methoxy-5-sufamoylbenzamide(442mg, 1.0mmol methyl iodide(710mg, 5.0mmol) and sodium carbonate(415mg, 3.0mmol in acetonitrile(10mL) was refluxed for 3 hours. After cooling to room temperature, the reaction mixture was poured into water and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous magnesium sulfate, the residue obtained by evaporating the solvent under reduced pressure was recrystallized from a mixed solvent of n-hexane and ethyl acetate (2:1) to give the title compound(207mg, 44.1%) as a white solid.

¹H-NMR(DMSO-d₆): δ 2.62(6H, s), 3.99(3H, s), 7.45(1H, d, J=9.0Hz), 7.85(1H, s), 7.91(1H, dd, J=8.7, 2.4Hz), 7.95(1H, d, J=2.4Hz), 8.43(2H, s), 10.90(1H, s).

(4) N-[3,5-Bis(trifluoromethyl)phenyl]-5-dimethylsufamoyl-2-hydroxybenzamide.

[0460] Using N-[3,5-bis(trifluoromethyl)phenyl]-5-dimethylsufamoyl-2-methoxybenzamide as the raw material, the same operation as the example 80(5) gave the title compound.

¹H-NMR(DMSO-d₆): δ 2.77(3H, d, J=4.5Hz), 4.37(1H, brs), 6.70(1H, d, J=3.6Hz), 7.04(2H, s).

Example 88: N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(pyrrole-1-sulfonyl)benzamide (Compound No. 87).

(1) N-[3,5-Bis(trifluoromethyl)phenyl]-2-methoxy-5-(pyrrole-1-sulfonyl)benzamide.

[0461] A mixture of N-[3,5-bis(trifluoromethyl)phenyl]-2-methoxy-5-sulfamoylbenzamide(442mg, 1mmol), 2,5-dimethoxytetrahydrofuran(159mg, 1.2mmol) and acetic acid(5mL) was refluxed for 2 hours. After cooling, the reaction mixture was poured into water and extracted with ethyl acetate. After the organic layer was washed with water, saturated aqueous sodium hydrogen carbonate and brine, dried over anhydrous magnesium sulfate, the residue obtained by evaporating the solvent under reduced pressure was purified by chromatography on silica gel(n-hexane:ethyl acetate=3:2) to give the title compound(436.5mg, 88.6%) as a white solid.

¹H-NMR(DMSO-d₆): δ 3.96(3H, s), 6.36(2H, d, J=2.4, 2.1Hz), 7.37(2H, dd, J=2.4, 2.1Hz), 7.42(1H, d, J=9.0Hz), 7.85(1H, s), 8.80(1H, dd, J=9.0, 2.4Hz), 8.18(1H, d, J=2.7Hz), 8.38(2H, s), 10.92(1H, s).

(2) N-[3,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-(pyrrole-1-sulfonyl)benzamide.

[0462] Using N-[3,5-bis(trifluoromethyl)phenyl]-2-methoxy-5-(pyrrole-1-sulfonyl)benzamide as the raw material, the same operation as the example 80(5) gave the title compound.

Yield: 79.4%.

¹H-NMR(DMSO-d₆, δ): 6.36(2H, dd, J=2.4, 2.1Hz), 7.18(1H, d, J=9.0Hz), 7.34(2H, d, J=2.4, 2.1Hz), 7.86(1H, s), 7.99(1H, dd, J=9.0, 2.7Hz), 8.31(1H, d, J=2.7Hz), 8.42(2H, s), 10.98(1H, s).

[0463] Using N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-nitrobenzamide as the raw material, the same operation as the example 84(4) gave the title compound. Yield: 98.0%.

5

[0464] Using 5-dimethylaminosalicylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

10

15

20

25

30

35

40

45

Yield: 11.3%.

50

55

¹H-NMR(DMSO-d₆): δ 6.87(1H, t, J=6.0Hz), 7.22(1H, d, J=8.7Hz), 7.21-7.23(1H, m), 7.77(1H, t, J=8.4Hz), 7.87(1H, t)

EP 1 352 650 A1

s), 7.95-7.98(3H, m), 8.03-8.07(4H, m), 8.47(1H, d, J=2.4Hz), 8.49(2H, s), 11.14(1H, s), 12.03(1H, br).

Example 96: 2-Acetoxy-N-[3,5-bis(trifluoromethyl)phenyl]-5-chlorobenzamide (Comopund No. 96).

[0470] N-[3,5-Bis(trifluoromethyl)phenyl]-5-chloro-2-hydroxybenzamide(1.51g, 3mmol) and pyridine(285mg, 3.6mmol) were dissolved in tetrahydrofuran(6mL). Acetyl chloride(234mg, 3.3mmol) was added dropwise under ice cooling, and the mixture was stirred at room temperature for 1 hour. The solvent was evaporated under reduced pressure. 2 N hydrochloric acid was added to the residue, and it was extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine, dried over anhydrous magnesium sulfate and concentrated, the residue was recrystallized from n-hexane-ethyl acetate to give the title compound(1.06g, 83.0%) as a white solid.

¹H-NMR(DMSO-d₆): δ 2.22(3H, s), 7.35(1H, d, J=9.0Hz), 7.71(1H, dd, J=8.7, 2.7Hz), 7.85(1H, s), 7.88(1H, d, J=2.7Hz), 8.37(2H, s), 11.05(1H, brs).

Example 97: 4-Acetylamino-N-(3,5-bis(trifluoromethyl)phenyl)-5-chloro-2-hydroxybenzamide (Comopund No. 97).

(1) 4-Acetylamino-5-chloro-2-methoxybenzoic acid.

[0471] Using 4-acetylamino-5-chloro-2-methoxybenzoic acid methyl ester as the raw material, the same operation as the example 82 gave the title compound.

Yield: 88.0%.

¹H-NMR(DMSO-d₆): δ 2.16(3H, s), 3.78(3H, s), 7.72(1H, s), 7.77(1H, s), 9.57(1H, s), 12.74(1H, s).

(2) 4-Acetylamino-N-[3,5-bis(trifluoromethyl)phenyl]-5-chloro-2-methoxybenzamide.

[0472] Using 4-acetylamino-5-chloro-2-methoxybenzoic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 24 gave the title compound.

Yield: 23.8%.

¹H-NMR(DMSO-d₆): δ 2.17(3H, s), 3.89(3H, s), 7.77-7.82(3H, m), 8.45-8.49(2H, m), 9.66(1H, s), 10.68(1H, s).

(3) 4-Acetylamino-N-[3,5-bis(trifluoromethyl)phenyl]-5-chloro-2-hydroxybenzamide.

[0473] Using 4-acetylamino-N-[3,5-bis(trifluoromethyl)phenyl]-5-chloro-2-methoxybenzamide as the raw material, the same operation as the example 80 gave the title compound.

Yield: 72.8%.

¹H-NMR(DMSO-d₆): δ 2.17(3H, s), 7.75(1H, s), 7.82(1H, s), 7.95(1H, s), 8.44(2H, s), 9.45(1H, s), 11.16(1H, brs), 11.63(1H, brs).

Example 98: N-[3,5-Bis(trifluoromethyl)phenyl]-4-chloro-2-hydroxybenzamide (Comopund No. 98).

[0474] Using 4-chlorosalicylic acid and 3,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 55.8%.

¹H-NMR(DMSO-d₆): δ 7.05-7.08(2H, m), 7.84-7.87(2H, m), 8.45(2H, s), 10.84(1H, s), 11.64(1H, brs).

Example 99: N-[3,5-Bis(trifluoromethyl)-2-bromophenyl]-5-chloro-2-hydroxybenzamide (Comopund No. 99).

[0475] Using 5-chlorosalicylic acid and 3,5-bis(trifluoromethyl)-2-bromoaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 14.5%.

¹H-NMR(DMSO-d₆): δ 7.11(1H, d, J=9.0Hz), 7.53(1H, dd, J=9.0, 2.7Hz), 7.91(1H, d, J=1.8Hz), 7.98(1H, d, J=2.7Hz), 9.03(1H, d, J=1.8Hz), 11.26(1H, brs).

Example 100: N-[2,5-Bis(trifluoromethyl)phenyl]-5-chloro-2-hydroxybenzamide (Comopund No. 100).

[0476] Using 5-chlorosalicylic acid and 2,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 3.6%.

¹H-NMR(CDCl₃): δ 7.03(1H, d, J=8.7Hz), 7.43-7.48(2H, m), 6.61(1H, d, J=8.1Hz), 7.85(1H, d, J=8.4Hz), 8.36(1H, brs),

EP 1 352 650 A1

8.60(1H, s), 11.31(1H, s).

Example 101: N-[2,5-Bis(trifluoromethyl)phenyl]-5-bromo-2-hydroxybenzamide (Comopund No. 101).

[0477] Using 5-bromosalicylic acid and 2,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 24.0%.

¹H-NMR(DMSO-d₆): δ 7.03(1H, d, J=8.7Hz), 7.65(1H, dd, J=8.7, 2.7Hz), 7.76(1H, d, J=8.4Hz), 8.03(1H, d, J=8.1Hz), 8.11(1H, d, J=2.7Hz), 8.74(1H, s), 11.02(1H, s), 12.34(1H, s).

Example 102: N-[2,5-Bis(trifluoromethyl)phenyl]-2-hydroxy-5-methylbenzamide (Comopund No. 102).

[0478] Using 5-methylsalicylic acid and 2,5-bis(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 1.5%.

¹H-NMR(CDCl₃): δ 2.36(3H, m), 6.97(1H, d, J=8.4Hz), 7.23(1H, s), 7.32(1H, dd, J=8.4, 1.5Hz), 7.57(1H, d, J=8.4Hz), 7.83(1H, d, J=8.4Hz), 8.46(1H, s), 8.69(1H, s), 11.19(1H, s).

Example 103: 2-Acetoxy-N-[2,5-bis(trifluoromethyl)phenyl]-5-chlorobenzamide (Comopund No. 103).

[0479] Using N-[2,5-bis(trifluoromethyl)phenyl]-5-chloro-2-hydroxybenzamide and acetyl chloride as the raw materials, the same operation as the example 96 gave the title compound.

Yield: 6.6%.

¹H-NMR(CDCl₃): δ 2.35(3H, s), 7.17(1H, d, J=8.7Hz), 7.54(1H, dd, J=8.7, 2.4Hz), 7.55(1H, d, J=8.1Hz), 7.80(1H, d, J=8.1Hz), 7.95(1H, d, J=2.4Hz), 8.60(1H, s), 8.73(1H, s).

Example 104: 5-Chloro-2-hydroxy-N-[2-(trifluoromethyl)phenyl]benzamide (Comopund No. 104).

[0480] Using 5-chlorosalicylic acid and 2-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 58.0%.

¹H-NMR(DMSO-d₆): δ 7.07(1H, d, J=8.7Hz), 7.42(1H, t, J=7.5Hz), 7.52(1H, d, J=8.7, 2.7Hz), 7.74(1H, t, J=8.1Hz), 7.77(1H, t, J=8.1Hz), 7.99(1H, d, J=2.7Hz), 8.18(1H, d, J=8.1Hz), 10.76(1H, s), 12.22(1H, s).

Example 105: 5-Chloro-N-[4-chloro-2-(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 105).

[0481] Using 5-chlorosalicylic acid and 4-chloro-2-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 21.5%.

¹H-NMR(DMSO-d₆): δ 7.07(1H, d, J=8.7Hz), 7.52(1H, dd, J=8.7, 2.7Hz), 7.80-7.85(2H, m), 7.97(1H, d, J=2.7Hz), 8.26(1H, d, J=8.4Hz), 10.80(1H, s), 12.26(1H, s).

Example 106: 5-Bromo-2-hydroxy-N-[3-(trifluoromethyl)phenyl]benzamide (Comopund No. 106).

[0482] Using 5-bromosalicylic acid and 3-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 50.3%.

¹H-NMR(DMSO-d₆, δ): 6.98(1H, d, J=8.7Hz), 7.48-7.52(1H, m), 7.59(1H, dd, J=8.7, 2.7Hz), 7.62(1H, t, J=8.1Hz), 7.92-7.96(1H, m), 8.02(1H, d, J=2.4Hz), 8.20(1H, s), 10.64(1H, s), 11.60(1H, s).

Example 107: 5-Chloro-N-[2-fluoro-3-(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 107).

[0483] Using 5-chlorosalicylic acid and 2-fluoro-3-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 71.7%, white solid.

¹H-NMR(DMSO-d₆): δ 7.07(1H, d, J=9.0Hz), 7.46(1H, t, J=7.8Hz), 7.52(1H, dd, J=9.0, 2.7Hz), 7.58(1H, t, J=7.2Hz), 7.96(1H, d, J=2.7Hz), 8.49(1H, t, J=7.2Hz), 10.82(1H, s), 12.13(1H, brs).

EP 1 352 650 A1

Example 108: 5-Chloro-N-[4-fluoro-3-(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 108).

[0484] Using 5-chlorosalicylic acid and 4-fluoro-3-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 72.1%, white solid.

¹H-NMR(DMSO-d₆): δ 7.03(1H, d, J=9.0Hz), 7.48(1H, dd, J=8.7, 2.7Hz), 7.56(1H, d, J=9.9Hz), 7.90(1H, d, J=2.7Hz), 7.99-8.03(1H, m), 8.21(1H, dd, J=6.6, 2.4Hz), 10.63(1H, s), 11.58(1H, s).

Example 109: 5-Bromo-N-[4-chloro-3-(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 109).

[0485] Using 5-bromosalicylic acid and 4-chloro-3-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 37.4%.

¹H-NMR(DMSO-d₆): δ 6.98(1H, d, J=8.7Hz), 7.59(1H, dd, J=8.7, 2.4Hz), 7.73(1H, d, J=8.7Hz), 7.98(1H, d, J=2.4Hz), 8.00(1H, dd, J=8.7, 2.4Hz), 8.31(1H, d, J=2.4Hz), 10.68(1H, s), 11.52(1H, brs).

Example 110: 5-Chloro-N-[3-fluoro-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 110).

[0486] Using 5-chlorosalicylic acid and 3-fluoro-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 62.0%.

¹H-NMR(DMSO-d₆): δ 7.04(1H, d, J=8.7Hz), 7.42(1H, d, J=8.4Hz), 7.48(1H, dd, J=9.0, 3.0Hz), 7.85(1H, d, J=2.4Hz), 7.94(1H, dd, J=11.4, 2.1Hz), 7.99(1H, s), 10.73(1H, s), 11.46(1H, s).

Example 111: 5-Bromo-N-[3-bromo-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 111).

[0487] Using 5-bromosalicylic acid and 3-bromo-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 73.3%.

¹H-NMR(DMSO-d₆): δ 6.99(1H, d, J=9.0Hz), 7.60(1H, dd, J=9.0, 2.4Hz), 7.72(1H, s), 7.97(1H, d, J=2.7Hz), 8.16(1H, s), 8.28(1H, s), 10.69(1H, s), 11.45(1H, s).

Example 112: 5-Chloro-N-[2-fluoro-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 112).

[0488] Using 5-chlorosalicylic acid and 2-fluoro-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 77.9%.

¹H-NMR(DMSO-d₆): δ 7.07(1H, d, J=9.0Hz), 7.52(1H, dd, J=9.0, 2.7Hz), 7.58-7.61(2H, m), 7.95(1H, d, J=2.7Hz), 8.71(1H, d, J=7.5Hz), 10.90(1H, s), 12.23(1H, s).

Example 113: 5-Chloro-N-[2-chloro-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 113).

[0489] Using 5-chlorosalicylic acid and 2-chloro-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 49.1%.

¹H-NMR(DMSO-d₆): δ 7.09(1H, d, J=9.0Hz), 7.53(1H, dd, J=9.0, 3.0Hz), 7.55(1H, dd, J=8.4, 2.7Hz), 7.83(1H, d, J=8.4Hz), 7.98(1H, d, J=3.0Hz), 8.88(1H, d, J=2.7Hz), 11.14(1H, s), 12.39(1H, s).

Example 114: 5-Bromo-N-[2-chloro-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 114).

[0490] Using 5-bromosalicylic acid and 2-chloro-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 34.2%.

¹H-NMR(DMSO-d₆): δ 7.04(1H, d, J=8.7Hz), 7.56(1H, ddd, J=8.1, 2.4, 1.2Hz), 7.64(1H, dd, J=8.7, 2.7Hz), 7.83(1H, dd, J=8.1, 1.2Hz), 8.11(1H, d, J=2.7Hz), 8.87(1H, d, J=2.4Hz), 11.12(1H, s), 12.42(1H, s).

EP 1 352 650 A1

Example 115: 5-Chloro-2-hydroxy-N-[4-nitro-3-(trifluoromethyl)phenyl]benzamide (Comopund No. 115).

[0491] Using 5-chlorosalicylic acid and 4-nitro-3-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 44.8%.

¹H-NMR(DMSO-d₆): δ 7.04(1H, d, J=9.0Hz), 7.49(1H, dd, J=9.0, 2.7Hz), 7.81(1H, d, J=2.7Hz), 8.23-8.24(2H, m), 8.43(1H, d, J=1.2Hz), 11.02(1H, s), 11.30(1H, br).

Example 116: 5-Chloro-2-hydroxy-N-[2-nitro-5-(trifluoromethyl)phenyl]benzamide (Comopund No. 116).

[0492] Using 5-chlorosalicylic acid and 2-nitro-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 8.1%.

¹H-NMR(DMSO-d₆): δ 7.08(1H, d, J=9.0Hz), 7.53(1H, dd, J=8.7, 2.7Hz), 7.73(1H, dd, J=8.4, 1.8Hz), 7.95(1H, d, J=3.0Hz), 8.36(1H, d, J=8.7Hz), 9.01(1H, d, J=1.8Hz), 12.04(1H, s), 12.20(1H, s).

Example 117: 5-Bromo-2-hydroxy-N-[4-nitro-3-(trifluoromethyl)phenyl]benzamide (Comopund No. 117).

[0493] Using 5-bromosalicylic acid and 4-nitro-3-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 49.7%.

¹H-NMR(DMSO-d₆): δ 6.99(1H, d, J=8.7Hz), 7.60(1H, dd, J=8.7, 2.4Hz), 7.92(1H, d, J=2.7Hz), 8.16(2H, s), 8.42(1H, s), 10.93(1H, s), 11.36(1H, s).

Example 118: 5-Chloro-2-hydroxy-N-[2-methyl-3-(trifluoromethyl)phenyl]benzamide (Comopund No. 118).

[0494] Using 5-chlorosalicylic acid and 2-methyl-3-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 14.5%.

¹H-NMR(DMSO-d₆): δ 2.36(3H, d, J=1.2Hz), 7.05(1H, d, J=8.7Hz), 7.46(1H, t, J=8.1Hz), 7.50(1H, dd, J=8.7, 2.7Hz), 7.60(1H, d, J=7.2Hz), 7.99(1H, d, J=7.2Hz), 8.00(1H, d, J=2.4Hz), 10.43(1H, s), 12.08(1H, s).

Example 119: 5-Chloro-2-hydroxy-N-[4-methyl-3-(trifluoromethyl)phenyl]benzamide (Comopund No. 119).

[0495] Using 5-chlorosalicylic acid and 4-methyl-3-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 80.2%.

¹H-NMR(DMSO-d₆): δ 7.01(1H, d, J=8.7Hz), 7.44(1H, d, J=8.4Hz), 7.47(1H, dd, J=9.0, 2.7Hz), 7.84(1H, dd, J=8.4, 2.1Hz), 7.92(1H, d, J=2.7Hz), 8.13(1H, d, J=2.1Hz), 10.65(1H, s), 11.68(1H, br).

Example 120: 5-Chloro-2-hydroxy-N-[2-methyl-5-(trifluoromethyl)phenyl]benzamide (Comopund No. 120).

[0496] Using 5-chlorosalicylic acid and 2-methyl-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 73.3%.

¹H-NMR(DMSO-d₆): 2.39(3H, s), 7.07(1H, d, J=8.7Hz), 7.44-7.54(3H, m), 7.99(1H, d, J=3.0Hz), 8.43(1H, s), 10.52(1H, s), 12.17(1H, brs).

Example 121: 5-Chloro-2-hydroxy-N-[4-methoxy-3-(trifluoromethyl)phenyl]benzamide (Comopund No. 121).

[0497] Using 5-chlorosalicylic acid and 4-methoxy-3-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 79.1%.

¹H-NMR(DMSO-d₆): δ 7.02(1H, d, J=9.0Hz), 7.30(1H, d, J=9.0Hz), 7.48(1H, dd, J=9.0, 3.0Hz), 7.92(1H, dd, J=9.0, 2.4Hz), 7.96(1H, d, J=2.7Hz), 8.04(1H, d, J=2.4Hz), 10.47(1H, s), 11.78(1H, s).

EP 1 352 650 A1

Example 122: 5-Bromo-2-hydroxy-N-[3-methoxy-5-(trifluoromethyl)phenyl]benzamide (Comopund No. 122).

[0498] Using 5-bromosalicylic acid and 3-methoxy-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 58.8%.

¹H-NMR(DMSO-d₆): δ 3.85(3H, s), 6.98(1H, d, J=8.7Hz), 7.03(1H, s), 7.57-7.61(2H, m), 7.77(1H, s), 8.00(1H, d, J=2.4Hz), 10.57(1H, s), 11.56(1H, s).

Example 123: 5-Bromo-2-hydroxy-N-(2-methoxy-5-(trifluoromethyl)phenyl)benzamide (Comopund No. 123).

[0499] Using 5-bromosalicylic acid and 2-methoxy-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 71.3%.

¹H-NMR(DMSO-d₆): δ 3.99(3H, s), 7.03(1H, d, J=9.0Hz), 7.30(1H, d, J=8.7Hz), 7.47-7.51(1H, m), 7.61(1H, dd, J=9.0, 2.4Hz), 8.10(1H, d, J=2.4Hz), 8.82(1H, d, J=2.1Hz), 11.03(1H, s), 12.19(1H, s).

Example 124: 5-Chloro-2-hydroxy-N-[2-methoxy-5-(trifluoromethyl)phenyl]-2benzamide (Comopund No. 124).

[0500] Using 5-chlorosalicylic acid and 2-methoxy-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 83.4%.

¹H-NMR(DMSO-d₆): δ 4.00(3H, s), 7.08(1H, d, J=9.0Hz), 7.30(1H, d, J=8.7Hz), 7.47-7.52(2H, m), 7.97(1H, d, J=2.7Hz), 8.83(1H, d, J=2.4Hz), 11.05(1H, s), 12.17(1H, s).

Example 125: 5-Chloro-2-hydroxy-N-[2-methylsulfanyl-5-(trifluoromethyl)phenyl]benzamide (Comopund No. 125).

[0501] Using 5-chlorosalicylic acid and 2-methyl-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 79.2%.

¹H-NMR(DMSO-d₆): δ 2.57(3H, s), 7.07(1H, d, J=8.7Hz), 7.52(1H, dd, J=8.7, 2.4Hz), 7.55(1H, dd, J=8.4, 1.5Hz), 7.63(1H, d, J=8.1Hz), 8.00(1H, d, J=2.4Hz), 8.48(1H, d, J=1.5Hz), 10.79(1H, s), 12.26(1H, s).

Example 126: 5-Bromo-2-hydroxy-N-[2-(1-pyrrolidinyl)-5-(trifluoromethyl)phenyl]benzamide (Comopund No. 126).

[0502] Using 5-bromosalicylic acid and 2-(1-pyrrolidinyl)-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound. Yield: 44.5%.

¹H-NMR(DMSO-d₆): δ 1.86-1.91(4H, m), 3.20-3.26(4H, m), 6.99(1H, d, J=8.7Hz), 7.07(1H, d, J=8.7Hz), 7.43(1H, dd, J=8.7, 2.1Hz), 7.62(1H, dd, J=8.7, 2.4Hz), 7.94(1H, d, J=2.1Hz), 8.17(1H, d, J=2.4Hz), 10.54(1H, s), 12.21(1H, s).

Example 127: 5-Bromo-2-hydroxy-N-[2-morpholino-5-(trifluoromethyl)phenyl]benzamide (Comopund No. 127).

[0503] Using 5-bromosalicylic acid and 2-morpholino-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound. Yield: 65.9%.

¹H-NMR(DMSO-d₆): δ 2.90(4H, dd, J=4.5, 4.2Hz), 3.84(4H, dd, J=4.8, 4.2Hz), 7.09(1H, d, J=8.4Hz), 7.48(2H, s), 7.61(1H, dd, J=8.4, 2.7Hz), 8.13(1H, d, J=2.7Hz), 8.90(1H, s), 11.21(1H, s), 12.04(1H, s).

Example 128: 5-Chloro-2-hydroxy-N-[4-(trifluoromethyl)phenyl]benzamide (Comopund No. 128).

[0504] Using 5-chlorosalicylic acid and 4-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 75.0%, white solid.

¹H-NMR(DMSO-d₆): δ 7.04(1H, d, J=9.0Hz), 7.48(1H, dd, J=8.7, 2.7Hz), 7.74(2H, d, J=8.7Hz), 7.90(1H, d, J=2.7Hz), 7.95(2H, d, J=9.0Hz), 10.65(1H, s), 11.59(1H, s).

Example 129: 5-Bromo-N-[2-chloro-4-(trifluoromethyl)phenyl]-2-hydroxybenzamide (Comopund No. 129).

[0505] Using 5-bromosalicylic acid and 2-chloro-4-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

EP 1 352 650 A1

Yield: 34.9%.

¹H-NMR(DMSO-d₆): δ 7.04(1H, d, J=8.7Hz), 7.64(1H, dd, J=8.7, 2.7Hz), 7.79(1H, dd, J=9.0, 2.1Hz), 7.99(1H, d, J=2.1Hz), 8.11(1H, d, J=2.4Hz), 8.73(1H, d, J=9.0Hz), 11.15(1H, s), 12.42(1H, s).

5 Example 130: 2-Acetoxy-5-chloro-N-[2-chloro-5-(trifluoromethyl)phenyl]benzamide (Comopund No. 130).

[0506] Using 5-chloro-N-[2-chloro-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide and acetyl chloride as the raw materials, the same operation as the example 96 gave the title compound.

Yield: 34.0%.

10 ¹H-NMR(CDCl₃): δ 2.39(3H, s), 7.16(1H, d, J=8.7Hz), 7.37(1H, ddd, J=8.7, 2.4, 0.6Hz), 7.51-7.56(2H, m), 7.97(1H, d, J=3.0Hz), 8.85(1H, s), 8.94(1H, d, J=1.8Hz).

Example 131: N-[2-Chloro-5-(trifluoromethyl)phenyl]-2-hydroxy-5-nitrobenzamide (Comopund No. 131).

15 **[0507]** Using 5-nitrosalicylic acid and 2-chloro-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 31.1%.

¹H-NMR(DMSO-d₆): δ 6.98(1H, d, J=9.3Hz), 7.52(1H, dd, J=8.4, 2.1Hz), 7.81(1H, d, J=8.4Hz), 8.21(1H, dd, J=9.0, 3.3Hz), 8.82(1H, d, J=3.0Hz), 8.93(1H, d, J=2.4Hz), 12.18(1H, s).

20

Example 132: N-[2-Chloro-5-(trifluoromethyl)phenyl]-2-hydroxy-5-methylbenzamide (Comopund No. 132).

[0508] Using 5-methylsalicylic acid and 2-chloro-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

25

Yield: 15.8%.

¹H-NMR(CDCl₃): δ 2.36(3H, s), 6.95(1H, d, J=8.1Hz), 7.26-7.31(2H, m), 7.37(1H, dd, J=8.4, 1.8Hz), 7.56(1H, d, J=8.4Hz), 8.65(1H, brs), 8.80(1H, d, J=1.8Hz), 11.33(1H, brs).

Example 133: N-[2-Chloro-5-(trifluoromethyl)phenyl]-2-hydroxy-5-methoxybenzamide (Comopund No. 133).

30

[0509] Using 5-methoxysalicylic acid and 2-chloro-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 56.4%.

35

¹H-NMR(DMSO-d₆): δ 3.77(3H, s), 6.91(1H, d, J=9.0Hz), 7.07(1H, dd, J=8.7, 3.0Hz), 7.20(1H, t, J=1.8Hz), 7.52-7.54(3H, m), 10.33(1H, s), 11.44(1H, s).

Example 134: N-[4-Chloro-3-(trifluoromethyl)phenyl]-2-hydroxy-5-methylbenzamide (Comopund No. 134).

40

[0510] Using 5-methylsalicylic acid and 4-chloro-3-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 70.4%.

¹H-NMR(DMSO-d₆): δ 2.29(3H, s), 6.9(1H, d, J=8.3Hz), 7.27(1H, ddd, J=8.3, 2.2, 0.6Hz), 7.71(1H, d, J=2.2Hz), 7.72(1H, d, J=8.5Hz), 8.02(1H, dd, J=8.5, 2.5Hz), 8.33(1H, d, J=2.5Hz), 10.64(1H, s), 11.25(1H, s).

45

Example 135: 2-Hydroxy-5-methyl-N-[4-methyl-3-(trifluoromethyl)phenyl]benzamide (Comopund No. 135).

[0511] Using 5-methylsalicylic acid and 4-methyl-3-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 63.7%.

50

¹H-NMR(DMSO-d₆): δ 2.29(3H, s), 2.42(3H, s), 6.89(1H, d, J=8.4Hz), 7.26(1H, ddd, J=8.4, 2.1, 0.6Hz), 7.44(1H, d, J=8.1Hz), 7.76(1H, d, J=2.1Hz), 7.86(1H, dd, J=8.4, 1.8Hz), 8.13(1H, d, J=2.1Hz), 10.50(1H, s), 11.42(1H, s).

Example 136: 2-Hydroxy-5-methyl-N-[2-methyl-5-(trifluoromethyl)phenyl]benzamide (Comopund No. 136).

55

[0512] Using 5-methylsalicylic acid and 2-methyl-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 14.2%, white solid.

¹H-NMR(DMSO-d₆): δ 2.29(3H, s), 2.38(3H, s), 6.94(1H, d, J=8.4Hz), 7.27(1H, ddd, J=8.4, 2.4, 0.6Hz), 7.44(1H, dd,

EP 1 352 650 A1

J=8.1, 1.5Hz), 7.52(1H, d, J=7.8Hz), 7.84(1H, d, J=2.4Hz), 8.46(1H, d, J=1.5Hz), 10.55(1H, s), 11.72(1H, s).

Example 137: 2-Hydroxy-N-[4-methoxy-3-(trifluoromethyl)phenyl]-5-methylbenzamide (Comopund No. 137).

5 [0513] Using 5-methylsalicylic acid and 4-methoxy-3-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 65.1%, slight yellow solid.

¹H-NMR(DMSO-d₆): δ 2.35(3H, s), 3.89(3H, s), 6.88(1H, d, J=8.4Hz), 7.26(1H, dd, J=8.1, 1.8Hz), 7.30(1H, d, J=8.4Hz), 7.77(1H, d, J=2.1Hz), 7.92(1H, dd, J=9.0, 2.1Hz), 8.04(1H, d, J=2.7Hz), 10.42(1H, s), 11.54(1H, s).

10 Example 138: 2-Hydroxy-N-[2-methoxy-5-(trifluoromethyl)phenyl]-5-methylbenzamide (Comopund No. 138).

[0514] Using 5-methylsalicylic acid and 2-methoxy-5-(trifluoromethyl)aniline as the raw materials, the same operation as the example 16 gave the title compound.

15 Yield: 77.9%.

¹H-NMR(CDCl₃): δ 2.35(3H, s), 4.02(3H, s), 6.93(1H, d, J=9.0Hz), 6.98(1H, d, J=8.4Hz), 7.25-7.28(2H, m), 7.36(1H, ddd, J=8.4, 2.1, 0.9Hz), 8.65(1H, brs), 8.73(1H, d, J=2.1Hz), 11.69(1H, s).

20 Example 139: 5-Bromo-2-hydroxy-N-phenylbenzamide (Comopund No. 139).

[0515] Using 5-bromosalicylic acid and aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 68.8%.

mp 229-230°C.

25 ¹H-NMR(DMSO-d₆): δ 6.96(1H, d, J=9.0Hz), 7.12-7.18(1H, m), 7.35-7.41(2H, m), 7.58(1H, dd, J=8.7, 2.7Hz), 7.67-7.71(2H, m), 8.08(1H, d, J=2.7Hz), 10.43(1H, s), 11.87(1H, s).

Example 140: 5-Bromo-N-(3-chlorophenyl)-2-hydroxybenzamide (Comopund No. 140).

30 [0516] Using 5-bromosalicylic acid and 3-chloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 63.1%.

mp 231-232°C.

35 ¹H-NMR(DMSO-d₆): δ 6.97(1H, d, J=8.7Hz), 7.19-7.22(1H, m), 7.38-7.43(1H, m), 7.57-7.63(2H, m), 7.91-7.92(1H, m), 8.01(1H, d, J=2.7Hz), 10.49(1H, s), 11.64(1H, s).

Example 141: 5-Bromo-N-(4-chlorophenyl)-2-hydroxybenzamide (Comopund No. 141).

[0517] This compound is a commercially available compound.

40 Supplier: Tokyo Kasei.

Catalog code number: B0897.

Example 142: 5-Chloro-N-(2,5-dichlorophenyl)-2-hydroxybenzamide (Comopund No. 142).

45 [0518] Using 5-chlorosalicylic acid and 2,5-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 10.8%.

¹H-NMR(DMSO-d₆): δ 7.08(1H, d, J=9.0Hz), 7.24-7.28(1H, m), 7.50-7.54(1H, m), 7.61(1H, dd, J=9.0, 3.0Hz), 7.97(1H, d, J=2.7Hz), 8.58(1H, d, J=2.4Hz), 11.02(1H, s), 12.35(1H, brs).

50 Example 143: 5-Bromo-N-(3,4-dichlorophenyl)-2-hydroxybenzamide (Comopund No. 143).

[0519] Using 5-bromosalicylic acid and 3,4-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

55 Yield: 58.2%.

mp 249-251°C.

¹H-NMR(DMSO-d₆): δ 6.97(1H, d, J=8.7Hz), 7.57-7.70(3H, m), 7.98(1H, d, J=2.7Hz), 8.10(1H, d, J=2.4Hz), 10.54(1H, s), 11.55(1H, s).

EP 1 352 650 A1

Example 144: 5-Bromo-N-(3,5-difluorophenyl)-2-hydroxybenzamide (Comopund No. 144).

[0520] Using 5-bromosalicylic acid and 3,5-difluoroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 36.3%.

mp 259-261°C

¹H-NMR(DMSO-d₆): δ 6.96-7.04(2H, m), 7.45-7.54(2H, m), 7.58(1H, dd, J=8.7, 2.7Hz), 7.94(1H, d, J=2.7Hz), 10.60(1H, s), 11.48(1H, s).

Example 145: 2-Acetoxy-N-(3,5-dichlorophenyl)benzamide (Comopund No. 172).

[0521] Using o-acetylsalicyloyl chloride and 3,5-dichloroaniline as the raw materials, the same operation as the example 2(1) gave the title compound.

Yield: 73.5%.

mp 167-168°C.

¹H-NMR(CDCl₃): δ 2.35(3H, s), 7.14-7.18(2H, m), 7.35-7.40(1H, m), 7.52-7.57(3H, m), 7.81(1H, dd, J=7.8, 1.8Hz), 8.05(1H, brs).

Example 146: N-(3,5-Dichlorophenyl)-2-hydroxybenzamide (Comopund No. 145).

[0522] Using 2-acetoxy-N-(3,5-dichlorophenyl)benzamide as the raw material, the same operation as the example 2(2) gave the title compound.

Yield: 60.3%.

mp 218-219°C.

¹H-NMR(DMSO-d₆): δ 6.95-7.02(2H, m), 7.35-7.36(1H, m), 7.42-7.47(1H, m), 7.83-7.87(3H, m), 10.54(1H, s), 11.35(1H, s).

Example 147: N-(3,5-Dichlorophenyl)-5-fluoro-2-hydroxybenzamide (Comopund No. 146).

[0523] Using 5-fluorosalicylic acid and 3,5-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 33.3%.

mp 258-260°C.

¹H-NMR(DMSO-d₆): δ 7.00-7.05(1H, m), 7.28-7.37(2H, m), 7.63(1H, dd, J=9.3, 3.3Hz), 7.84(2H, d, J=2.1Hz), 10.56(1H, s), 11.23(1H, s).

Example 148: 5-Chloro-N-(3,5-dichlorophenyl)-2-hydroxybenzamide (Comopund No. 147).

[0524] Using 5-chlorosalicylic acid and 3,5-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 41.2%.

¹H-NMR(DMSO-d₆): δ 7.03(1H, d, J=9.0Hz), 7.36-7.37(1H, m), 7.48(1H, dd, J=8.7, 2.7Hz), 7.83-7.84(3H, m), 10.56(1H, s), 11.44(1H, s).

Example 149: 5-Bromo-N-(3,5-dichlorophenyl)-2-hydroxybenzamide (Comopund No. 148).

[0525] Using 5-bromosalicylic acid and 3,5-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 61.6%.

mp 243-244°C.

¹H-NMR(DMSO-d₆): δ 6.98(1H, d, J=8.7Hz), 7.36-7.37(1H, m), 7.59(1H, dd, J=9.0, 2.4Hz), 7.83(2H, d, J=1.8Hz), 7.95(1H, d, J=2.4Hz), 10.56(1H, s), 11.46(1H, s).

Example 150: N-(3,5-Dichlorophenyl)-2-hydroxy-5-iodobenzamide (Comopund No. 149).

[0526] Using 5-iodosalicylic acid and 3,5-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 65.4%.

EP 1 352 650 A1

mp 244-245°C.

¹H-NMR(DMSO-d₆): δ 6.84(1H, d, J=9.0Hz), 7.35-7.37(1H, m), 7.72(1H, dd, J=9.0, 2.1Hz), 7.83(2H, d, J=1.8Hz), 8.09(1H, d, J=2.1Hz), 10.55(1H, s), 11.45(1H, s).

5 Example 151: 3,5-Dibromo-N-(3,5-dichlorophenyl)-2-hydroxybenzamide (Comopund No. 150).

[0527] Using 3,5-dibromosalicylic acid and 3,5-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 44.2%.

10 mp 181-182°C.

¹H-NMR(DMSO-d₆): δ 7.42-7.43(1H, m), 7.80(2H, d, J=1.8Hz), 8.03(1H, d, J=2.1Hz), 8.17(1H, d, J=2.1Hz), 10.82(1H, s).

Example 152: 4-Chloro-N-(3,5-dichlorophenyl)-2-hydroxybenzamide (Comopund No. 151).

15

[0528] Using 4-chlorosalicylic acid and 3,5-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 57.2%.

mp 255-256°C.

20 ¹H-NMR(DMSO-d₆): δ 7.03-7.06(2H, m), 7.34-7.36(1H, m), 7.82-7.85(3H, m), 10.51(1H, s), 11.70(1H, brs).

Example 153: N-(3,5-Dichlorophenyl)-2-hydroxy-5-nitrobenzamide (Comopund No. 152).

25 [0529] Using 5-nitrosalicylic acid and 3,5-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 83.1%.

mp 232-233.

30 ¹H-NMR(DMSO-d₆): δ 7.16(1H, d, J=9.6Hz), 7.37-7.39(1H, m), 7.84(1H, d, J=2.1Hz), 8.29(1H, dd, J=9.0, 3.0Hz), 8.65(1H, d, J=3.0Hz), 10.83(1H, s).

Example 154: N-(3,5-Dichlorophenyl)-2-hydroxy-5-methylbenzamide (Comopund No. 153).

[0530] Using 5-methylsalicylic acid and 3,5-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

35 Yield: 71.0%.

mp 216-217°C.

¹H-NMR(DMSO-d₆): δ 2.28(3H, s), 6.90(1H, d, J=8.4Hz), 7.26(1H, dd, J=8.7, 1.8Hz), 7.34-7.36(1H, m), 7.67(1H, d, J=1.5Hz), 7.85(2H, d, J=1.8Hz), 10.52(1H, s), 11.15(1H, s).

40 Example 155: N-(3,5-Dichlorophenyl)-2-hydroxy-5-methoxybenzamide (Comopund No. 154).

[0531] Using 5-methoxysalicylic acid and 3,5-dichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 29.8%.

45 mp 230-232°C.

¹H-NMR(DMSO-d₆): δ 3.76(3H, s), 6.95(1H, d, J=8.7Hz), 7.08(1H, dd, J=9.0, 3.0Hz), 7.35-7.36(1H, m), 7.40(1H, d, J=3.0Hz), 7.85(2H, d, J=1.5Hz), 10.55(1H, s), 10.95(1H, s).

Example 156: 5-Bromo-2-hydroxy-N-(3,4,5-trichlorophenyl)benzamide (Comopund No. 155).

50

[0532] Using 5-bromosalicylic acid and 3,4,5-trichloroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 78.6%.

mp 297-299°C.

55 ¹H-NMR(DMSO-d₆): δ 6.98(1H, d, J=9.0Hz), 7.58(1H, dd, J=8.4, 2.4Hz), 7.95(1H, d, J=2.4Hz), 8.03(1H, s), 10.58(1H, s), 11.49(1H, s).

EP 1 352 650 A1

Example 157: 5-Bromo-2-hydroxy-N-(3,5-dichloro-4-hydroxyphenyl)benzamide (Comopund No. 156).

[0533] Using 5-bromosalicylic acid and 3,5-dichloro-4-hydroxyaniline as the raw materials, the same operation as the example 16 gave the title compound. 22.5%.

¹H-NMR(DMSO-d₆): δ 6.96(1H, d, J=8.7Hz), 7.58(1H, dd, J=8.7, 2.4Hz), 7.76(2H, s), 8.01(1H, d, J=2.4Hz), 10.03(1H, s), 10.36(1H, s), 11.67(1H, brs).

Example 158: 5-Chloro-2-hydroxy-N-(2,3,4,5,6-pentafluorophenyl)benzamide (Comopund No. 157).

[0534] Using 5-chlorosalicylic acid and 2,3,4,5,6-pentafluoroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 58.6%.

¹H-NMR(DMSO-d₆): δ 7.07(1H, d, J=8.7Hz), 7.53(1H, dd, J=8.7, 2.7Hz), 7.91(1H, d, J=2.7Hz), 10.38(1H, brs), 11.74(1H, brs).

Example 159: 5-Bromo-N-(3,5-dinitrophenyl)-2-hydroxybenzamide (Comopund No. 158).

[0535] Using 5-bromosalicylic acid and 3,5-dinitroaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 32.2%.

mp 258-260°C.

¹H-NMR(DMSO-d₆): δ 6.98-7.02(1H, m), 7.59-7.63(1H, m), 7.96-7.97(1H, m), 8.56-8.58(1H, m), 9.03-9.05(2H, m), 11.04(1H, s), 11.39(1H, brs).

Example 160: N-[2,5-Bis[(1,1-dimethyl)ethyl]phenyl]-5-chloro-2-hydroxybenzamide (Comopund No. 159).

[0536] Using 5-chlorosalicylic acid and 2,5-bis[(1,1-dimethyl)ethyl]aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 75.7%.

¹H-NMR(DMSO-d₆): δ 1.27(9H, s), 1.33(9H, s), 7.04(1H, d, J=9.0Hz), 7.26(1H, dd, J=8.4, 2.1Hz), 7.35-7.38(2H, m), 7.49(1H, dd, J=8.7, 2.7Hz), 8.07(1H, d, J=2.4Hz), 10.22(1H, s), 12.38(1H, brs).

Example 161: 5-Chloro-N-[5-(1,1-dimethyl)ethyl-2-methoxyphenyl]-2-hydroxybenzamide (Comopund No. 160).

[0537] Using 5-chlorosalicylic acid and 5-[(1,1-dimethyl)ethyl]-2-methoxyaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 89.5%.

¹H-NMR(DMSO-d₆): δ 1.28(9H, s), 3.33(3H, s), 7.01(1H, d, J=8.7Hz), 7.05(1H, d, J=9.0Hz), 7.11(1H, dd, J=8.7, 2.4Hz), 7.47(1H, dd, J=9.0, 3.0Hz), 7.99(1H, d, J=3.0Hz), 8.49(1H, d, J=2.4Hz), 10.78(1H, s), 12.03(1H, s).

Example 162: 5-Bromo-N-(3,5-dimethylphenyl)-2-hydroxybenzamide (Comopund No. 161).

[0538] Using 5-bromosalicylic acid and 3,5-dimethylaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 58.1%.

mp 188-190°C.

¹H-NMR(DMSO-d₆): δ 2.28(6H, s), 6.80(1H, s), 6.96(1H, d, J=8.7Hz), 7.33(2H, s), 7.58(1H, dd, J=9.0, 2.4Hz), 8.10(1H, d, J=2.4Hz), 10.29(1H, s), 11.93(1H, brs).

Example 163: N-[3,5-Bis[(1,1-dimethyl)ethyl]phenyl]-5-chloro-2-hydroxybenzamide (Comopund No. 162).

[0539] Using 5-chlorosalicylic acid and 3,5-bis[(1,1-dimethyl)ethyl]aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 34.1%.

¹H-NMR(CDCl₃): δ 1.26(18H, s), 6.99(1H, d, J=8.7Hz), 7.29(1H, t, J=1.8Hz), 7.39(1H, dd, J=9.0, 2.4Hz), 7.41(2H, d, J=1.5Hz), 7.51(1H, d, J=2.1Hz), 7.81(1H, brs), 12.01(1H, s).

EP 1 352 650 A1

Example 164: N-(3,5-Bis[(1,1-dimethyl)ethyl]phenyl)-5-bromo-2-hydroxybenzamide (Comopund No. 163).

[0540] Using 5-bromosalicylic acid and 3,5-bis[(1,1-dimethyl)ethyl]aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 45.2%.

¹H-NMR(DMSO-d₆, δ): 1.30(18H, s), 6.95(1H, d, J=8.7Hz), 7.20(1H, t, J=1.5Hz), 7.56(2H, d, J=1.5Hz), 7.58(1H, dd, J=8.7, 2.4Hz), 8.12(1H, d, J=2.7Hz), 10.39(1H, s), 11.98(1H, s).

Example 165: 5-Chloro-2-hydroxy-(3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydronaphthalen-2-yl)benzamide (Comopund No. 164).

[0541] Using 5-chlorosalicylic acid and 2-amino-3,5,5,8,8-pentamethyl-5,6,7,8-tetrahydronaphthalene as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 77.5%.

¹H-NMR(DMSO-d₆, δ): 1.23(6H, s), 1.24(6H, s), 1.64(4H, s), 2.19(3H, s), 7.13(1H, d, J=9.0Hz), 7.20(1H, s), 7.49(1H, dd, J=8.7, 2.7Hz), 7.67(1H, s), 8.04(1H, d, J=2.7Hz), 10.23(1H, s), 12.26(1H, s).

Example 166: N-(Biphenyl-3-yl)-5-chloro-2-hydroxybenzamide (Comopund No. 165).

[0542] Using 5-chlorosalicylic acid and 3-aminobiphenyl as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 75.6%.

¹H-NMR(DMSO-d₆, δ): 7.04(1H, d, J=8.7Hz), 7.35-7.44(1H, m), 7.45-7.54(5H, m), 7.65-7.68(2H, m), 7.72(1H, dt, J=7.2, 2.1Hz), 7.99(1H, d, J=3.0Hz), 8.03(1H, m), 10.50(1H, s), 11.83(1H, brs).

Example 167: 5-Chloro-2-hydroxy-N-(4-methoxybiphenyl-3-yl)benzamide (Comopund No. 166).

[0543] Using 5-chlorosalicylic acid and 3-amino-4-methoxybiphenyl as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 37.0%.

¹H-NMR(DMSO-d₆, δ): 3.95(3H, s), 7.08(1H, d, J=8.7Hz), 7.20(1H, d, J=8.4Hz), 7.34(1H, t, J=7.2Hz), 7.40-7.50(4H, m), 7.62(1H, d, J=8.7Hz), 8.00(1H, d, J=3.0Hz), 8.77(1H, d, J=2.1Hz), 10.92(1H, s), 12.09(1H, s).

Example 168: 5-Bromo-N-(2,5-dimethoxyphenyl)-2-hydroxybenzamide (Comopund No. 167).

[0544] Using 5-bromosalicylic acid and 2,5-dimethoxyaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 39.7%.

¹H-NMR(DMSO-d₆, δ): 3.72(3H, s), 3.84(3H, s), 6.66(1H, ddd, J=9.0, 3.0, 0.6Hz), 6.99-7.03(2H, m), 7.58(1H, ddd, J=9.0, 2.7, 0.6Hz), 8.10(1H, dd, J=2.4, 0.6Hz), 8.12(1H, d, J=3.0Hz), 10.87(1H, s), 12.08(1H, s).

Example 169: 5-Bromo-N-(3,5-dimethoxyphenyl)-2-hydroxybenzamide (Comopund No. 168).

[0545] Using 5-bromosalicylic acid and 3,5-dimethoxyaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 40.3%.

mp 207-209°C.

¹H-NMR(DMSO-d₆, δ): 3.75(6H, s), 6.30-6.32(1H, m), 6.94-6.97(3H, m), 7.57(1H, dd, J=8.7, 2.4Hz), 8.04(1H, d, J=2.4Hz), 10.32(1H, s), 11.78(1H, s).

Example 170: 5-Chloro-N-(3-acetylphenyl)-2-hydroxybenzamide (Comopund No. 169).

[0546] Using 5-chlorosalicylic acid and 3-acetylaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 80.0%.

¹H-NMR(DMSO-d₆, δ): 2.60(3H, s), 7.03(1H, d, J=9.0Hz), 7.49(1H, dd, J=9.0, 3.0Hz), 7.54(1H, t, J=8.1Hz), 7.76(1H, dq, J=7.8, 0.9Hz), 7.96-8.00(2H, m), 8.30(1H, t, J=1.8Hz), 10.56(1H, s), 11.75(1H, s).

EP 1 352 650 A1

Example 171: 5-[[[(5-Bromo-2-hydroxy)benzoyl]amino]isophthalic acid dimethyl ester (Comopund No. 170).

[0547] Using 5-bromosalicylic acid and 5-aminoisophthalic acid dimethyl ester as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 74.1%.

mp 254-2561.

¹H-NMR(DMSO-d₆): δ 3.92(6H, s), 6.97(1H, d, J=9.0Hz), 7.60(1H, dd, J=9.0, 2.4Hz), 8.06(1H, d, J=2.4Hz), 8.24-8.25(1H, m), 8.62(2H, m), 10.71(1H, s), 11.57(1H, s).

Example 172: N-[4-[3-(2,3-Dichlorophenyl)thioureido]phenyl]-2-hydroxybenzamide (Comopund No. 171).

[0548] This compound is a commercially available compound.

Seller: Maybridge.

Catalog code number: RDR 01434

Example 173: N-(2,5-Bis[(1,1-dimethyl)ethyl]phenyl)-2-hydroxy-5-methylbenzamide (Comopund No. 173).

[0549] Using 5-methylsalicylic acid and 2,5-bis[(1,1-dimethyl)ethyl]aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 61.1%.

¹H-NMR(DMSO-d₆): δ 1.27(9H, s), 1.33(9H, s), 2.28(3H, s), 6.89(1H, d, J=8.1Hz), 7.24(1H, d, J=2.1Hz), 7.27(1H, d, J=2.1Hz), 7.32(1H, d, J=2.4Hz), 7.37(1H, d, J=8.4Hz), 7.88(1H, d, J=1.5Hz), 10.15(1H, s), 11.98(1H, brs).

Example 174: 2-Acetoxy-N-(3,5-bis[(1,1-dimethyl)ethyl]phenyl)-5-chlorobenzamide (Comopund No. 174).

[0550] Using N-(3,5-bis[(1,1-dimethyl)ethyl]phenyl)-5-chloro-2-hydroxybenzamide and acetyl chloride as the raw materials, the same operation as the example 96 gave the title compound.

Yield: 66.1%.

¹H-NMR(CDCl₃): δ 1.34(18H, s), 2.36(3H, s), 7.12(1H, d, J=8.4Hz), 7.25(1H, d, J=1.5Hz), 7.44(2H, d, J=1.2Hz), 7.47(1H, dd, J=8.7, 2.7Hz), 7.87(1H, d, J=2.4Hz), 7.98(1H, s).

Example 175: N-(3,5-Bis[(1,1-dimethyl)ethyl]phenyl)-2-hydroxy-5-nitrobenzamide (Comopund No. 175).

[0551] Using 5-nitrosalicylic acid and 3,5-bis[(1,1-dimethyl)ethyl]aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 46.7%.

¹H-NMR(CDCl₃): δ 1.37(18H, s), 7.13(1H, d, J=9.3Hz), 7.32(1H, t, J=1.8Hz), 7.46(2H, d, J=1.8Hz), 8.07(1H, s), 8.33(1H, dd, J=9.3, 2.1Hz), 8.59(1H, d, J=2.4Hz), 13.14(1H, s).

Example 176: N-(3,5-Bis[(1,1-dimethyl)ethyl]phenyl)-2-hydroxy-5-methylbenzamide (Comopund No. 176).

[0552] Using 5-methylsalicylic acid and 3,5-bis[(1,1-dimethyl)ethyl]aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 16.3%.

¹H-NMR(CDCl₃): δ 1.35(18H, s), 2.35(3H, s), 6.94(1H, d, J=8.4Hz), 7.23-7.28(2H, m), 7.31(1H, s), 7.42(1H, d, J=1.8Hz), 7.88(1H, s), 11.86(1H, s).

Example 177: N-(3,5-Bis[(1,1-dimethyl)ethyl]phenyl)-2-hydroxy-5-methoxybenzamide (Comopund No. 177).

[0553] Using 5-methoxysalicylic acid and 3,5-bis[(1,1-dimethyl)ethyl]aniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 12.7%.

¹H-NMR(DMSO-d₆): δ 3.56(3H, s), 7.01(1H, d, J=9.0Hz), 7.11(1H, dd, J=9.0, 3.0Hz), 7.52-7.56(2H, m), 7.83(1H, d, J=8.1Hz), 8.95(1H, d, J=1.5Hz), 11.29(1H, s), 11.63(1H, s).

Example 178: 2-Acetoxy-5-chloro-N-[5-(1,1-dimethyl)ethyl-2-methoxyphenyl]benzamide (Comopund No. 178).

[0554] Using 5-chloro-N-[5-(1,1-dimethyl)ethyl-2-methoxyphenyl]ethyl-2-hydroxybenzamide and acetyl chloride as

EP 1 352 650 A1

the raw materials, the same operation as the example 96 gave the title compound.

Yield: 87.5%.

¹H-NMR(CDCl₃): δ 1.35(9H, s), 2.37(3H, s), 3.91(3H, s), 6.86(1H, d, 8.7Hz), 7.12(1H, dd, J=8.7, 2.4Hz), 7.13(1H, d, J=9.0Hz), 7.47(1H, dd, J=9.0, 2.4Hz), 8.02(1H, d, J=2.7Hz), 8.66(1H, d, J=2.4Hz), 8.93(1H, s).

Example 179: N-[5(1,1-Dimethyl)ethyl-2-methoxyphenyl]-2-hydroxy-5-methylbenzamide (Comopund No. 178).

[0555] Using 5-methylsalicylic acid and 5-(1,1-dimethyl)ethyl-2-methoxyaniline as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 84.7%.

¹H-NMR(CDCl₃): δ 1.35(9H, s), 2.34(3H, s), 3.93(3H, s), 6.86(1H, d, J=8.7Hz), 6.93(1H, d, J=8.4Hz), 7.12(1H, dd, J=8.7, 2.4Hz), 7.24(1H, dd, J=8.4, 1.8Hz), 7.27(1H, brs), 8.48(1H, d, J=2.4Hz), 8.61(1H, brs), 11.95(1H, s).

Example 180: 5-Bromo-2-hydroxy-N-(thiazol-2-yl)benzamide (Comopund No. 180).

[0556] Using 5-bromosalicylic acid and 2-aminothiazole as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 12.0%.

mp 212°C(dec.).

¹H-NMR(DMSO-d₆): δ 6.94(1H, brd, J=8.0Hz), 7.25(1H, brd, J=3.2Hz), 7.56(2H, m), 8.05(1H, d, J=2.8Hz).

Example 181: 5-Bromo-N-{4-[(1,1-dimethyl)ethyl]thiazol-2-yl}-2-hydroxybenzamide (Comopund No. 186).

(1) 2-Amino-4-[(1,1-dimethyl)ethyl]thiazole.

[0557] A mixture of 1-bromo-3,3-dimethyl-2-butanone(5.03g, 28.1mmol), thiourea(2.35g, 30.9mmol) and ethanol (30mL) was refluxed for 1.5 hours. After cooling, the reaction mixture was poured into saturated aqueous sodium hydrogen carbonate and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel(hexane:ethyl acetate=2:1→ 1:1) to give the title compound(3.99g, 90.9%) as-a yellowish white powder.

¹H-NMR(CDCl₃): δ 1.26(9H, s), 4.96(2H, brs), 6.09(1H, s).

(2) 2-Acetoxy-5-bromo-N-{4-[(1,1-dimethyl)ethyl]thiazol-2-yl}benzamide.

[0558] Using 2-acetoxy-5-bromobenzoic acid and 2-amino-4-[(1,1-dimethyl)ethyl]thiazole as the raw materials, the same operation as the example 24 gave the title compound.

Yield: 59.4%.

¹H-NMR(CDCl₃): δ 1.31(9H, s), 2.44(3H, s), 6.60(1H, s), 7.13(1H, d, J=8.4Hz), 7.68(1H, dd, J=8.7, 2.4Hz), 8.17(1H, d, J=2.4Hz), 9.72(1H, brs).

(3) 5-Bromo-N-{4-[(1,1-dimethyl)ethyl]thiazol-2-yl}-2-hydroxybenzamide.

[0559] 2-Acetoxy-5-bromo-N-{4-[(1,1-dimethyl)ethyl]thiazol-2-yl}benzamide(100.1mg, 0.25mmol) was dissolved in tetrahydrofuran(3mL). 2 N sodium hydroxide(0.2ml) was added, and the mixture was stirred at room temperature for 20 minutes. The reaction mixture was poured into diluted hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was crystallized(isopropyl ether/n-hexane) to give the title compound(70.1mg, 78.9%) as a light gray solid.

¹H-NMR(DMSO-d₆): δ 1.30(9H, s), 6.80(1H, brs), 6.95(1H, brs), 7.57(1H, brs), 8.06(1H, d, J=2.4Hz), 11.82(1H, brs), 13.27(1H, brs).

Example 182: 5-Bromo-N-{5-bromo-4-[(1,1-dimethyl)ethyl]thiazol-2-yl}-2-hydroxybenzamide (Comopund No. 181).

(1) 2-Acetoxy-5-bromo-N-{5-bromo-4-[(1,1-dimethyl)ethyl]thiazol-2-yl}benzamide.

[0560] 2-Acetoxy-5-bromo-N-{4-[(1,1-dimethyl)ethyl]imidazol-2-yl}benzamide(0.20g, 0.50mmol) was dissolved in acetonitrile(10mL). N-Bromosuccinimide(97.9mg, 0.55mmol) was added, and the mixture was stirred at room temperature.

EP 1 352 650 A1

ture for 1 hour. The reaction mixture was concentrated under reduced pressure, and the obtained residue was purified by chromatography on silica gel(n-hexane:ethyl acetate=3:1) to give the title compound as a crude product.

(2) 5-Bromo-N-(5-bromo-4-((1,1-dimethyl)ethyl)thiazol-2-yl)-2-hydroxybenzamide.

[0561] Using 2-acetoxy-5-bromo-N-(5-bromo-4-((1,1-dimethyl)ethyl)thiazol-2-yl)benzamide as the raw material, the same operation as the example 2(2) gave the title compound.

Yield: 90.9%(2steps).

mp 212°C (dec.).

¹H-NMR(DMSO-d₆): δ 1.42(9H, s), 6.99(1H, d, J=8.7Hz), 7.61(1H, dd, J=8.7, 2.7Hz), 8.02(1H, d, J=2.4Hz), 11.79(1H, brs), 12.00(1H, brs).

Example 183: 5-Bromo-N-[5-bromo-4-(trifluoromethyl)thiazol-2-yl]-2-hydroxybenzamide (Comopund No. 182).

[0562] Using 5-bromosalicylic acid and 2-amino-5-bromo-4-(trifluoromethyl)thiazole as the raw materials, the same operation as the example 16 gave the title compound. (2-Amino-5-bromo-4-(trifluoromethyl)thiazole: refer to J. Heterocycl. Chem., 1991, 28, 1017.)

Yield: 22.4%.

mp 215°C (dec.).

¹H-NMR(DMSO-d₆): δ 7.00(1H, d, J=8.8Hz), 7.61(1H, dd, J=8.8, 2.8Hz), 7.97(1H, d, J=2.4Hz).

Example 184: 5-Chloro-N-(5-cyano-4-((1,1-dimethyl)ethyl)thiazol-2-yl)-2-hydroxybenzamide.

(1) α-Bromo-pivaloylacetoneitrile.

[0563] Pivaloylacetoneitrile(1.00g, 7.99mmol) was dissolved in carbon tetrachloride(15mL). N-Bromosuccinimide (1.42g, 7.99mmol) was added, and the mixture was refluxed for 15 minutes. After cooling, the insoluble matter was filtered off, and the residue obtained by evaporation of the filtrate under reduced pressure was purified by chromatography on silica gel(n-hexane:ethyl acetate=4:1) to give the title compound(1.43g, 87.9%) as a yellowish brown oil.

¹H-NMR(CDCl₃): δ 1.33(9H, s), 5.10(1H, s).

(2) 2-Amino-5-cyano-4-((1,1-dimethyl)ethyl)thiazole

[0564] Using α-bromo-pivaloylacetoneitrile and thiourea as the raw materials, the same operation as the example 181 (1) gave the title compound.

Yield: 66.3%.

¹H-NMR(CDCl₃): δ 1.41(9H, s), 5.32(2H, s).

(3) 5-Chloro-N-(5-cyano-4-((1,1-dimethyl)ethyl)thiazol-2-yl)-2-hydroxybenzamide.

[0565] Using 5-chlorosalicylic acid and 2-amino-5-cyano-4-((1,1-dimethyl)ethyl)thiazole as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 63.4%.

¹H-NMR(DMSO-d₆): δ 1.43(9H, s), 7.06(1H, d, J=8.7Hz), 7.51(1H, dd, J=8.7, 3.0Hz), 7.85(1H, d, J=2.7Hz), 12.31(2H, br).

Example 185: 5-Bromo-N-(5-cyano-4-((1,1-dimethyl)ethyl)thiazol-2-yl)-2-hydroxybenzamide (Comopund No. 184).

[0566] Using 5-bromosalicylic acid and 2-amino-5-cyano-4-((1,1-dimethyl)ethyl)thiazole as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 61.3%.

¹H-NMR(DMSO-d₆): δ 1.43(9H, s), 7.00(1H, d, J=8.7Hz), 7.62(1H, dd, J=8.7, 2.7Hz), 7.97(1H, d, J=2.7Hz), 11.75(1H, br), 12.43(1H, br).

Example 186: 5-Bromo-2-hydroxy-N-(5-methylthiazol-2-yl)benzamide (Comopund No. 185).

[0567] Using 5-bromosalicylic acid and 2-amino-5-methylthiazole as the raw materials, the same operation as the example 16 gave the title compound.

EP 1 352 650 A1

Yield: 12.9%.

¹H-NMR(DMSO-d₆): δ 2.33(3H, s), 6.91(1H, d, J=7.6Hz), 7.26(1H, s), 7.54(1H, d, J=9.6Hz), 8.03(1H, d, J=2.8Hz).

Example 187: 5-Bromo-N-(4,5-dimethylthiazol-2-yl)-2-hydroxybenzamide (Comopund No. 187).

[0568] Using 5-bromosalicylic acid and 2-amino-4,5-dimethylthiazole as the raw materials, the same operation as the example 16 gave the title compound.

Yield: 14.4%.

¹H-NMR(DMSO-d₆): δ 2.18(3H, s), 2.22(3H, s), 6.89(1H, d, J=8.8Hz), 7.51(1H, d, J=6.8Hz), 8.02(1H, d, J=2.8Hz), 13.23(1H, brs).

Example 188: 5-Bromo-N-(5-methyl-4-phenylthiazol-2-yl)-2-hydroxybenzamide (Comopund No. 188).

[0569] Using 5-bromosalicylic acid and 2-amino-5-methyl-4-phenylthiazole as the raw materials, the same operation as the example 16 gave the title compound. (2-Amino-5-methyl-4-phenylthiazole: refer to Yakugaku Zasshi, 1961, 81, 1456.)

Yield: 27.7%.

mp 243-244°C.

¹H-NMR(CD₃OD): δ 2.47(3H, s), 6.92(1H, d, J=8.7Hz), 7.36-7.41(1H, m), 7.44-7.50(2H, m), 7.53(1H, dd, J=9.0, 2.7Hz), 7.57-7.61(2H, m), 8.16(1H, d, J=2.7Hz).

Example 189: 5-Bromo-N-(4-methyl-5-(4-fluorophenyl)thiazol-2-yl)-2-hydroxybenzamide (Comopund No. 189).

[0570] Using (4-fluorophenyl)acetone as the raw material, the same operation as the examples 184(1)-(3) gave the title compound.

Yield: 28.8%(3steps).

(1) α-Bromo-(4-fluorophenyl)acetone.

[0571] ¹H-NMR(CDCl₃): δ 2.33(3H, s), 5.41(1H, s), 7.07(2H, t, J=8.7Hz), 7.43(2H, dd, J=8.7, 5.1Hz).

(2) 2-Amino-4-methyl-5-(4-fluorophenyl)thiazole.

[0572] ¹H-NMR(CDCl₃): δ 2.27(3H, s), 4.88(2H, s), 7.07(2H, t, J=8.7Hz), 7.32(2H, dd, J=8.7, 5.4Hz).

(3) 5-Bromo-N-[4-methyl-5-(4-fluorophenyl)thiazol-2-yl]-2-hydroxybenzamide.

[0573] ¹H-NMR(DMSO-d₆): δ 2.36(3H, s), 6.95(1H, d, J=8.4Hz), 7.33(2H, t, J=8.7Hz), 7.52-7.59(3H, m), 8.06(1H, d, J=3.0Hz), 12.01-13.65(2H, br).

Example 190: 5-Bromo-N-[4-methyl-5-[3-(trifluoromethyl)phenyl]thiazol-2-yl]-2-hydroxybenzamide (Comopund No. 190).

[0574] Using 3-(trifluoromethyl)phenylacetone as the raw material, the same operation as the examples 184(1)-(3) gave the title compound.

Yield: 39.8%(3steps).

(1) α-Bromo-3-(trifluoromethyl)phenylacetone.

[0575] ¹H-NMR(CDCl₃): δ 2.38(3H, s), 5.43(1H, s), 7.52(1H, t, J=7.8Hz), 7.61-7.66(2H, m), 7.69-7.70(1H, m).

(2) 2-Amino-4-methyl-5-[3-(trifluoromethyl)phenyl]thiazole.

[0576] ¹H-NMR(CDCl₃): δ 2.32(3H, s), 4.95(2H, s), 7.46-7.56(3H, m), 7.59-7.61(1H, m).

(3) 5-Bromo-N-[4-methyl-5-[3-(trifluoromethyl)phenyl]thiazol-2-yl]-2-hydroxybenzamide.

[0577] ¹H-NMR(DMSO-d₆): δ 2.40(3H, s), 6.97(1H, d, J=8.7Hz), 7.59(1H, dd, J=8.7, 2.4Hz), 7.71-7.84(4H, m), 12.01-13.65(2H, br).

EP 1 352 650 A1

m), 8.06(1H, d, J=2.4Hz), 12.09(1H, br), 12.91-13.63(1H, br).

Example 191: 5-Bromo-N-(4-[(1,1-dimethyl)ethyl]-5-ethylthiazol-2-yl)-2-hydroxybenzamide (Comopund No. 191).

- 5 [0578] Using 2,2-dimethyl-3-hexanone as the raw material, the same operation as the examples 184(1)-(3) gave the title compound.
Yield: 17.0%(3steps).

(2) 2-Amino-4-[(1,1-dimethyl)ethyl]-5-ethylthiazole.

10

[0579] ¹H-NMR(CDCl₃): δ 1.21(3H, t, J=7.5Hz), 1.32(9H, s), 2.79(2H, q, J=7.5Hz), 4.63(2H, brs).

(3) 5-Bromo-N-(4-[(1,1-dimethyl)ethyl]-5-ethylthiazol-2-yl)-2-hydroxybenzamide.

- 15 [0580] ¹H-NMR(CDCl₃): δ 1.32(3H, t, J=7.5Hz), 1.41(9H, s), 2.88(2H, q, J=7.5Hz), 6.84(1H, d, J=9.0Hz), 7.44(1H, dd, J=8.7, 2.4Hz), 8.05(1H, d, J=2.7Hz), 11.46(2H, br).

Example 192: 5-Bromo-N-(4-ethyl-5-phenylthiazol-2-yl)-2-hydroxybenzamide (Comopund No. 192).

- 20 [0581] Using 5-bromosalicylic acid and 2-amino-4-ethyl-5-phenylthiazole as the raw materials, the same operation as the example 16 gave the title compound.
Yield: 17.4%.
mp 224-225°C.
¹H-NMR(DMSO-d₆): δ 1.24(3H, t, J=7.6Hz), 2.70(2H, q, J=7.6Hz), 6.95(1H, brd, J=7.6Hz), 7.39-7.42(1H, m), 7.45-7.51
25 (4H, m), 7.56(1H, brd, J=8.0Hz), 8.06(1H, d, J=2.8Hz), 11.98(1H, brs).

Example 193: 5-Bromo-N-(4-ethyl-5-isopropylthiazol-2-yl)-2-hydroxybenzamide (Comopund No. 193).

- [0582] Using benzyl isopropyl ketone as the raw material, the same operation as the examples 184(1)-(3) gave the
30 title compound.
Yield: 4.4%(3steps).

(2) 2-Amino-4-ethyl-5-isopropylthiazole.

- 35 [0583] ¹H-NMR(CDCl₃): δ 1.23(6H, d, J=6.6Hz), 3.05(1H, m), 4.94(2H, s), 7.28-7.41(5H, m).

(3) 5-Bromo-N-(4-ethyl-5-isopropylthiazol-2-yl)-2-hydroxybenzamide.

- [0584] ¹H-NMR(DMSO-d₆): δ 1.26(6H, d, J=6.0Hz), 3.15(1H, m), 6.98(1H, brs), 7.43-7.53(5H, m), 7.59(1H, brs),
40 8.08(1H, d, J=2.7Hz), 11.90(1H, brs), 13.33(1H, brs).

Example 194: 5-Bromo-N-(4-butyl-5-phenylthiazol-2-yl)-2-hydroxybenzamide (Comopund No. 194).

- [0585] Using 1-phenyl-2-hexanone as the raw material, the same operation as the examples 184(1)-(3) gave the
45 title compound.
Yield: 52.6%(3steps).

(1) α-Bromo-1-phenyl-2-hexanone.

- 50 [0586] ¹H-NMR(CDCl₃): δ 0.85(3H, t, J=7.2Hz), 1.19-1.32(2H, m), 1.50-1.60(2H, m), 2.59(2H, td, J=7.5, 3.9Hz), 5.44
(1H, s), 7.34-7.45(5H, m).

(2) 2-Amino-4-butyl-5-phenylthiazole.

- 55 [0587] ¹H-NMR(CDCl₃, δ): 0.89(3H, t, J=7.5Hz), 1.28-1.41(2H, m), 1.61-1.71(2H, m), 2.56-2.61(2H, m), 4.87(2H, s), 7.25-7.40(5H, m).

EP 1 352 650 A1

(3) 5-Bromo-N-(4-butyl-5-phenylthiazol-2-yl)-2-hydroxybenzamide.

[0588] ¹H-NMR(DMSO-d₆): δ 0.85(3H, t, J=7.2Hz), 1.23-1.35(2H, m), 1.59-1.69(2H, m), 2.70(2H, t, J=7.2Hz), 6.96(1H, d, J=6.9Hz), 7.39-7.59(6H, m), 8.07(1H, d, J=2.4Hz), 11.93(1H, br), 13.18-13.59(1H, br).

Example 195: 5-Chloro-N-(4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl)-2-hydroxybenzamide (Comopund No. 195).

(1) α-Bromo-dipivaloylmethane.

[0589] Dipivaloylmethane(1.00g, 5.42mmol) was dissolved in carbon tetrachloride (10mL). N-Bromosuccinimide (965.8mg, 5.42mmol) was added, and the mixture was refluxed for 2 hours. After cooling, the insoluble matter was filtered off, and the filtrate was evaporated under reduced pressure to give the title compound(1.42g, quant.) as a white crystal.

¹H-NMR(CDCl₃, δ): 1.27(18H, s), 5.67(1H, s).

(2) 2-Amino-4-[(1,1-dimethyl)ethyl]-5-[(2,2-dimethyl)propionyl]thiazole.

[0590] A mixture of α-bromo-dipivaloylmethane(1.42g), thiourea(451.8mg) and ethanol(15m was refluxed for 2 hours. After cooling, the reaction mixture was poured into saturated aqueous sodium hydrogen carbonate and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was crystallized(dichloromethane/hexane) to give the title compound(1.23g, 94.5%) as a white crystal.

¹H-NMR (CDCl₃, δ): 1.26(9H, s), 1.29(9H, s), 5.03(2H, s).

(3) 5-Chloro-N-(4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl)-2-hydroxybenzamide.

[0591] A mixture of 5-bromosalicylic acid (0.20g, 0.92mmol), 2-amino-4-[(1,1-dimethyl)ethyl]-5-[(2,2-dimethyl)propionyl]thiazole(221.5mg, 0.92mmol), phosphorus trichloride (40 μl, 0.46mmol) and chlorobenzene(5mL) was refluxed for 3 hours. The residue obtained by concentration of the reaction mixture under reduced pressure was purified by chromatography on silica gel(n-hexane:ethyl acetate=2:1) to give the title compound(96.2mg, 23.8%) as a white powder.

¹H-NMR(CDCl₃, δ): 1.33(9H, s), 1.35(9H, s), 6.94(1H, d, J=8.7Hz), 7.55(1H, dd, J=8.7, 2.1Hz), 7.85(1H, d, J=2.1Hz), 10.51(2H, br).

Example 196: 5-Bromo-N-(4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl)-2-hydroxybenzamide (Comopund No. 196).

[0592] Using 5-bromosalicylic acid and 2-amino-4-[(1,1-dimethyl)ethyl]-5-[(2,2-dimethyl)propionyl]thiazole as the raw material, the same operation as the example 195(3) gave the title compound.

Yield: 23.8%.

¹H-NMR(CDCl₃, δ): 1.33(9H, s), 1.35(9H, s), 6.94(1H, d, J=8.7Hz), 7.55(1H, dd, J=8.7, 2.1Hz), 7.85(1H, d, J=2.1Hz), 10.51(2H, br).

Example 197: 2-(5-Bromo-2-hydroxybenzoyl)amino-4-[(1,1-dimethyl)ethyl]thiazole-5-carboxylic acid ethyl ester (Comopund No. 197).

[0593] Using pivaloyl acetic acid ethyl ester as the raw material, the same operation as the examples 195(1)-(3) gave the title compound.

Yield: 45.7%(3steps).

(1) α-Bromo-pivaloyl acetic acid ethyl ester.

[0594] ¹H-NMR(CDCl₃, δ): 1.28(9H, s), 1.29(3H, t, J=7.2Hz), 4.26(2H, q, J=7.2Hz), 5.24(1H, s).

(2) 2-Amino-4-[(1,1-dimethyl)ethyl]thiazole-5-carboxylic acid ethyl ester.

[0595] ¹H-NMR(CDCl₃, δ): 1.32(3H, t, J=7.2Hz), 1.43(9H, s), 4.24(2H, q, J=7.2Hz), 5.18(2H, s).

EP 1 352 650 A1

(3) 2-(5-Bromo-2-hydroxybenzoyl)amino-4-[(1,1-dimethyl)ethyl]thiazole-5-carboxylic acid ethyl ester.

[0596] ¹H-NMR(DMSO-d₆): δ 1.30(3H, t, J=7.2Hz), 1.44(9H, s), 4.27(2H, q, J=6.9Hz), 7.00(1H, d, J=8.7Hz), 7.63(1H, dd, J=8.7, 2.7Hz), 8.02(1H, d, J=2.4Hz), 11.80(1H, br), 12.12(1H, br).

Example 198: 5-Bromo-N-[4-(1,1-dimethyl)ethyl-5-piperidinethiazol-2-yl]-2-hydroxybenzamide (Comopund No. 198)

(1) 2-Amino-5-bromo-4-[(1,1-dimethyl)ethyl]thiazole.

[0597] 2-Amino-4-[(1,1-dimethyl)ethyl]thiazole(0.87g, 5.6mmol) was dissolved in carbon tetrachloride(9mL). N-Bromosuccinimide(1.00g, 5.6mmol) was added, and the mixture was stirred at room temperature for 1 hour. Hexane was added to the reaction mixture, the insoluble matter was filtered off, and the residue obtained by evaporation of the filtrate under reduced pressure was purified by chromatography on silica gel(hexane:ethyl acetate=2:1) to give the title compound(1.23g, 93.7%) as a yellowish gray powder.

¹H-NMR(CDCl₃): δ 1.39(9H, s), 4.81(2H, brs).

(2) 2-Amino-4-[(1,1-dimethyl)ethyl]-5-piperidinethiazole.

[0598] A mixture of 2-amino-5-bromo-4-[(1,1-dimethyl)ethyl]thiazole(0.10g, 0.42mmol), piperidine(0.1mL potassium carbonate(0.20g) and acetonitrile(4mL) was refluxed for 3 hours. The reaction mixture was poured into water and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel(n-hexane:ethyl acetate=2:1) to give the title compound(80.7mg, 79.3%) as a yellow crystal.

¹H-NMR(CDCl₃): δ 1.32(9H, s), 1.64(4H, t, J=5.7Hz), 1.71-1.77(2H, m), 2.35(2H, brs), 2.99(2H, brs), 4.68(2H, s).

(3) 2-Acetoxy-5-bromo-N-[4-(1,1-dimethyl)ethyl-5-piperidinethiazol-2-yl]benzamide.

[0599] Under argon atmosphere, phosphorus oxychloride(46 μl, 0.50mmol) was added to a mixture of 2-acetoxy-5-bromobenzoic acid(J. Med. Cehm. 31, 861-874 1996)(90.3mg, 0.35mmol), the thiazole(80.7mg, 0.34mmol), pyridine(0.1mL) and THF(3mL), and the mixture was stirred at room temperature for 2 hours. The reaction mixture was poured into 2N hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel(n-hexane:ethyl acetate=3:1) to give the title compound(84.3mg) as a crude product.

(4) 5-Bromo-N-[4-(1,1-dimethyl)ethyl-5-piperidinethiazol-2-yl]-2-hydroxybenzamide.

[0600] 2-Acetoxy-5-bromo-N-[4-(1,1-dimethyl)ethyl-5-piperidinethiazol-2-yl]benzamide(crude product, 84.3mg) was dissolved in ethanol(3mL). 2 N aqueous sodium hydroxide (0.1mL) was added, and the mixture was stirred at room temperature for 1 hour. The reaction mixture was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel(n-hexane:ethyl acetate=4:1) to give the title compound(54.1mg, 36.3%; 2steps) as a white powder.

¹H-NMR(CDCl₃): δ 1.41(9H, s), 1.56(2H, brs), 1.67-1.74(4H, m), 2.79(4H, brs), 6.85(1H, d, J=9.0Hz), 7.45(1H, dd, J=9.0, 2.4Hz), 8.06(1H, d, J=2.4Hz), 11.70(2H, br).

Example 199: 5-Bromo-N-[4-(1,1-dimethyl)ethyl-5-morpholinethiazol-2-yl]-2-hydroxybenzamide (Comopund No. 199).

[0601] Using morpholine as the raw material, the same operation as the examples 198(2)-(4) gave the title compound. Yield: 17.1%.

(2) 2-Amino-4-[(1,1-dimethyl)ethyl]-5-morpholinethiazole.

[0602] ¹H-NMR(CDCl₃): δ 1.33(9H, s), 2.76(4H, brs), 3.79(4H, brs), 4.66(2H, s).

(3) 2-Acetoxy-5-bromo-N-[4-(1,1-dimethyl)ethyl-5-morpholinethiazol-2-yl]benzamide.

[0603] The product was used for the next reaction as a crude product.

EP 1 352 650 A1

(4) 5-Bromo-N-[4-(1,1-dimethyl)ethyl-5-morpholinothiazol-2-yl]-2-hydroxybenzamide.

[0604] ¹H-NMR(CDCl₃): δ 1.24(9H, s), 2.89(4H, dd, J=4.8, 4.2Hz), 3.83(4H, dd, J=4.5, 4.2Hz), 6.89(1H, d, J=9.0Hz), 7.49(1H, dd, J=9.0, 2.4Hz), 7.98(1H, d, J=2.1Hz), 11.20(2H, br).

Example 200: 5-Bromo-N-[4-(1,1-dimethyl)ethyl-5-(4-methylpiperazin-1-yl)thiazol-2-yl]-2-hydroxybenzamide (Comopund No. 200).

[0605] Using 4-methylpiperazine as the raw material, the same operation as the examples 198(2)-(4) gave the title compound.
Yield: 6.9%.

(2) 2-Amino-4-(1,1-dimethyl)ethyl-5-(4-methylpiperazin-1-yl)thiazole.

[0606] ¹H-NMR(DMSO-d₆): δ 1.25(9H, s), 2.12(2H, brs), 2.19(3H, s), 2.57(2H, brs), 2.72(4H, brs), 6.51(2H, s).

(3) 2-Acetoxy-N-[4-(1,1-dimethyl)ethyl-5-(4-methylpiperazin-1-yl)thiazol-2-yl]benzamide.

[0607] The product was used for the next reaction as a crude product.

(4) 5-Bromo-N-[4-(1,1-dimethyl)ethyl-5-(4-methylpiperazin-1-yl)thiazol-2-yl]-2-hydroxybenzamide.

[0608] ¹H-NMR(CD₃OD): δ 1.41(9H, s), 2.55(3H, s), 2.87(4H, brs), 3.03(4H, brs), 6.88(1H, d, J=8.7Hz), 7.49(1H, dd, J=8.7, 2.7Hz), 8.11(1H, d, J=2.7Hz).

Example 201: 5-Bromo-N-[4-(1,1-dimethyl)ethyl-5-(4-phenylpiperazin-1-yl)thiazol-2-yl]-2-hydroxybenzamide (Comopund No. 201).

[0609] Using 4-phenylpiperazine as the raw material, the same operation as the examples 198(2)-(4) gave the title compound.
Yield: 6.9%.

(2) 2-Amino-4-(1,1-dimethyl)ethyl-5-(4-phenylpiperazin-1-yl)thiazole.

[0610] ¹H-NMR(CDCl₃): δ 1.34(9H, s), 2.80(2H, brs), 3.03(4H, brs), 3.55(2H, brs), 4.69(2H, s), 6.88(1H, tt, J=7.2, 1.2Hz), 6.95(2H, dd, J=9.0, 1.2Hz), 7.28(2H, dd, J=8.7, 7.2Hz).

(3) 2-Acetoxy-5-bromo-N-[4-(1,1-dimethyl)ethyl-5-(4-phenylpiperazin-1-yl)thiazol-2-yl]benzamide.

[0611] The product was used for the next reaction as a crude product.

(4) 5-Bromo-N-[4-(1,1-dimethyl)ethyl-5-(4-phenylpiperazin-1-yl)thiazol-2-yl]-2-hydroxybenzamide.

[0612] ¹H-NMR(DMSO-d₆): δ 1.39(9H, s), 2.97(4H, s), 3.30(4H, s), 6.82(1H, t, J=7.5Hz), 6.97(2H, brs), 6.99(2H, t, J=7.5Hz), 7.58(1H, brs), 8.05(1H, d, J=2.4Hz), 11.69(1H, brs), 11.82(1H, brs).

Example 202: 5-Bromo-N-[4-phenylthiazol-2-yl]-2-hydroxybenzamide (Comopund No. 202).

[0613] Using 5-bromosalicylic acid and 2-amino-4-phenylthiazole as the raw materials, the same operation as the example 195(3) gave the title compound.
Yield: 16.0%.

mp 239°C(dec.).

¹H-NMR(DMSO-d₆): δ 7.02(1H, d, J=8.4Hz), 7.34(1H, t, J=7.6Hz), 7.44(2H, t, J=7.6Hz), 7.62(1H, dd, J=8.4, 2.8Hz), 7.67(1H, s), 7.92(2H, d, J=7.2Hz), 8.08(1H, d, J=2.8Hz), 11.88(1H, brs), 12.05(1H, brs).

EP 1 352 650 A1

Example 203: {2-[(5-Bromo-2-hydroxybenzoyl)amino]-4-phenylthiazol-5-yl}acetic acid (Comopund No. 203).

(1) {2-[(5-Bromo-2-hydroxybenzoyl)amino]-4-phenylthiazol-5-yl}acetic acid methyl ester.

[0614] Using 5-bromosalicylic acid and 2-amino-4-phenylthiazole-5-acetic acid methyl ester as the raw materials, the same operation as the example 195(3) gave the title compound.

Yield: 32.1%.

mp 288.5-229.5°C.

¹H-NMR(DMSO-d₆): δ 3.66(3H, s), 3.95(2H, s), 6.99(1H, d, J=8.0Hz), 7.42(1H, d, J=6.0Hz), 7.48(2H, brt, J=7.6Hz), 7.56-7.61(3H, m), 8.07(1H, d, J=2.4Hz), 11.85(1H, brs), 11.98(1H, brs).

(2) {2-[(5-Bromo-2-hydroxybenzoyl)amino]-4-phenylthiazol-5-yl}acetic acid.

[0615] {2-[(5-Bromo-2-hydroxybenzoyl)amino]-4-phenylthiazol-5-yl}acetic acid methyl ester(75mg, 0.17mmol) was dissolved in methanol(5mL). 2 N sodium hydroxide(0.5mL, 1mmol) was added, and the mixture was stirred at room temperature for 12 hours. The reaction mixture was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine one after another, dried over anhydrous sodium sulfate, the solvent was evaporated under reduced pressure. The obtained residue was suspended and washed with n-hexane-ethyl acetate under heating at reflux to give the title compound(56mg, 77.3%) as a light yellow white crystal. mp 284-286°C.

¹H-NMR(DMSO-d₆): δ 3.84(2H, s), 6.98(1H, d, J=8.8Hz), 7.42(1H, d, J=6.8Hz), 7.49(2H, t, J=7.6Hz), 7.58-7.61(3H, m), 8.07(1H, d, J=2.8Hz), 12.25(H, brs).

Example 204: 5-Bromo-N-(4,5-diphenylthiazol-2-yl)-2-hydroxybenzamide (Comopund No. 204).

[0616] Using 5-bromosalicylic acid and 2-amino-4,5-diphenylthiazole as the raw materials, the same operation as the example 195(3) gave the title compound.

(2-Amino-4,5-diphenylthiazole: refer to Nihon Kagaku Zasshi, 1962, 83, 209.)

Yield: 25.9%.

mp 262-263°C.

¹H-NMR(DMSO-d₆): δ 7.02(1H, d, J=8.1Hz), 7.34-7.47(10H, m), 7.63(1H, d, J=6.9Hz), 8.08(1H, d, J=2.4Hz), 11.88(1H, brs), 12.08(1H, brs).

Example 205: 5-Bromo-N-(4-benzyl-5-phenylthiazol-2-yl)-2-hydroxybenzamide (Comopund No. 205).

[0617] Using 5-bromosalicylic acid and 2-amino-4-benzyl-5-phenylthiazole as the raw materials, the same operation as the example 195(3) gave the title compound.

(2-Amino-4-benzyl-5-phenylthiazole: refer to Chem. Pharm. Bull., 1962, 10, 376.)

Yield: 28.1%.

mp 198-200°C.

¹H-NMR(DMSO-d₆): δ 4.08(2H, s), 6.95(1H, d, J=8.8Hz), 7.15-7.22(3H, m), 7.30(2H, t, J=7.6Hz), 7.38-7.43(1H, m), 7.47(4H, d, J=4.4Hz), 7.57(1H, brd, J=8.8Hz), 8.05(1H, d, J=2.4Hz), 11.98(1H, brs).

Example 206: 5-Bromo-N-[5-phenyl-4-(trifluoromethyl)thiazol-2-yl]-2-hydroxybenzamide (Comopund No. 206).

[0618] Using 5-bromosalicylic acid and 2-amino-5-phenyl-4-(trifluoromethyl)thiazole as the raw materials, the same operation as the example 195(3) gave the title compound.

Yield: 33.2%.

mp 250°C(dec.).

¹H-NMR(DMSO-d₆): δ 7.02(1H, d, J=8.8Hz), 7.51(5H, s), 7.63(1H, dd, J=8.8, 2.4Hz), 8.02(1H, d, J=2.8Hz), 12.38(1H, brs).

Example 207: 5-Bromo-N-[5-acetyl-4-phenylthiazol-2-yl]-2-hydroxybenzamide (Comopund No. 207).

[0619] Using 1-phenyl-1,3-butanedione as the raw material, the same operation as the examples 195(1)-(3) gave the title compound.

Yield: 8.9%(3 steps).

EP 1 352 650 A1

(1) α -Bromo-1-phenyl-1,3-butanedione.

[0620] $^1\text{H-NMR}(\text{CDCl}_3)$: δ 2.46(3H, s), 5.62(1H, s), 7.48-7.54(2H, m), 7.64(1H, tt, $J=7.5, 2.1\text{Hz}$), 7.97-8.01(2H, m).

(2) 2-Amino-5-acetyl-4-phenylthiazole.

[0621] $^1\text{H-NMR}(\text{CDMSO}-d_6)$: δ 2.18(3H, s), 7.50-7.55(2H, m), 7.59-7.68(3H, m), 8.69(2H, brs).

(3) 5-Bromo-N-[5-acetyl-4-phenylthiazol-2-yl]-2-hydroxybenzamide.

[0622] $^1\text{H-NMR}(\text{DMSO}-d_6)$: δ 2.44(3H, s), 6.99(1H, d, $J=9.0\text{Hz}$), 7.55-7.71(4H, m), 7.76-7.80(2H, m), 8.01(1H, d, $J=2.4\text{Hz}$), 12.36(2H, br).

Example 208: 5-Bromo-N-[5-benzoyl-4-phenylthiazol-2-yl]-2-hydroxybenzamide (Comopund No. 208).

[0623] Using 1,3-diphenyl-1,3-propanedione as the raw material, the same operation as the examples 195(1)-(3) gave the title compound.

Yield: 49.7%.

(1) α -Bromo-1,3-diphenyl-1,3-propanedione.

[0624] $^1\text{H-NMR}(\text{CDCl}_3, \delta)$: 6.55(1H, s), 7.45-7.50(4H, m), 7.61(2H, tt, $J=7.2, 2.1\text{Hz}$), 7.98-8.01(4H, m).

(2) 2-Amino-5-benzoyl-4-phenylthiazole.

[0625] $^1\text{H-NMR}(\text{DMSO}-d_6)$: δ 7.04-7.18(5H, m), 7.22-7.32(3H, m), 7.35-7.38(2H, m), 8.02(2H, s).

(3) 5-Bromo-N-[5-benzoyl-4-phenylthiazol-2-yl]-2-hydroxybenzamide.

[0626] $^1\text{H-NMR}(\text{DMSO}-d_6)$: δ 7.03(1H, d, $J=8.7\text{Hz}$), 7.17-7.30(5H, m), 7.39-7.47(3H, m), 7.57-7.60(2H, m), 7.64(1H, dd, $J=8.7, 2.7\text{Hz}$), 8.05(1H, d, $J=2.4\text{Hz}$), 11.82(1H, brs), 12.35(1H, brs).

Example 209: 2-(5-Chloro-2-hydroxybenzoyl)amino-4-phenylthiazole-5-carboxylic acid ethyl ester (Comopund No. 209).

[0627] Using 5-chlorosalicylic acid and 2-amino-4-phenylthiazole-5-carboxylic acid ethyl ester as the raw materials, the same operation as the example 195(3) gave the title compound.

Yield: 69.4%.

$^1\text{H-NMR}(\text{DMSO}-d_6)$: δ 1.22(3H, t, $J=7.5\text{Hz}$), 4.21(2H, q, $J=7.5\text{Hz}$), 7.07(1H, d, $J=8.7\text{Hz}$), 7.43-7.47(3H, m), 7.53(1H, dd, $J=8.7, 2.4\text{Hz}$), 7.70-7.74(2H, m), 7.92(1H, d, $J=3.0\text{Hz}$), 11.88(1H, br), 12.29(1H, brs).

Example 210: 2-(5-Bromo-2-hydroxybenzoyl)amino-4-phenylthiazole-5-carboxylic acid ethyl ester (Comopund No. 210).

[0628] Using 5-bromosalicylic acid and 2-amino-4-phenylthiazole-5-carboxylic acid ethyl ester as the raw materials, the same operation as the example 195(3) gave the title compound.

Yield: 28.6%.

mp 197-199°C.

$^1\text{H-NMR}(\text{DMSO}-d_6)$: δ 1.21(3H, t, $J=6.8\text{Hz}$), 4.20(2H, q, $J=6.8\text{Hz}$), 7.01(1H, d, $J=8.8\text{Hz}$), 7.43-7.48(3H, m), 7.63(1H, dd, $J=8.8, 2.4\text{Hz}$), 7.70-7.72(2H, m), 8.04(1H, d, $J=2.4\text{Hz}$), 12.33(1H, brs).

Example 211: 2-(5-Bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)thiazole-5-carboxylic acid ethyl ester (Comopund No. 211).

[0629] Using pentafluorobenzoylacetic acid as the raw material, the same operation as the examples 195(1)-(3) gave the title compound.

Yield: 40.0%(3steps).

EP 1 352 650 A1

(1) α -Bromo-pentafluorobenzoylacetic acid ethyl ester.

[0630] It was used for the next reaction as a crude product.

(2) 2-Amino-4-(pentafluorophenyl)thiazole-5-carboxylic acid ethyl ester.

[0631] $^1\text{H-NMR}(\text{CDCl}_3)$: δ 1.23(3H, t, $J=7.2\text{Hz}$), 4.21(2H, q, $J=7.2\text{Hz}$), 5.41(2H, s).

(3) 2-(5-Bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)thiazole-5-carboxylic acid ethyl ester.

[0632] $^1\text{H-NMR}(\text{DMSO}-d_6)$: δ 1.20(3H, t, $J=7.2\text{Hz}$), 2.51(2H, q, $J=7.2\text{Hz}$), 7.02(1H, d, $J=8.7\text{Hz}$), 7.64(1H, dd, $J=8.7, 2.7\text{Hz}$), 7.90(1H, d, $J=3.0\text{Hz}$), 11.92(1H, br), 12.58(1H, br).

Example 212: [2-(5-Bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)thiazol-5-yl]-N-methylcarboxamide (Compound No. 212).

(1) 2-(5-Bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)thiazole-5-carboxylic acid.

[0633] Using 2-(5-bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)-thiazole-5-carboxylic acid ethyl ester as the raw material, the same operation as the example 82 gave the title compound.

(2) [2-(5-Bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)thiazol-5-yl]-N-methylcarboxamide.

[0634] A mixture of 2-(5-bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)-thiazole-5-carboxylic acid (0.20g, 0.48mmol), methylamine 40% methanol solution (0.2ml), 1-hydroxybenzotriazole hydrate (96.7mg, 0.72mmol), WSC HCl (137.2mg, 0.72mmol) and tetrahydrofuran (15mL) was stirred at room temperature for 18 hours. The reaction mixture was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the organic layer was washed with water and brine, dried over anhydrous sodium sulfate, the residue obtained by evaporation under reduced pressure was purified by chromatography on silica gel (n-hexane:ethyl acetate=1:2), and crystallized (dichloromethane/n-hexane) to give the title compound (87.9mg, 42.6%) as a white powder.

$^1\text{H-NMR}(\text{DMSO}-d_6)$: δ 2.70(3H, d, $J=4.5\text{Hz}$), 7.02(1H, d, $J=9.0\text{Hz}$), 7.40-7.48(3H, m), 7.63(1H, dd, $J=9.0, 2.4\text{Hz}$), 7.68-7.71(2H, m), 8.06(1H, d, $J=2.4\text{Hz}$), 8.16(1H, t, $J=4.5\text{Hz}$), 11.88(1H, br), 12.15(1H, brs).

Example 213:

[2-(5-Bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)thiazol-5-yl]-N-ethylcarboxamide (Compound No. 213).

[0635] Using 2-(5-bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)thiazole-5-carboxylic acid and 70% aqueous ethylamine solution as the raw materials, the same operation as the example 212(2) gave the title compound. Yield: 62.5%.

$^1\text{H-NMR}(\text{DMSO}-d_6)$: δ 1.05(3H, t, $J=6.9\text{Hz}$), 3.15-3.24(2H, m), 7.02(1H, d, $J=8.7\text{Hz}$), 7.40-7.47(3H, m), 7.63(1H, dd, $J=8.7, 3.0\text{Hz}$), 7.69-7.72(2H, m), 8.06(1H, d, $J=2.4\text{Hz}$), 8.20(1H, t, $J=5.4\text{Hz}$), 11.84(1H, br), 12.14(1H, brs).

Example 214: [2-(5-Bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)thiazol-5-yl]-N-isopropylcarboxamide (Compound No. 214).

[0636] Using 2-(5-bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)thiazole-5-carboxylic acid and isopropylamine as the raw materials, the same operation as the example 212(2) gave the title compound. Yield: 23.9%.

$^1\text{H-NMR}(\text{DMSO}-d_6)$: δ 1.07(6H, d, $J=6.3\text{Hz}$), 4.02(1H, m), 7.02(1H, d, $J=9.0\text{Hz}$), 7.40-7.52(3H, m), 7.64(1H, dd, $J=8.7, 2.7\text{Hz}$), 7.69-7.73(2H, m), 8.06(1H, d, $J=2.7\text{Hz}$), 11.89(1H, br), 12.14(1H, brs).

Example 215: [2-(5-Bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)thiazol-5-yl]-N-(2-phenethyl)carboxamide (Compound No. 215).

[0637] Using 2-(5-bromo-2-hydroxybenzoyl)amino-4-(pentafluorophenyl)thiazole-5-carboxylic acid and 2-phenethylamine as the raw materials, the same operation as the example 212 gave the title compound. Yield: 62.2%.

EP 1 352 650 A1

¹H-NMR(DMSO-d₆): δ 2.78(2H, t, J=7.5Hz), 3.43(2H, q, J=7.5Hz), 7.02(1H, d, J=9.0Hz), 7.19-7.24(3H, m), 7.27-7.33(2H, m), 7.39-7.41(3H, m), 7.61-7.65(3H, m), 8.06(1H, d, J=2.4Hz), 8.25(1H, t, J=6.0Hz), 11.85(1H, brs), 12.15(1H, brs).

Example 216: 2-(5-Bromo-2-hydroxybenzoyl)amino-4-(trifluoromethyl)thiazole-5-carboxylic acid ethyl ester
(Comopund No. 216).

[0638] Using 5-bromosalicylic acid and 2-amino-4-(trifluoromethyl)thiazole-5-carboxylic acid ethyl ester as the raw materials, the same operation as the example 195(3) gave the title compound.

Yield: 88.7%.

¹H-NMR(DMSO-d₆): δ 1.32(3H, t, J=7.2Hz), 4.33(2H, q, J=7.2Hz), 7.01(1H, d, J=8.7Hz), 7.63(1H, dd, J=8.7, 2.7Hz), 7.98(1H, d, J=2.4Hz), 12.64(1H, br).

Example 217: 2-Acetoxy-5-chloro-N-{4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl}benzamide.

[0639] Using 5-chloro-N-{4-(1,1-dimethyl)ethyl-5-[(2,2-dimethyl)propionyl]thiazol-2-yl}-2-hydroxybenzamide and acetyl chloride as the raw materials, the same operation as the example 96 gave the title compound.

Yield: 65.3%.

¹H-NMR(CDCl₃): δ 1.32(9H, s), 1.33(9H, s), 2.46(3H, s), 7.22(1H, d, J=8.4Hz), 7.56(1H, dd, J=8.7, 2.4Hz), 8.05(1H, d, J=2.7Hz), 9.82(1H, brs).

Example 218: 2-[(4-Hydroxybiphenyl)-3-carbonyl]amino-4-phenylthiazole-5-carboxylic acid ethyl ester (Comopund No. 218).

[0640] Using 4-hydroxybiphenyl-3-carboxylic acid and 2-amino-4-phenylthiazole-5-carboxylic acid ethyl ester as the raw materials, the same operation as the example 195(3) gave the title compound. (4-Hydroxybiphenyl-3-carboxylic acid: refer to Tetrahedron, 1997, 53, 11437.)

Yield: 61.7%.

mp 207-208°C.

¹H-NMR(DMSO-d₆): δ 1.23(3H, t, J=7.2Hz), 4.22(2H, q, J=7.2Hz), 7.16(1H, d, J=8.7Hz), 7.36(1H, t, J=7.5Hz), 7.45-7.50(5H, m), 7.69-7.76(4H, m), 7.85(1H, dd, J=8.7, 2.4Hz), 8.31(1H, d, J=2.4Hz), 11.73(1H, brs), 12.60(1H, brs).

Example 219: 2-[(4'-Fluoro-4-hydroxybiphenyl)-3-carbonyl]amino-4-phenylthiazole-5-carboxylic acid ethyl ester (Comopund No. 219).

[0641] Using (4'-fluoro-4-hydroxybiphenyl)-3-carboxylic acid and 2-amino-4-phenylthiazole-5-carboxylic acid ethyl ester as the raw materials, the same operation as the example 195(3) gave the title compound. ((4'-Fluoro-4-hydroxybiphenyl)-3-carboxylic acid: refer to Tetrahedron, 1997, 53, 11437.)

Yield: 62.7%.

mp 237-238°C.

¹H-NMR(DMSO-d₆): δ 1.22(3H, t, J=7.2Hz), 4.21(2H, q, J=7.2Hz), 7.13(1H, d, J=8.4Hz), 7.28(2H, t, J=8.8Hz), 7.44-7.45(3H, m), 7.71-7.75(4H, m), 7.81(1H, dd, J=8.8, 2.4Hz), 8.27(1H, d, J=2.4Hz), 11.67(1H, brs), 12.58(1H, brs).

Example 220: 2-[(2',4'-Difluoro-4-hydroxybiphenyl)-3-carbonyl]amino-4-phenylthiazole-5-carboxylic acid ethyl ester (Comopund No. 220).

[0642] Using (2',4'-difluoro-4-hydroxybiphenyl)-3-carboxylic acid and 2-amino-4-phenylthiazole-5-carboxylic acid ethyl ester as the raw materials, the same operation as the example 195(3) gave the title compound.

Yield: 45.6%.

mp 206-207°C.

¹H-NMR(DMSO-d₆): δ 1.22(3H, t, J=7.2Hz), 4.22(2H, q, J=7.2Hz), 7.17(1H, d, J=9.0Hz), 7.21(1H, td, J=8.7, 2.4Hz), 7.38(1H, ddd, J=11.7, 9.3, 2.4Hz), 7.44-7.46(3H, m), 7.60-7.75(4H, m), 8.13-8.14(1H, m), 11.86(1H, brs), 12.46(1H, brs).

Example 221: 2-([4-Hydroxy-4'-(trifluoromethyl)biphenyl]-3-carbonyl)amino-4-phenylthiazole-5-carboxylic acid ethyl ester (Comopund No. 221).

(1) [4'-(Trifluoromethyl)-4-hydroxybiphenyl]-3-carboxylic acid.

[0643] A mixture of 5-bromosalicylic acid(500mg, 2.30mmol), dihydroxy-4-(trifluoromethyl)phenylborane(488mg, 2.57mmol), palladium acetate(10mg 0.040mmol) and 1M sodium carbonate(7mL) was stirred at 80°C for 1 hour. The reaction mixture was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine one after another, dried over anhydrous sodium sulfate, the solvent was evaporated under reduced pressure. According to the fixed procedure, the obtained residue was methyl-esterified by trimethylsilyldiazomethane and methanol, and purified by column chromatography on silica gel(n-hexane:ethyl acetate=5:1) to give a colourless liquid(563mg). This liquid was dissolved in methanol(10mL). 2 N sodium hydroxide(3mL) was added, and the mixture was stirred at 60°C for 1 hour. After the reaction mixture was cooled to room temperature, it was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the ethyl acetate layer was washed with water and saturated brine one after another, dried over MgSO₄, the solvent was evaporated under reduced pressure. The obtained residue was suspended and washed with n-hexane-dichloromethane under heating at reflux to give the title compound (458mg, 70.4%) as a white crystal.

mp 185°C(dec).

¹H-NMR(DMSO-d₆): δ 7.09(1H, d, J=8.8Hz), 7.77(2H, d, J=8.0Hz), 7.85(2H, d, J=8.0Hz), 7.90(1H, dd, J=8.8, 2.0Hz), 8.10(1H, d, J=2.4Hz), 11.80(brs).

(2) 2-([4-Hydroxy-4'-(trifluoromethyl)biphenyl]-3-carbonyl)amino-4-phenylthiazole-5-carboxylic acid ethyl ester.

[0644] Using [4'-(trifluoromethyl)-4-hydroxybiphenyl]-3-carboxylic acid and 2-amino-4-phenylthiazole-5-carboxylic acid ethyl ester as the raw materials, the same operation as the example 195(3) gave the title compound.

Yield: 41.7%.

mp 236-237°C.

¹H-NMR(DMSO-d₆): δ 1.22(3H, t, J=7.2Hz), 4.21(2H, q, J=7.2Hz), 7.18(1H, d, J=8.8Hz), 7.44-7.45(3H, m), 7.72-7.74(2H, m), 7.81(2H, d, J=8.4Hz), 7.91(1H, dd, J=8.8, 2.4Hz), 7.93(2H, d, J=8.4Hz), 8.36(1H, d, J=2.4Hz), 11.78(1H, brs), 12.62(1H, brs).

Example 222: 2-[2-Hydroxy-5-(1-pyrrolyl)benzoyl] amino-4-phenylthiazole-5-carboxylic acid ethyl ester (Comopund No. 222).

[0645] Using 2-hydroxy-5-(1-pyrrolyl)benzoic acid and 2-amino-4-phenylthiazole-5-carboxylic acid ethyl ester as the raw materials, the same operation as the example 195(3) gave the title compound.

Yield: 55.0%.

¹H-NMR(DMSO-d₆): δ 1.22(3H, t, J=7.2Hz), 4.22(2H, q, J=7.2Hz), 6.26(2H, t, J=2.1Hz), 7.13(1H, d, J=8.7Hz), 7.32(2H, t, J=2.1Hz), 7.43-7.47(3H, m), 7.70-7.75(3H, m), 8.09(1H, d, J=2.7Hz), 11.58(1H, brs), 12.55(1H, brs).

Example 223: 2-[2-Hydroxy-5-(2-thienyl)benzoyl]amino-4-phenylthiazole-5-carboxylic acid ethyl ester (Comopund No. 223).

(1) 2-Hydroxy-5-(2-thienyl)benzoic acid.

[0646] 5-Bromosalicylic acid(500mg, 2.30mmol) was dissolved in 1,2-dimethoxyethane(5mL). Tetrakis(triphenylphosphine)palladium(80mg, 0.07mmol) was added under argon atmosphere, and the mixture was stirred at room temperature for 10 minutes. Then dihydroxy-2-thienylborane(324mg, 2.53mmol) and 1M sodium carbonate(7mL) were added, and the mixture was refluxed for 2 hours. After the reaction mixture was cooled to room temperature, it was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine one after another, dried over anhydrous sodium sulfate, the solvent was evaporated under reduced pressure. According to the fixed procedure, the obtained residue was methyl-esterified by trimethylsilyldiazomethane and methanol, and purified by column chromatography on silica gel(n-hexane:ethyl acetate=5:1) to give a yellow liquid(277mg). This was dissolved in methanol(5mL). 2 N sodium hydroxide(1.5mL) was added, and the mixture was stirred at 60°C for 1 hour. After the reaction mixture was cooled to room temperature, it was poured into 2 N hydrochloric acid and extracted with ethyl acetate. After the ethyl acetate layer was washed with water and brine one after another, dried over anhydrous magnesium sulfate, the residue obtained by evaporating the solvent under reduced pressure was crystallized from n-hexane-dichloromethane to give the title compound(58mg, 11.5%) as a white crystal.

EP 1 352 650 A1

¹H-NMR(DMSO-d₆): δ 6.95(1H, d, J=8.8Hz), 7.09(1H, dd, J=4.8, 3.6Hz), 7.37(1H, dd, J=4.0, 1.2Hz), 7.45(1H, dd, J=5.2, 1.2Hz), 7.74(1H, dd, J=8.8, 2.8Hz), 7.96(1H, d, J=2.8Hz).

(2) 2-[2-Hydroxy-5-(2-thienyl)benzoyl]amino-4-phenylthiazole-5-carboxylic acid ethyl ester.

[0647] Using 2-hydroxy-5-(2-thienyl)benzoic acid and 2-amino-4-phenylthiazole-5-carboxylic acid ethyl ester as the raw materials, the same operation as the example 195(3) gave the title compound.

Yield: 58.2%.

mp 213-2149°C.

¹H-NMR(DMSO-d₆): δ 1.22(3H, t, J=7.2Hz), 4.21(2H, q, J=7.2Hz), 7.10(1H, d, J=9.2Hz), 7.12(1H, dd, J=4.8, 3.6Hz), 7.44-7.46(4H, m), 7.50(1H, dd, J=4.8, 1.2Hz), 7.71-7.74(2H, m), 7.79(1H, dd, J=8.8, 2.4Hz), 8.21(1H, d, J=2.4Hz), 11.78(1H, brs), 12.44(1H, brs).

Test Example: Measurement of inhibitory activity of NF-κB activation.

[0648] Inhibitory activity of NF-κB activation was measured referring to the method of Hill et al. (Hill C. S., et al., Cell, 73, 395-406(1993)). Using a transfection reagent(Effectene; QIAGEN), the human hepatoma cell strain HepG2 or the human hystercarcinoma cell strain HeLa was transfected with the firefly luciferase gene(Luc) contained plasmid (pNF-κB-Luc Reporter Plasmid; STRATAGENE) which contained oligonucleotide having five tandem copies of NF-κB binding sequences(TGGGGACTTTCCGC) on a upstream region of Luc, according to the QUAGEN's protocol and it was incubated for 6-24 hours. After addition of TNF-α (40ng/ml) with or without the test compound, the cells were incubated for 4 hours, and intracellular luciferase activity was measured with PicaGene LT (TOYO INK MFG Co., Ltd.) and chemical luminescence measurement device (SPECTRAFluor Plus; TECAN). The inhibition ratio was measured as a ratio to the value of the luciferase activity without the test compound. The inhibition ratio of NF-κB activity with the test compound 10μg/ml or 1μg/ml were shown in the following table.

Compound Number	The rate of inhibitory activity against NF-κB activation (%)	
	Concentration of the agent: 10 μg/mL	Concentration of the agent: 1 μg/mL
1	54.4	-33.6
2	83.2	18.6
3	68.4	54.2
4	94.1	42.9
5	98.0	33.3
6	61.9	27.8
7	68.7	30.4
8	59.9	35.3
9	99.2	21.9
10	78.6	7.1
11	44.1	28.4
12	87.3	68.6
13	63.8	-7.1
14	98.9	21.7
15	70.4	15.2
16	91.6	36.4
17	96.5	19.9
18	90.2	85.3
19	95.1	-55.4
20	86.8	-12.1

EP 1 352 650 A1

(continued)

Compound Number	The rate of inhibitory activity against NF- κ B activation (%)	
	Concentration of the agent: 10 μ g/mL	Concentration of the agent: 1 μ g/mL
21	95.0	89.6
22	92.9	37.0
23	96.6	75.7
24	82.2	58.1
25	86.9	85.4
27	47.3	68.5
28	41.7	16.3
29	73.0	46.3
30	98.1	76.5
31	93.2	13.3
32	96.3	89.3
33	99.5	95.1
34	98.5	90.5
35	85.4	88.2
36	84.7	26.6
37	63.1	29.1
38	81.8	-10.1
39	56.0	21.4
40	81.9	3.9
41	90.3	26.1
42	92.3	14.3
43	78.9	25.5
44	65.8	36.7
45	91.3	61.7
46	85.7	-43.7
47	99.4	91.3
48	95.6	93.3
49	94.3	81.5
50	99.5	96.3
51	98.6	94.9
52	85.4	86.6
53	99.2	92.0
54	99.6	92.2
55	99.4	95.8
56	98.3	92.9
57	96.0	76.8
58	98.3	94.7

EP 1 352 650 A1

(continued)

Compound Number	The rate of inhibitory activity against NF- κ B activation (%)	
	Concentration of the agent: 10 μ g/mL	Concentration of the agent: 1 μ g/mL
59	99.2	94.5
60	99.4	42.7
61	98.5	59.7
62	99.1	74.9
63	96.9	95.5
64	90.1	53.3
65	97.1	83.9
66	94.9	91.1
67	96.8	91.8
68	98.3	92.3
69	99.6	96.4
70	95.4	93.3
71	97.9	93.8
72	97.8	79.5
73	92.9	81.7
74	95.3	82.1
76	99.0	90.4
77	97.0	30.7
78	99.2	86.3
79	98.7	90.7
81	96.4	88.2
82	94.5	-8.7
83	87.1	16.0
84	82.2	23.7
85	96.0	44.9
86	95.9	42.2
87	98.1	84.4
89	67.5	-21.6
90	63.4	1.0
91	88.4	20.5
92	97.2	51.8
93	98.7	96.2
94	89.1	19.4
95	97.1	90.9
96	99.2	96.5
97	96.0	69.9
98	98.2	90.5

EP 1 352 650 A1

(continued)

Compound Number	The rate of inhibitory activity against NF- κ B activation (%)	
	Concentration of the agent: 10 μ g/mL	Concentration of the agent: 1 μ g/mL
101	98.3	95.7
104	96.9	76.2
105	93.9	89.6
106	93.3	80.7
107	95.0	92.3
108	97.6	94.7
109	88.8	83.0
110	98.9	94.7
111	98.7	96.7
112	95.9	93.1
113	97.1	94.8
114	94.1	88.9
115	94.3	89.0
116	96.7	86.3
117	93.0	89.2
118	96.3	94.1
119	91.7	88.1
120	97.9	93.8
121	96.5	85.5
122	97.2	84.5
123	93.4	76.6
125	99.1	94.6
126	97.8	95.8
127	86.4	81.8
128	95.0	87.2
129	85.8	75.4
139	60.2	48.2
140	96.7	94.2
141	96.4	83.3
142	96.9	95.1
143	93.8	91.6
144	96.8	91.8
145	95.5	92.9
146	97.0	94.2
147	96.8	84.5
148	92.8	77.1
149	97.1	85.4

EP 1 352 650 A1

(continued)

Compound Number	The rate of inhibitory activity against NF- κ B activation (%)	
	Concentration of the agent: 10 μ g/mL	Concentration of the agent: 1 μ g/mL
150	95.1	91.4
151	71.8	-42.8
152	70.6	-56.8
153	88.7	49.1
154	48.2	-31.0
155	94.1	85.6
156	74.9	7.3
157	98.1	86.2
158	95.6	91.0
159	96.3	89.1
160	99.2	86.2
161	92.6	86.3
163	82.0	70.9
164	98.6	94.9
165	95.1	88.2
166	97.9	82.4
167	95.7	32.4
168	96.8	38.3
169	88.1	14.5
170	56.4	-40.0
171	95.8	33.7
172	97.5	88.6
180	42.8	-23.1
181	98.7	96.5
182	94.4	85.3
183	92.4	92.6
184	93.8	20.0
185	69.7	-1.5
186	95.2	88.4
187	67.2	4.6
188	94.4	83.6
189	82.0	-8.4
190	71.7	-32.4
191	98.1	90.5
192	87.6	28.8
193	96.1	70.1
194	88.7	46.1

EP 1 352 650 A1

(continued)


Compound Number	The rate of inhibitory activity against NF- κ B activation (%)	
	Concentration of the agent: 10 μ g/mL	Concentration of the agent: 1 μ g/mL
195	98.3	95.7
196	97.5	86.8
197	92.4	84.5
198	97.8	93.6
199	96.8	87.8
200	89.6	36.3
201	95.9	92.5
202	78.8	-41.8
203	72.1	2.4
204	67.0	-5.8
205	95.0	79.7
206	89.4	85.1
207	95.9	70.2
208	97.3	90.7
209	82.8	55.8
210	94.2	80.7
211	96.0	82.2
212	58.6	50.8
213	84.0	51.9
214	91.3	49.6
215	60.4	33.3
216	96.5	87.6
217	97.7	95.0
218	78.6	34.6
219	85.8	45.0
220	90.3	31.8
221	90.0	66.9
222	90.1	74.0
223	84.8	40.8

Industrial Applicability

[0649] The medicaments of the present invention have inhibitory activity against the activation of transcription factor NF- κ B, furthermore they have suppressing activity against the production and release of inflammatory cytokines. Therefore, the medicaments of the present invention are useful as preventive and/or therapeutic agent for treatment of diseases caused by activation of NF- κ B and for diseases caused by overproduction of inflammatory cytokines.

Claims



1. A medicament having inhibitory activity against NF- κ B which comprises as an active ingredient a substances selected from the group consisting of a compound represented by the following general formula (I) and a pharmaco-





wherein

- X represents a connecting group wherein a number of atoms in a main chain is 2 to 4 (said connecting group may be substituted),
 A represents hydrogen atom or acetyl group,
 E represents an aryl group which may be substituted or a heteroaryl group which may be substituted,
 ring Z represents an arene which may be substituted in addition to the group represented by formula -O-A wherein A has the same meaning as that defined above and the group represented by formula -X-E wherein each of X and E has the same meaning as that defined above, or ring Z represents a heteroarene which may be substituted in addition to the group represented by formula -O-A wherein A has the same meaning as that defined above and the group represented by formula -X-E wherein each of X and E has the same meaning as that defined above.
2. The medicament according to claim 1, wherein X is a group selected from the following connecting group α (said group may be substituted):


[Connecting Group α] The groups of the following formulas:





wherein a bond at the left end binds to ring Z and a bond at the right end binds to E.

3. The medicament according to claim 1, wherein X is a group represented by the following formula (said group may be substituted):



wherein a bond at the left end binds to ring Z and a bond at the right end binds to E.

4. The medicament according to any one of claims 1 to 3, wherein A is a hydrogen atom.
5. The medicament according to any one of claims 1 to 4, wherein ring Z is a C₆ to C₁₀ arene which may be substituted in addition to the group represented by the formula -O-A wherein A has the same meaning as that defined in the general formula (I) and the group represented by the formula -X-E wherein each of X and E has the same meaning as that defined in the general formula (I), or ring Z is a 6 to 13-membered heteroarene which may be substituted in addition to the group represented by the formula -O-A wherein A has the same meaning as that defined in the general formula (I) and the group represented by the formula -X-E wherein each of X and E has the same meaning as that defined in the general formula (I).
6. The medicament according to any one of claims 1 to 5, wherein ring Z is selected from the following ring group β:

[Ring Group β] benzene ring, naphthalene ring, pyridine ring, indole ring, quinoxaline ring, and carbazole ring

wherein said ring may be substituted in addition to the group represented by the formula -O-A wherein A has the same meaning as that defined in the general formula (I) and the group represented by the formula -X-E wherein each of X and E has the same meaning as that defined in the general formula (I).

7. The medicament according to claim 6, wherein ring Z is a benzene ring which may be substituted in addition to the group represented by the formula -O-A wherein A has the same meaning as that defined in the general formula (I) and the group represented by the formula -X-E wherein each of X and E has the same meaning as that defined in the general formula (I).
8. The medicament according to claim 7, wherein ring Z is a benzene ring which may be substituted with halogen atom(s) in addition to the group represented by the formula -O-A wherein A has the same meaning as that defined in the general formula (I) and the group represented by the formula -X-E wherein each of X and E has the same meaning as that defined in the general formula (I).

EP 1 352 650 A1

9. The medicament according to any one of claims 1 to 8, wherein E is a C₆ to C₁₀ aryl group which may be substituted or a 5 to 13 membered heteroaryl group which may be substituted.

10. The medicament according to claim 9, wherein E is a phenyl group which may be substituted.

11. The medicament according to claim 10, wherein E is 3,5-ditrifluoromethylphenyl group.


12. The medicament according to claim 8, wherein E is a 5 membered heteroaryl group which may be substituted.

13. The medicament according to any one of claims 1 to 12, which is an inhibitor against expression of a gene for one or more substances selected from the following substance group δ :

[Substance group δ] tumor necrosis factor (TNF), interleukin-1, interleukin-2, interleukin-6, interleukin-8, granulocyte colony-stimulating factor, interferon β , cell adhesion factor ICAM-1, VCAM-1, ELAM-1, nitric oxide synthetase, major histocompatibility antigen family class I, major histocompatibility antigen family class II, β 2-microglobulin, immunoglobulin light chain, serum amyloid A, angiotensinogen, complement B, complement C4, c-myc, transcript derived from HIV gene, transcript derived from HTLV gene, transcript derived from simian virus 40 gene, transcript derived from cytomegalovirus gene, and transcript derived from adenovirus gene.

14. The medicament according to any one of claims 1 to 12, which is an inhibitor against production and release of an inflammatory cytokine or an immuno suppressive agent.


15. A compound represented by the following general formula (I-1) or a salt thereof, or a hydrate thereof or a solvate thereof:



wherein Z¹ represents 2-hydroxyphenyl group which may be substituted in the 5-position or 2-acetoxyphenyl group which may be substituted in the 5-position, E¹ represents a phenyl group substituted with two C₁ to C₆ halogenated alkyl groups wherein said phenyl group may be substituted in addition to the two C₁ to C₆ halogenated alkyl groups, provided that the following compounds:

N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxybenzamide,
N-[3,5-bis(trifluoromethyl)phenyl]-5-chloro-2-hydroxybenzamide,
N-[3,5-bis(trifluoromethyl)phenyl]-5-bromo-2-hydroxybenzamide,
N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-iodobenzamide,
N-[3,5-bis(trifluoromethyl)phenyl]-2-hydroxy-5-nitrobenzamide, and
2-hydroxy-N-[2,3,5-tris(trifluoromethyl)phenyl]benzamide are excluded.

16. A compound represented by the following general formula (I-2) or a salt thereof, or a hydrate thereof or a solvate thereof




wherein Z² represents 2-hydroxyphenyl group which may be substituted in the 5-position or 2-acetoxyphenyl group


which may be substituted in the 5-position, E² represents a phenyl group whose 3-position or 5-position is substituted with a C₁ to C₆ halogenated alkyl group wherein said phenyl group may be substituted (except when a substituent is a C₁ to C₆ halogenated alkyl group) in addition to the C₁ to C₆ halogenated alkyl group in the 3-position or 5-position, provided that the following compounds:

5-chloro-2-hydroxy-N-[3-(trifluoromethyl)phenyl]benzamide,
 5-bromo-2-hydroxy-N-[3-(trifluoromethyl)phenyl]benzamide,
 5-hydroxy-5-iodo-N-[3-(trifluoromethyl)phenyl]benzamide,
 5-chloro-N-[4-chloro-3-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-chloro-N-[5-chloro-3-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-chloro-2-hydroxy-N-[4-nitro-3-(trifluoromethyl)phenyl]benzamide,
 5-fluoro-2-hydroxy-N-[2-(2,2,2-trifluoroethoxy)-5-(trifluoromethyl)phenyl]benzamide,
 5-fluoro-2-hydroxy-N-[2-(6,6,6-trifluorohexyloxy)-5-(trifluoromethyl)phenyl]benzamide,
 5-chloro-2-hydroxy-N-[3-trifluoromethyl-4-[[4-(trifluoromethyl)sulfanyl]phenoxy]phenyl]benzamide,
 N-[4-(benzothiazole-2-yl)sulfanyl-3-(trifluoromethyl)phenyl]-5-chloro-2-hydroxybenzamide,
 5-chloro-N-[2-(4-chlorophenoxy)-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-chloro-2-hydroxy-N-[2-(4-methylphenoxy)-5-(trifluoromethyl)phenyl]benzamide,
 5-chloro-N-[2-(4-chlorophenyl)sulfanyl-5-(trifluoromethyl)phenyl]-2-hydroxybenzamide,
 5-chloro-2-hydroxy-N-[2-(1-naphthyloxy)-5-(trifluoromethyl)phenyl]benzamide, and
 5-chloro-2-hydroxy-N-[2-(2-naphthyloxy)-5-(trifluoromethyl)phenyl]benzamide are excluded.

17. A compound represented by the following general formula (I-3) or a salt thereof, or a hydrate thereof or a solvate thereof




wherein Z³ represents 2-hydroxyphenyl group which may be substituted in the 5-position or 2-acetoxyphenyl group which may be substituted in the 5-position, E³ is a group represented by the following formula :




wherein R^{3e2} and R^{3e3} may be the same or different and each represents hydrogen atom, a hydrocarbon group which may be substituted, or hydroxyl group which may be substituted, provided that both of R^{3e2} and R^{3e3} are not simultaneously hydrogen atoms, and R^{3e5} represents a C₂ to C₆ hydrocarbon group which may be substituted.

18. A compound represented by the following general formula (I-4) or a salt thereof, or a hydrate thereof or a solvate thereof



10 wherein Z^4 represents 2-hydroxyphenyl group which may be substituted in the 5-position or 2-acetoxyphenyl group which may be substituted in the 5-position, E^4 is a group represented by the following formula :



20 wherein R^{4e4} represents a hydrocarbon group which may be substituted, R^{4e5} represents a halogen atom, cyano group, an acyl group which may be substituted, or a heterocyclic group which may be substituted.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP01/11084

A. CLASSIFICATION OF SUBJECT MATTER

(See extra sheet.)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
(See extra sheet.)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAPLUS (STN), REGISTRY (STN)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO, 99/65449, A2 (Smithkline Beecham Corp.), 23 December, 1999 (23.12.99), (Full text) & AU 9940255 A	1-18
X	Madan Babita et al., "2'-Hydroxychalcone inhibits nuclear factor- κ B and blocks tumor necrosis factor- α and lipopolysaccharide-induced adhesion of neutrophils to human umbilical vein endothelial cells", Mol. Pharmacol., Vol.58, No.3 (October 2000), pages 526 to 533 (Particularly, abstract, page 531)	1,2,4-10,13, 14
X A	EP, 221211, A1 (Unilever PLC.), 13 May, 1987 (13.05.87), (Family: none) (Particularly, page 6, lines 1, 3, 6)	16 1-15,17,18

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

<ul style="list-style-type: none"> * Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed 	<ul style="list-style-type: none"> "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family
---	--

Date of the actual completion of the international search
21 February, 2002 (21.02.02)Date of mailing of the international search report
12 March, 2002 (12.03.02)Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/JP01/11084

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X A	JP, 62-99329, A (Unilever N.V.), 08 May, 1987 (08.05.87), (Particularly, claim 7) & US 4560549 A & US 4725590 A & US 4742083 A	16 1-15, 17, 18
X A	US, 6117859, A (The Research Foundation of State University of New York), 12 September, 2000 (12.09.00), (Particularly, column 4, line 17 to column 6, line 19) & US 5958911 A	16 1-15, 17, 18
X	US, 4358443, A (The Research Foundation of State University of New York), 09 November, 1982 (09.11.82), (Particularly, column 10; Example 4; Table 1) & US 4287191 A & EP 38192 A1 & JP 57-112360 A	16
A	Naoki MATSUMOTO et al., "Synthesis of NK-kB Activation Inhibitors Derived from Epoxyquinomicin C", Bioorg. Med. Chemical Lett., Vol.10, No.9 (May 2000), Pages 865 to 869	1-18

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP01/11084

Continuation of A. CLASSIFICATION OF SUBJECT MATTER
(International Patent Classification (IPC))

Int.Cl⁷ A61K31/055, 31/166, 31/12, 31/18, 31/167, 31/136, 31/17, 31/695, 31/5375, 31/357, 31/404, 31/44, 31/498, 31/403, 31/415, 31/421, 31/422, 31/433, 31/428, 31/505, 31/47, 31/40, 31/381, 31/426, 31/437, 31/4402, 31/4453, 31/445, C07D295/192, 321/10, 215/12, 213/82, 209/42, 241/44, 209/88, 333/68, 231/40, 263/48, 413/14, 285/12, 213/75, 239/47, 209/08, 209/42, 215/38, 209/88, 207/325, 333/24, 277/30, 471/04, 213/56, 295/192, 211/16, 207/48, 213/76, 277/46, 277/56, 413/14, C07C235/60, 233/75, A61P43/00, 19/02, 29/00, 37/06, 17/00, 21/00, 21/04, 9/00, 1/04, 1/16, 13/12, 1/18, 19/06, 9/10, 17/06, 37/08, 11/00, 3/04, 3/10, 3/06, 27/02, 25/28, 35/00, 35/02, 15/00
(According to International Patent Classification (IPC) or to both national classification and IPC)

Continuation of B. FIELDS SEARCHED
Minimum Documentation searched (International Patent Classification (IPC))

Int.Cl⁷ A61K31/055, 31/166, 31/12, 31/18, 31/167, 31/136, 31/17, 31/695, 31/5375, 31/357, 31/404, 31/44, 31/498, 31/403, 31/415, 31/421, 31/422, 31/433, 31/428, 31/505, 31/47, 31/40, 31/381, 31/426, 31/437, 31/4402, 31/4453, 31/445, C07D295/192, 321/10, 215/12, 213/82, 209/42, 241/44, 209/88, 333/68, 231/40, 263/48, 413/14, 285/12, 213/75, 239/47, 209/08, 209/42, 215/38, 209/88, 207/325, 333/24, 277/30, 471/04, 213/56, 295/192, 211/16, 207/48, 213/76, 277/46, 277/56, 413/14, C07C235/60, 233/75, A61P43/00, 19/02, 29/00, 37/06, 17/00, 21/00, 21/04, 9/00, 1/04, 1/16, 13/12, 1/18, 19/06, 9/10, 17/06, 37/08, 11/00, 3/04, 3/10, 3/06, 27/02, 25/28, 35/00, 35/02, 15/00
Minimum Documentation Searched(classification system followed by classification symbols)

REMARK CONCERNING THE SUBJECT OF SEARCH:

Each of claims 1-14 relates to medicines containing an extraordinary variety of compounds as the active ingredient.

However, only a small part of the compounds represented by the general formula (I) are supported by the description within the meaning of Article 6 of the PCT and disclosed within the meaning of Article 5 of the PCT.

Therefore, this search has been carried out with priority given to the part supported by the description and disclosed therein, that is, to Examples.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.